

References of systematic review of the relationship between SHS and breastfeeding

Those references with an asterisk are included in the systematic review.

National Institute on Drug Abuse. (2012). Tobacco/Nicotine. *Research Report Series*.

Retrieved from <http://www.who.int/mediacentre/factsheets/fs339/en/>

*Baheiraei, A., Ghafoori, F., Rahimi Foroushani, A., & Nedjat, S. (2014). The effects of maternal exposure to second-hand smoke on breast-feeding duration: A prospective cohort study. *Journal of Public Health (Germany)*, 22(1), 13-22.

Baron, J. A., Bulbrook, R. D., Wang, D. Y., & Kwa, H. G. (1986). Cigarette smoking and prolactin in women. *British Medical Journal (Clinical research ed.)*, 293(6545), 482-483.

Ben-Jonathan, N., Hugo, E. R., Brandebourg, T. D., & LaPensee, C. R. (2006). Focus on prolactin as a metabolic hormone. *Trends in Endocrinology & Metabolism*, 17(3), 110-116. doi:<https://doi.org/10.1016/j.tem.2006.02.005>

Benkaddour, Y. A., Fatih, B., Majdi, F., & Soummami, A. (2016). Passive Smoking and Other Principal Risk Factors Associated with Low Birth Weight. *Open Journal of Obstetrics and Gynecology*, 6(07), 390.

Blake, C. A., & Sawyer, C. H. (1972). Nicotine Blocks the Suckling-Induced Rise in Circulating Prolactin in Lactating Rats. *Science*, 177, 619-621.

Casetta, B., Videla, A. J., Bardach, A., Morello, P., Soto, N., Lee, K., . . . Ciapponi, A. (2017). Association Between Cigarette Smoking Prevalence and Income Level: A Systematic Review and Meta-Analysis. *Nicotine & Tobacco Research*, 19(12), 1401-1407. doi:10.1093/ntr/ntw266

- *Chou, S. Y., Hsu, H. H., Kuo, H. H., & Kuo, H. W. (2008). Association between exposure to environmental tobacco smoke (ETS) and breastfeeding behaviour. *Acta Paediatrica*, 97(1), 76-80.
- Daly, J. B., Mackenzie, L. J., Freund, M., Wolfenden, L., Roseby, R., & Wiggers, J. H. (2016). Interventions by health care professionals who provide routine child health care to reduce tobacco smoke exposure in children: A review and meta-analysis. *JAMA Pediatrics*, 170(2), 138-147.
doi:10.1001/jamapediatrics.2015.3342
- Dorea, J. G. (2007). Maternal smoking and infant feeding: breastfeeding is better and safer. *Maternal & Child Health Journal*, 11(3), 287-291.
- Eidelman, A. I., Schanler, R. J., Johnston, M., Landers, S., Noble, L., Szucs, K., & Viehmann, L. (2012). Breastfeeding and the use of human milk. *Pediatrics*, 129(3), e827-e841.
- Farber, H. J., Nelson, K. E., Groner, J. A., & Walley, S. C. (2015). Public Policy to Protect Children From Tobacco, Nicotine, and Tobacco Smoke. *Pediatrics*, 136(5), 998-1007. doi:10.1542/peds.2015-3109
- Hawsawi, A. M., Bryant, L. O., & Goodfellow, L. T. (2015). Association between exposure to secondhand smoke during pregnancy and low birthweight: a narrative review. *Respiratory Care*, 60(1), 135-140.
- Horta, B. L., Loret de Mola, C., & Victora, C. G. (2015). Long-term consequences of breastfeeding on cholesterol, obesity, systolic blood pressure and type 2 diabetes: a systematic review and meta-analysis. *Acta Paediatrica*, 104, 30-37.

doi:10.1111/apa.13133

- *Horta, B. L., Victora, C. G., Menezes, A. M., & Barros, F. C. (1997). Environmental tobacco smoke and breastfeeding duration. *American Journal of Epidemiology*, 146(2), 128-133.
- Jaddoe, V. W., Troe, E. J., Hofman, A., Mackenbach, J. P., Moll, H. A., Steegers, E. A., & Witteman, J. C. (2008). Active and passive maternal smoking during pregnancy and the risks of low birthweight and preterm birth: the Generation R Study. *Paediatric Perinatal Epidemiology*, 22(2), 162-171.
- *Jedrychowski, W., Perera, F., Mroz, E., Edwards, S., Flak, E., Rauh, V., . . . Musial, A. (2008). Prenatal exposure to passive smoking and duration of breastfeeding in nonsmoking women: Krakow inner city prospective cohort study. *Archives of Gynecology & Obstetrics*, 278(5), 411-417.
- Khader, Y. S., Al-Akour, N., Alzubi, I. M., & Lataifeh, I. (2011). The association between second hand smoke and low birth weight and preterm delivery. *Maternal Child Health Journal*, 15(4), 453-459. doi:10.1007/s10995-010-0599-2
- Kharrazi, M., DeLorenze, G. N., Kaufman, F. L., Eskenazi, B., Bernert, J. T., Jr., Graham, S., . . . Pirkle, J. (2004). Environmental tobacco smoke and pregnancy outcome. *Epidemiology*, 15(6), 660-670.
- Kim, S. Y., Park, J. E., Lee, Y. J., Seo, H.-J., Sheen, S.-S., Hahn, S., . . . Son, H.-J. (2011). Testing a tool for assessing the risk of bias for nonrandomized studies showed moderate reliability and promising validity. *Journal of Clinical Epidemiology*, 64(10), 1075-1082.

66(4), 408-414. doi:10.1016/j.jclinepi.2012.09.016

*Kwok, M. K., Schooling, C. M., Lam, T. H., & Leung, G. M. (2010). Paternal smoking and childhood overweight: evidence from the Hong Kong "Children of 1997". *Pediatrics*, 126(1), e46-56.

*Lemke, M., Hartert, T. V., Gebretsadik, T., & Carroll, K. N. (2013). Relationship of secondhand smoke and infant lower respiratory tract infection severity by familial atopy status. *Annals of Allergy, Asthma, & Immunology*, 110(6), 433-437.

Leonardi-Bee, J., Britton, J., & Venn, A. (2011). Secondhand smoke and adverse fetal outcomes in nonsmoking pregnant women: a meta-analysis. *Pediatrics*, 127(4), 734-741. doi:10.1542/peds.2010-3041

Leonardi-Bee, J., Smyth, A., Britton, J., & Coleman, T. (2008). Environmental tobacco smoke and fetal health: systematic review and meta-analysis. *Archives Diseases Children Fetal Neonatal Education*, 93(5), F351-361.
doi:10.1136/adc.2007.133553

Lindbohm, M.-L., Sallmén, M., & Taskinen, H. (2002). Effects of exposure to environmental tobacco smoke on reproductive health. *Scandinavian Journal Work Environment Health*, 28, 84-96.

Mennella, J. A., Yourshaw, L. M., & Morgan, L. K. (2007). Breastfeeding and Smoking: Short-term Effects on Infant Feeding and Sleep. *Pediatrics*, 120(3), 497-502. doi:10.1542/peds.2007-0488

Minh, H. V., Ngan, T. T., Mai, V. Q., My, N. T., Chung le, H., Kien, V. D., . . . Giang,

- K. B. (2016). Tobacco Control Policies in Vietnam: Review on MPOWER Implementation Progress and Challenges. *Asian Pacific Journal Cancer Prevention, 17*(S1), 1-9.
- Minichiello, A., Lefkowitz, A. R., Firestone, M., Smylie, J. K., & Schwartz, R. (2016). Effective strategies to reduce commercial tobacco use in Indigenous communities globally: A systematic review. *BMC Public Health, 16*, 21. doi:10.1186/s12889-015-2645-x
- Rosen-Carole, C. B., Auinger, P., Howard, C. R., Brownell, E. A., & Lanphear, B. P. (2017). Low-Level Prenatal Toxin Exposures and Breastfeeding Duration: A Prospective Cohort Study. *Maternal Child Health Journal*. doi:10.1007/s10995-017-2346-4
- *Salama, R. H. M., Abdel-Aal, D. E. D. M., Eshra, D. K., Nagieb, S., & Arief, A. F. (2013). Clinical and biochemical effects of environmental tobacco smoking on pregnancy outcome. *Indian Journal of Clinical Biochemistry, 28*(4), 368-373.
- U. S Department of Health & Human Services. BeTobaccoFree.gov.
Retrieved from <https://betobaccofree.hhs.gov/laws/index.html>
- Sopori, M. (2002). Effects of cigarette smoke on the immune system. *Nature Reviews Immunology, 2*(5), 372-377.
- Thakur, S. K., Roy, S. K., Paul, K., Khanam, M., Khatun, W., & Sarker, D. (2012). Effect of nutrition education on exclusive breastfeeding for nutritional outcome of low birth weight babies. *European Journal Clinical Nutrition, 66*(3), 376-381. doi:10.1038/ejcn.2011.182

Department of Health. Tobacco Control Office, The Government of the Hong Kong

Special Administrative Region. (2012). Tobacco Control Legislation.

Retrieved from

http://gamapserver.who.int/gho/interactive_charts/tobacco/use/atlas.html

Vila Candel, R., Soriano-Vidal, F. J., Hevilla Cucarella, E., Castro-Sanchez, E., & Martin-Moreno, J. M. (2015a). Tobacco use in the third trimester of pregnancy and its relationship to birth weight. A prospective study in Spain. *Women & Birth*, 28(4), e134-139. doi:10.1016/j.wombi.2015.06.003

Vila Candel, R., Soriano-Vidal, F. J., Hevilla Cucarella, E., Castro-Sanchez, E., & Martin-Moreno, J. M. (2015b). Tobacco use in the third trimester of pregnancy and its relationship to birth weight. A prospective study in Spain. *Women & Birth*, 28(4), e134-139.

*Wdowiak, A., Wiktor, H., & Wdowiak, L. (2009). Maternal passive smoking during pregnancy and neonatal health. *Annals Agricultural Environmental Medicine*, 16(2), 309-312.

World Health Organization. (2002). *Global strategy on Infant and young child feeding*.

Retrieved from

http://apps.who.int/gb/archive/pdf_files/WHA55/ea5515.pdf?ua=1

World Health Organization. (2016). *WHO media center tobacco fact sheet*.

Retrieved from <http://www.who.int/mediacentre/factsheets/fs339/en/>

World Health Organization. (2016). *Prevalence of tobacco smoking*. Retrieved from

http://gamapserver.who.int/gho/interactive_charts/tobacco/use/atlas.html

World Health Organization. (2017). *WHO report on the global tobacco epidemic, 2017: monitoring tobacco use and prevention policies*. Retrieved from <http://apps.who.int/iris/bitstream/10665/255874/1/9789241512824-eng.pdf?ua=1&ua=1>

Zhou, S., Rosenthal, D. G., Sherman, S., Zelikoff, J., Gordon, T., & Weitzman, M. (2014). Physical, behavioral, and cognitive effects of prenatal tobacco and postnatal secondhand smoke exposure. *Current Problems in Pediatric & Adolescent Health Care*, 44(8), 219-241.

Reference systematic review of the relationship between SHS and depressive symptoms

Those references with an asterisk are included in the systematic review.

- *Alibekova, R., Huang, J. P., Lee, T. S. H., Au, H. K., & Chen, Y. H. (2016). Effects of smoking on perinatal depression and anxiety in mothers and fathers: A prospective cohort study. *Journal of Affective Disorders*, 193, 18-26.
- Bandiera, F. C., Arheart, K. L., Caban-Martinez, A. J., Fleming, L. E., McCollister, K., Dietz, N. A., . . . Lee, D. J. (2010). Secondhand Smoke Exposure and Depressive Symptoms. *Psychosomatic Medicine*, 72(1), 68-72.
doi:10.1097/PSY.0b013e3181c6c8b5
- Chen, J., Wang, M.-P., Wang, X., Viswanath, K., Lam, T.-H., & Chan, S. S. (2015). Secondhand smoke exposure (SHS) and health-related quality of life (HRQoL) in Chinese never smokers in Hong Kong. *BMJ Open*, 5(9), e007694.
doi:10.1136/bmjopen-2015-007694
- Gim, W., Yoo, J.-H., Shin, J.-Y., & Goo, A.-J. (2016). Relationship between Secondhand Smoking with Depressive Symptom and Suicidal Ideation in Korean Non-Smoker Adults: The Korean National Health and Nutrition Examination Survey 2010–2012. *Korean Journal of Family Medicine*, 37(2), 97-104. doi:10.4082/kjfm.2016.37.2.97
- Healey, C., Morriss, R., Henshaw, C., Wadoo, O., Sajjad, A., Scholefield, H., & Kinderman, P. (2013). Self-harm in postpartum depression and referrals to a perinatal mental health team: an audit study. *Archives of women's mental health*,

- 16(3), 237-245. doi:10.1007/s00737-013-0335-1
- Hiscock, R., Bauld, L., Amos, A., Fidler, J. A., & Munafò, M. (2012). Socioeconomic status and smoking: a review. *Annals of the New York Academy of Sciences*, 1248(1), 107-123. doi:10.1111/j.1749-6632.2011.06202.x
- Hoebel, J., Maske, U. E., Zeeb, H., & Lampert, T. (2017). Social Inequalities and Depressive Symptoms in Adults: The Role of Objective and Subjective Socioeconomic Status. *PLoS One*, 12(1), e0169764. doi:10.1371/journal.pone.0169764
- Hughes, J. R. (2008). Smoking and Suicide: A Brief Overview. *Drug and alcohol dependence*, 98(3), 169-178. doi:10.1016/j.drugalcdep.2008.06.003
- Jung, S. J., Shin, A., & Kang, D. (2015). Active smoking and exposure to secondhand smoke and their relationship to depressive symptoms in the Korea national health and nutrition examination survey (KNHANES). *BMC Public Health*, 15, 1053. doi:10.1186/s12889-015-2402-1
- *Khan, S., Arif, A. A., Laditka, J. N., & Racine, E. F. (2015). Prenatal exposure to secondhand smoke may increase the risk of postpartum depressive symptoms. *Journal of Public Health*, 37(3), 406-411.
- Kim, S. Y., Park, J. E., Lee, Y. J., Seo, H.-J., Sheen, S.-S., Hahn, S., . . . Son, H.-J. Testing a tool for assessing the risk of bias for nonrandomized studies showed moderate reliability and promising validity. *Journal of Clinical Epidemiology*, 66(4), 408-414. doi:10.1016/j.jclinepi.2012.09.016
- Laaksonen, M., Rahkonen, O., Karvonen, S., & Lahelma, E. (2005). Socioeconomic

- status and smokingAnalysing inequalities with multiple indicators. *European journal of public health*, 15(3), 262-269. doi:10.1093/eurpub/cki115
- Lisa C. Dierker, Shelli Avenevoli, Marilyn Stolar, & K.R. Merikangas. (2002). Smoking and Depression: An Examination of Mechanisms of Comorbidity. *American Journal of Psychiatry*, 159(6), 947-953. doi:10.1176/appi.ajp.159.6.947
- LORANT, V., CROUX, C., WEICH, S., DELIÈGE, D., MACKENBACH, J., & ANSSEAU, M. (2007). Depression and socio-economic risk factors: 7-year longitudinal population study. *The British Journal of Psychiatry*, 190(4), 293-298. doi:10.1192/bjp.bp.105.020040
- *Mbah, A. K., Salihu, H. M., Dagne, G., Wilson, R. E., & Bruder, K. (2013). Exposure to environmental tobacco smoke and risk of antenatal depression: application of latent variable modeling. *Archives of women's mental health*, 16(4), 293-302.
- *Miyake, Y., Tanaka, K., & Arakawa, M. (2012). Employment, income, and education and prevalence of depressive symptoms during pregnancy: the Kyushu Okinawa Maternal and Child Health Study. *BMC psychiatry*, 12, 117.
- Mojtabai, R., & Crum, R. M. (2013). Cigarette Smoking and Onset of Mood and Anxiety Disorders. *American journal of public health*, 103(9), 1656-1665. doi:10.2105/AJPH.2012.300911
- Pierce, J. P., White, V. M., & Emery, S. L. (2012). What public health strategies are needed to reduce smoking initiation? *Tobacco control*, 21(2), 258-264. doi:10.1136/tobaccocontrol-2011-050359

- Rhoades, R. R., & Beebe, L. A. (2015). Tobacco Control and Prevention in Oklahoma: Best Practices in a Preemptive State. *American journal of preventive medicine*, 48(1, Supplement 1), S6-S12. doi:<https://doi.org/10.1016/j.amepre.2014.09.001>
- Salimi, S., Terplan, M., Cheng, D., & Chisolm, M. S. (2015). The Relationship Between Postpartum Depression and Perinatal Cigarette Smoking: An Analysis of PRAMS Data. *Journal of substance abuse treatment*, 56, 34-38. doi:10.1016/j.jsat.2015.03.004
- Satoh, A., Kitamiya, C., Yaegashi, Y., Ohse, F., Lee, S., Kishi, C., . . . Sasaki, H. (2013). Influence of smoking on postpartum depression in Japan. *Health*, 05(08), 1254-1260. doi:10.4236/health.2013.58170
- Sit, D., Luther, J., Buysse, D., Dills, J. L., Eng, H., Okun, M., . . . Wisner, K. L. (2015). Suicidal Ideation in Depressed Postpartum Women: Associations with Childhood Trauma, Sleep Disturbance and Anxiety. *Journal of psychiatric research*, 0, 95-104. doi:10.1016/j.jpsychires.2015.04.021
- Stephoe, A., ardle, J., Tsuda, A., & Tanaka, Y. (2007). Depressive symptoms, socio-economic background, sense of control, and cultural factors in University students from 23 Countries. *International Journal of Behavioral Medicine*, 14(2), 97-107. doi:10.1007/bf03004175
- Swanson, E., Scott, E., & Thomas, M. (2017). Smoking and its effects on postpartum depression. *Advances in Pharmacy: Journal of Student Solutions to Pharmacy Challenges*, 1(1).
- *Tan, S., Courtney, L. P., El-Mohandes, A. A., Gantz, M. G., Blake, S. M., Thornberry,

- J., . . . Kiely, M. (2011). Relationships between self-reported smoking, household environmental tobacco smoke exposure and depressive symptoms in a pregnant minority population. *Maternal & Child Health Journal, 15 Suppl 1*, S65-74.
- Taylor, G., McNeill, A., Girling, A., Farley, A., Lindson-Hawley, N., & Aveyard, P. (2014). Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ : British Medical Journal, 348*. doi:10.1136/bmj.g1151
- United Nations, (2015). Transforming our World: The 2030 Agenda for Sustainable Development.
- *Vivilaki, V. G., Diamanti, A., Tzeli, M., Patelarou, E., Bick, D., Papadakis, S., . . . Katsaounou, P. (2016). Exposure to active and passive smoking among Greek pregnant women. *Tobacco induced diseases, 14*, 12.
- Votruba, N., Thornicroft, G., & the FundaMental, S. D. G. S. G. (2016). Sustainable development goals and mental health: learnings from the contribution of the FundaMentalSDG global initiative. *Global Mental Health, 3*, e26. doi:10.1017/gmh.2016.20
- Wagenknecht, L. E., Perkins, L. L., Cutter, G. R., Sidney, S., Burke, G. L., Manolio, T. A., . . . Hulley, S. B. (1990). Cigarette smoking behavior is strongly related to educational status: The cardia study. *Preventive medicine, 19*(2), 158-169. doi:https://doi.org/10.1016/0091-7435(90)90017-E
- *Weng, S. C., Huang, J. P., Huang, Y. L., Lee, T. S., & Chen, Y. H. (2016). Effects of tobacco exposure on perinatal suicidal ideation, depression, and anxiety. *BMC*

Public Health, 16, 623. doi:10.1186/s12889-016-3254-z

World Health Organization, Maternal mental health. *Mental health*. Retrieved from

http://www.who.int/mental_health/maternal-child/maternal_mental_health/en/

World Health Organization, (2013a). Mental health action plan 2013 - 2020

World Health Organization, (2013b). WHO recommendations for the prevention and management of tobacco use and second-hand smoke exposure in pregnancy.

World Health Organization, (2016). Prevalence of tobacco smoking. Retrieved from

http://gamapsserver.who.int/gho/interactive_charts/tobacco/use/atlas.html

World Health Organization, (2017a). Depression fact sheet. *WHO Media center*.

Retrieved from <http://www.who.int/mediacentre/factsheets/fs369/en/>

World Health Organization, (2017b). Suicide data. *Mental health*. Retrieved from

http://www.who.int/mental_health/prevention/suicide/suicideprevent/en/

Reference of this study

- Alibekova, R., Huang, J. P., Lee, T. S. H., Au, H. K., & Chen, Y. H. (2016). Effects of smoking on perinatal depression and anxiety in mothers and fathers: A prospective cohort study. *Journal of Affective Disorders, 193*, 18-26.
- Baheiraei, A., Ghafoori, F., Rahimi Foroushani, A., & Nedjat, S. (2014). The effects of maternal exposure to second-hand smoke on breast-feeding duration: A prospective cohort study. *Journal of Public Health (Germany), 22*(1), 13-22.
- Benkaddour, Y. A., Fatih, B., Majdi, F., & Soummami, A. (2016). Passive smoking and other principal risk factors associated with low birth weight. *Open Journal of Obstetrics and Gynecology, 6*(07), 390.
- Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. (2004). [WHO Expert Consultation] *The Lancet, 363*(9403), 157-163. doi:[https://doi.org/10.1016/S0140-6736\(03\)15268-3](https://doi.org/10.1016/S0140-6736(03)15268-3)
- Department of Health, Government of Western Australia. (2006). *Edinburgh Postnatal Depression Scale (EPDS): Translated versions – validated*. (Perth, Western Australia: State Perinatal Mental Health Reference Group.) Retrieved from <https://www.mcpapformoms.org/Docs/Edinburgh%20Depression%20Scale%20Translated%20Government%20of%20Western%20Australia%20Department%20of%20Health.pdf>
- United Nations Statistic Divisions. (2018). UN data profile Indonesia. Retrieved from <http://data.un.org/CountryProfile.aspx?crName=indonesia>

- Dubray, J., Schwartz, R., Chaiton, M., O'Connor, S., & Cohen, J. E. (2015). The effect of MPOWER on smoking prevalence. *Tobacco Control*, 24(6), 540-542.
doi:10.1136/tobaccocontrol-2014-051834
- Hawsawi, A. M., Bryant, L. O., & Goodfellow, L. T. (2015). Association between exposure to secondhand smoke during pregnancy and low birthweight: A narrative review. *Respiratory Care*, 60(1), 135-140.
- Hillenbrand, K. M., & Larsen, P. G. (2002). Effect of an educational intervention about breastfeeding on the knowledge, confidence, and behaviors of pediatric resident physicians. *Pediatrics*, 110(5), e59-e59. doi:10.1542/peds.110.5.e59
- Hiscock, R., Bauld, L., Amos, A., Fidler, J. A., & Munafò, M. (2012). Socioeconomic status and smoking: A review. *Annals of the New York Academy of Sciences*, 1248(1), 107-123. doi:10.1111/j.1749-6632.2011.06202.x
- Hughes, J. R. (2008). Smoking and suicide: A brief overview. *Drug Alcohol Dependency*, 98(3), 169-178. doi:10.1016/j.drugalcdep.2008.06.003
- Jacobs, M., Alonso, A. M., Sherin, K. M., Koh, Y., Dhamija, A., & Lowe, A. L. (2013). Policies to restrict secondhand smoke exposure: American College of Preventive Medicine Position Statement. *American Journal Preventive Medicine*, 45(3), 360-367. doi:10.1016/j.amepre.2013.05.007
- Jaddoe, V. W., Troe, E. J., Hofman, A., Mackenbach, J. P., Moll, H. A., Steegers, E. A., & Witteman, J. C. (2008). Active and passive maternal smoking during pregnancy and the risks of low birthweight and preterm birth: The generation R Study. *Paediatric Perinatal Epidemiology*, 22(2), 162-171. doi:10.1111/j.1365-

3016.2007.00916.x

Kaufman, M. R., Merritt, A. P., Rimbatmaja, R., & Cohen, J. E. (2015). 'Excuse me, sir. Please don't smoke here'. A qualitative study of social enforcement of smoke-free policies in Indonesia. *Health Policy Planning*, 30(8), 995-1002.
doi:10.1093/heapol/czu103

Khader, Y. S., Al-Akour, N., Alzubi, I. M., & Lataifeh, I. (2011). The association between second hand smoke and low birth weight and preterm delivery. *Maternal Child Health Journal*, 15(4), 453-459. doi:10.1007/s10995-010-0599-2

Laaksonen, M., Rahkonen, O., Karvonen, S., & Lahelma, E. (2005). Socioeconomic status and smoking: Analysing inequalities with multiple indicators. *European Journal Public Health*, 15(3), 262-269. doi:10.1093/eurpub/cki115

Leonardi-Bee, J., Britton, J., & Venn, A. (2011). Secondhand smoke and adverse fetal outcomes in nonsmoking pregnant women: A meta-analysis. *Pediatrics*, 127(4), 734-741. doi:10.1542/peds.2010-3041

Levy, D. T., Yuan, Z., Luo, Y., & Mays, D. (2016). Seven years of progress in tobacco control: An evaluation of the effect of nations meeting the highest level MPOWER measures between 2007 and 2014. *Tobacco Control*.
doi:10.1136/tobaccocontrol-2016-053381

Markou, A., Kosten, T. R., & Koob, G. F. (1998). Neurobiological similarities in depression and drug dependence: A self-medication hypothesis. *Neuropsychopharmacology*, 18, 135. doi:10.1016/S0893-133X(97)00113-9

- Naeye, R. L. (1987). Do placental weights have clinical significance? *Human Pathology*, 18(4), 387-391. doi:[https://doi.org/10.1016/S0046-8177\(87\)80170-3](https://doi.org/10.1016/S0046-8177(87)80170-3)
- Ngo, A., Cheng, K.-W., Chaloupka, F. J., & Shang, C. (2017). The effect of MPOWER scores on cigarette smoking prevalence and consumption. *Preventive Medicine*, 105(Supplement), S10-S14. doi:<https://doi.org/10.1016/j.ypmed.2017.05.006>
- Nichter, M., Padmawati, S., Danardono, M., Ng, N., Prabandari, Y., & Nichter, M. (2009). Reading culture from tobacco advertisements in Indonesia. *Tobacco Control*, 18(2), 98-107. doi:10.1136/tc.2008.025809
- Niu, Z., Xie, C., Wen, X., Tian, F., Ding, P., He, Y., . . . Chen, W. Q. (2015). Placenta mediates the association between maternal second-hand smoke exposure during pregnancy and small for gestational age. *Placenta*, 36(8), 876-880. doi:10.1016/j.placenta.2015.05.005
- Niu, Z., Xie, C., Wen, X., Tian, F., Yuan, S., Jia, D., & Chen, W. Q. (2016). Potential pathways by which maternal second-hand smoke exposure during pregnancy causes full-term low birth weight. *Scientific Reports*, 6, 24987.
- Öberg, M., Woodward, A., Jaakkola, M. S., Peruga, A., & Prüss-Ustün, A. (2011). WHO Global estimate of the burden of disease from second-hand smoke. Retrieved from: http://apps.who.int/iris/bitstream/10665/44426/1/9789241564076_eng.pdf
- Prochaska, J. O., & Velicer, W. F. (1997). The Transtheoretical Model of Health Behavior Change. *American Journal of Health Promotion*, 12(1), 38-48. doi:10.4278/0890-1171-12.1.38

- Stanković, A., Nikolić, M., & Arandelović, M. (2012). Exposure to environmental tobacco smoke and absence from work in women in Niš, Serbia. *Central European Journal Public Health*, 20(1), 24-28.
- Taylor, G., McNeill, A., Girling, A., Farley, A., Lindson-Hawley, N., & Aveyard, P. (2014). Change in mental health after smoking cessation: Systematic review and meta-analysis. *BMJ : British Medical Journal*, 348. doi:10.1136/bmj.g1151
- Venkatesh, K., Ziotnick, C., Triche, E., Ware, C. & Phipps, M. (2014). Accuracy of brief screening tools for identifying postpartum depression among adolescent mothers. *Pediatrics*, 113(1), e45-e53. doi: 10.1542/peds.2013-1628
- Vila Candel, R., Soriano-Vidal, F. J., Hevilla Cucarella, E., Castro-Sanchez, E., & Martin-Moreno, J. M. (2015). Tobacco use in the third trimester of pregnancy and its relationship to birth weight. A prospective study in Spain. *Women Birth*, 28(4), e134-139. doi:10.1016/j.wombi.2015.06.003
- Wahabi, H. A., Alzeidan, R. A., Fayed, A. A., Mandil, A., Al-Shaikh, G., & Esmaeil, S. A. (2013). Effects of secondhand smoke on the birth weight of term infants and the demographic profile of Saudi exposed women. *BMC Public Health*, 13, 341. doi:10.1186/1471-2458-13-341
- Wardlaw, T. M. Blanc, A., Zupan, J., & Ahman, E. (2004). *Low birthweight: Country, regional and global estimates*. (UNICEF.)
https://www.unicef.org/publications/files/low_birthweight_from_EY.pdf
- Weng, S. C., Huang, J. P., Huang, Y. L., Lee, T. S., & Chen, Y. H. (2016). Effects of tobacco exposure on perinatal suicidal ideation, depression, and anxiety. *BMC*

Public Health, 16, 623. doi:10.1186/s12889-016-3254-z

Wipfli, H., Avila-Tang, E., Navas-Acien, A., Kim, S., Onicescu, G., Yuan, J., . . . for the Famri Homes Study, I. (2008). Secondhand smoke exposure among women and children: Evidence from 31 countries. *American Journal of Public Health*, 98(4), 672-679. doi:10.2105/AJPH.2007.126631

World Health Organization. (2008). *WHO report on the global tobacco epidemic: The MPOWER package*. (World Health Organization.)

World Health Organization. (2012a). *Global adult tobacco survey: Indonesia report 2011*. Geneva: WHO.

World Health Organization. (2012b). *WHO global report: Mortality attributable to tobacco* (Report No. ISBN 978 92 4 156443 4). Geneva, Switzerland: WHO.

World Health Organization. (2014). *Global Nutrition Targets 2025: Policy brief series: low birth weight*. Retrieved from http://www.who.int/nutrition/topics/globaltargets_lowbirthweight_policybrief.pdf

World Health Organization. (2015a). *Health in 2015: From MDGs to SDGs: General introduction*.

World Health Organization. (2015b). *WHO report on the global tobacco epidemic, 2015: Raising taxes on tobacco*.

World Health Organization. (2016). *WHO Media center tobacco fact sheet*. Retrieved from <http://www.who.int/mediacentre/factsheets/fs339/en/>

Zhou, S., Rosenthal, D. G., Sherman, S., Zelikoff, J., Gordon, T., & Weitzman, M.

(2014). Physical, behavioral, and cognitive effects of prenatal tobacco and postnatal secondhand smoke exposure. *Current Problems in Pediatric & Adolescent Health Care*, 44(8), 219-241.