

## 参考文献

- Acar, N., Ozcelik, H., Cevik, A.A., Ozakin, E., Yorulmaz, G., Kebapci, N., ... Bilgin, M. (2014). Low perfusion index affects the difference in glucose level between capillary and venous blood. *Therapeutics and Clinical Risk Management*, 20 (10), 985-991.
- Apfelbaum, J.L., Hagberg, C.A., Caplan, R.A., ... Ovassapian, A. (2013). Practice Guidelines for Management of the Difficult airway: An Updated Report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology*, 118(2), 251-270.
- Applegate, R.L., Dorotta, I.L., Wells, B., Juma, D., & Applegate, P.M. (2016). The relationship between oxygen reserve index and arterial partial pressure of oxygen during surgery. *Anesthesia and Analgesia*, 123(3), 626-633.
- Baraka, A.S., & Salem, M.R. (2012). Hagberg, C.A. (ed), Preoxygenation. *Benumof and Hagberg's Airway Management*, 3rd ed (pp.280-300). Elsevier Saunders.
- Baraka, A.S., Taha, S.K., Aouad, M.T., El-Khatib, M.F., & Kawkabani, N.I. (1999). Preoxygenation Comparison of Maximal Breathing and Tidal Volume Breathing Techniques. *Anesthesiology*, 91, 612-616.
- Barrett, K.E., Barman, S.M., Boitano, S., & Brooks, H.L. (2012/2014). 岡田泰伸 (監訳), ギャノン生理学 原著 24 版 (pp.730). 丸善出版.
- Benumof, J.L. (1999). Preoxygenation: best method for both efficacy and efficiency. *Anesthesiology*, 91, 603-605.

- Berry, C.B., & Myles, P.S. (1994). Pre-oxygenation in healthy volunteers—a graph of oxygen “washin” using end tidal oxygraphy. *British Journal of Anaesthesia*, 72, 116–118.
- Bhatia, P.K., Bhandari, S.C., Tulsiani, K.L., & Kumar, Y. (1997). End-tidal oxygraphy and safe duration of apnoea in young adults and elderly patients. *Anaesthesia*, 52, 175-178.
- Bouroche, G., & Bourgain, J.L. (2015). Preoxygenation and general anesthesia: a review. *Minerva Anesthesiologica*, 81(8), 910-920.
- Campbell, I.T., & Beatty, P.C. (1994). Monitoring preoxygenation. *British Journal of Anaesthesia*, 72, 3-4.
- Drummond, G.B., & Park, G.R. (1984). Arterial Oxygen Saturation Before Intubation of Trachea: An assessment of oxygenation techniques. *British Journal of Anaesthesia*, 56, 987.
- Edmark, L., Aherdan, K.K., Enlund, M., & Hedenstierna, G. (2003). Optimal oxygen concentration during induction of general anesthesia. *Anesthesiology*, 98, 28–33.
- Edmark, L., Auner, U., Enlund, M., Ostberg, E., & Hedenstierna, G. (2011). Oxygen concentration and characteristics of progressive atelectasis formation during anaesthesia. *Acta Anaesthesiologica Scandinavica*, 55, 75–81.

Ehrenfeld, J.M., Funk, L.M., Schalkwyk, J.V., Merry, A.F., Sandberg, W.S., & Gawande, A. (2010). The incidence of hypoxemia during surgery: evidence from two institutions. *Canadian Journal of Anesthesia*, 57(10), 888–97. doi: 10.1007/s12630-010-9366-5.

Gagnon, C., Fortier, L.P., & Donati, F. (2006). When a leak is unavoidable, preoxygenation is equally ineffective with vital capacity or tidal volume breathing. *Canadian Journal of Anesthesia*, 53(1), 86–91.

Gambee, A.M., Hertzka, R.E., & Fisher, D.M. (1987). Preoxygenation techniques: Comparison of three minutes and 4 breaths. *Anesthesia and Analgesia*, 66, 468–470.

Hamilton, W.K., & Eastwood, D.W. (1955). A study of denitrogenation with some inhalation anesthetic systems. *Anesthesiology*, 16, 861-867.

半沢篤, 福田謙一, 小野寺隆昭, 本田藍, 半田俊之, 高北義彦, 一戸達也. (2015). プロポフォール・レミフェンタニルによる麻酔導入における術前不安の影響 上下顎移動術を受ける患者を対象として. *日本歯科麻酔学会雑誌*, 43 (2), 283-285.

Hedenstierna, G. (2012). Oxygen and anesthesia: what lung do we deliver to the post-operative ward? *Acta Anaesthesiologica Scandinavica*, 56 (6), 675–85. doi:10.1111/j.1399-6576.2012.02689.x.

Hedenstierna, G., & Edmark, L. (2010). Mechanisms of atelectasis in the perioperative period. *Best Practice & Research: Clin Anaesthesiology*, 24, 157-169.

Hedenstierna G, Edmark, L., & Aherdan, K.K. (2000). Time to reconsider the pre-oxygenation during induction of anaesthesia. *Minerva Anestesiologica*, 66, 293-296.

Herriger, A., Frascarolo, P., Spahn, D.R., & Magnusson, L. (2004). The effect of positive airway pressure during pre-oxygenation and induction of anaesthesia upon duration of non-hypoxic apnoea. *Anaesthesia*, 59, 243-247.

平田直之, 茶木友浩, 山蔭道明. (2016). Oxygen reserve index: 新しい酸素化モニターの可能性. *臨床麻酔*, 40 (10), 1391-1397.

Kundra, P., Stephen, S., & Vinayagam, S., (2013). Techniques of preoxygenation in patients with ineffective face mask seal. *Indian Journal of Anaesthesia*, 57(2), 175–179. doi: 10.4103/0019-5049.111847

Lane, S., Saunders, D., Schofield, A., Padmanabhan, R., Hildreth, A. & Laws, D. (2005). A prospective, randomised controlled trial comparing the efficacy of pre-oxygenation in the 20° head-up vs supine position. *Anaesthesia*, 60, 1064–1067.

Lye, S.T., Liaw, C.M., Seet, E., & Koh, K.F. (2013). Comparison of results from novice and trained personnel using the Macintosh laryngoscope, Pentax AWS<sup>®</sup>, C-MAC<sup>™</sup> and Bonfils intubation fibroscope: a manikin study. *Singapore Medical Journal*, 54, 64–68. doi: 10.11622/smedj.2013026.

Martin, D.S., & Grocott, M.P. (2013). Oxygen therapy in anaesthesia: the yin and yang of O<sub>2</sub>. *British Journal of Anaesthesia*, 111 (6), 867-871. doi: 10.1093/bja/aet291.

Masimo Whitepaper: 灌流指標 (Perfusion Index). (2007).

<http://www.masimo.com/pdf/whitepaper/LAB3410F.pdf>. (検索日 2017 年 6 月 25 日)

Masimo Whitepaper: Oxygen Reserve Index (ORi<sup>™</sup>).

[http://www.masimo.co.uk/pdf/ori/LAB8543A\\_Whitepaper\\_ORI\\_British.pdf](http://www.masimo.co.uk/pdf/ori/LAB8543A_Whitepaper_ORI_British.pdf). (検索  
日 2017 年 6 月 25 日)

Mavridou, P., Dimitriou, V., Manataki, A., Arnaoutoglou, E., & Papadopoulos, G.  
(2013). Patient's anxiety and fear of anesthesia: effect of gender, age, education, and  
previous experience of anesthesia: a survey of 400 patients. *J Anesth*, 27(1), 104-108.  
doi:10.1007/s00540-012-1460-0

McCrory, J.W., & Matthews, J.N. (1990). Comparison of Four Methods of  
Preoxygenation. *British Journal of Anaesthesia*, 64, 571-576.

McGowan, P., & Skinner, A. (1995). Preoxygenation: The importance of a good face  
mask seal. *British Journal of Anaesthesia*, 75, 777-778.

Miller, R.D. (2004/2007). 武田純三 (監). ミラー麻酔科学 第6版. (p548, 557).  
メディカル・サイエンス・インターナショナル.

日本呼吸器学会肺生理専門委員会 (編). (2014). *Q&A パルスオキシメータハンド  
ブック* (pp.12). 法研.

Nimmagadda, U., Chiravuri, S.D., Salem, M.R., Joseph, N.J., Wafai, Y., Crystal, G.J.,  
& El-Orbany, M.I. (2001). Preoxygenation with Tidal Volume and Deep Breathing  
Techniques: The Impact of Duration of Breathing and Fresh Gas Flow. *Anesthesia  
and Analgesia*, 92(5), 1337-1341.

Nimmagadda, U., Salem, M.R., & Crystal, G.J. (2017). Preoxygenation: Physiologic  
Basis, Benefits, and Potential Risks. *Anesthesia and Analgesia*, 124(2), 507-517.

Nimmagadda, U., Salem, M.R., Joseph, N.J., Lopez, G., Megally, M., Lang, D.J., & Wafai, Y. (2000). Efficacy of preoxygenation with tidal volume breathing: Comparison of breathing systems. *Anesthesiology*, 93, 693-698.

Pandit, J.J., Duncan, T., & Robbins, P.A. (2003). Total Oxygen Uptake with Two Maximal Breathing Techniques and the Tidal Volume Breathing Technique: A Physiologic Study of Preoxygenation. *Anesthesiology*, 99, 841–846.

Pedersen, T., Nicholson, A., Hovhannisyan, K., Moller, A.M., Smith, A.F., & Lewis, S.R. (2014). Pulse oximetry for perioperative monitoring. *Cochrane Database Systematic Reviews* 2014, 3.

Rooney, M.J. (1994). Pre-oxygenation: a comparison of two techniques using a Bain system. *Anaesthesia*, 49, 629-632.

Rothen, H.U., Neumann, P., Berglund, J.E., Valtysson, J., Magnusson, A., & Hedenstierna, G. (1999). Dynamics of re-expansion of atelectasis during general anaesthesia. *British Journal of Anaesthesia*, 82, 551–556.

Rothen, H.U., Sporre, B., Engberg, G., Wegenius, G., Reber, A., & Hedenstierna, G. (1995). Prevention of atelectasis during general anaesthesia. *Lancet*, 345, 1387-1391.

Russell, G.N., Smith, C.L., Snowdon, S.L., & Bryson, T.H. (1987). Pre-oxygenation and the parturient patient. *Anaesthesia*, 42, 346–51.

Samain, E., Farah, E., Delefosse, D., & Marty, J. (2000). End-tidal oxygraphy during pre-oxygenation in patients with severe diffuse emphysema. *Anaesthesia*, 55, 841-846.

Scheeren, T.W.L., Belda, F. J., & Perel, A. (2017). The oxygen reserve index (ORI): a new tool to monitor oxygen therapy. *Journal of Clinical Monitoring and Computing*, 8. doi: 10.1007/s10877-017-0049-4.

Shah, U., Wong, J., Wong, D.T., & Chung, F. (2016). Preoxygenation and intraoperative ventilation strategies in obese patients: a comprehensive review. *Current Opinion Anaesthesiology*, 29 (1), 109-118.

Singh, B., Afzal, L., Kaur, B., & Osahan, N. K. (2006). Comparison of Pre-Oxygenation by Maximal Breathing and Tidal Volume Breathing Techniques. *Indian Journal of Anaesthesia*, 50(3), 209-213.

Szmuk, P., Steiner, J.W., Olomu, P.N., Ploski, R.P., Sessler, D.I., & Ezri, T. (2016). Oxygen reserve index: A novel noninvasive measure of oxygen reserve- A pilot study. *Anesthesiology*, 124 (4), 779-784.

Tanoubi, I., Drolet, P., & Donati, F. (2009). Optimizing preoxygenation in adults. *Canadian Journal of Anesthesia*, 56(6), 449-466.

内田慎也, 荻野祐一, 齋藤繁. (2011). エアウェイスコープ®の習熟度評価と挿管困難対策用具としての有用性について. *日本臨床麻酔学会誌*, 31 (4), 678-674.

吉田奏, 宮坂勝之. (2013). 手術室におけるチーム医療 周麻酔期看護とは. *日本手術医学会誌*, 34 (2), 121-125.