

引用文献

- Absalom, A. R., Mani, V., De Smet, T., & Struys, M. M. (2009). Pharmacokinetic models for propofol--defining and illuminating the devil in the detail. *British Journal of Anaesthesia*, 103(1), 26-37.
- Ahmetovic-Djug, J., Hasukic, S., Djug, H., Hasukic, B., & Jahic, A. (2017). Impact of preoperative anxiety in patients on hemodynamic changes and a dose of anesthetic during induction of anesthesia. *Medical Archives (Sarajevo, Bosnia and Herzegovina)*, 71(5), 330-333.
- Badner, N. H., Nielson, W. R., Munk, S., Kwiatkowska, C., & Gelb, A. W. (1990). Preoperative anxiety: Detection and contributing factors. *Canadian Journal of Anaesthesia = Journal Canadien d'Anesthesie*, 37(4 Pt 1), 444-447.
- Bradshaw, P., Hariharan, S., & Chen, D. (2016). Does preoperative psychological status of patients affect postoperative pain? A prospective study from the caribbean. *British Journal of Pain*, 10(2), 108-115.
- Bradt, J., Dileo, C., & Shim, M. (2013). Music interventions for preoperative anxiety. *The Cochrane Database of Systematic Reviews*, (6):CD006908. doi(6), CD006908.
- Choong, E., Loryan, I., Lindqvist, M., Nordling, A., el Bouazzaoui, S., van Schaik, R. H., et al. (2013). Sex difference in formation of propofol metabolites: A replication study. *Basic & Clinical Pharmacology & Toxicology*, 113(2), 126-131.
- Domar, A. D., Everett, L. L., & Keller, M. G. (1989). Preoperative anxiety: Is it a predictable entity? *Anesthesia and Analgesia*, 69(6), 763-767.
- Domschke, K., Tidow, N., Schrempf, M., Schwarte, K., Klauke, B., Reif, A., et al. (2013). Epigenetic signature of panic disorder: A role of glutamate decarboxylase 1 (GAD1) DNA hypomethylation? *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 46, 189-196.
- Erden, V., Yangin, Z., Erkalp, K., Delatioglu, H., Bahceci, F., & Seyhan, A. (2005). Increased progesterone production during the luteal phase of menstruation may decrease anesthetic requirement. *Anesthesia and Analgesia*, 101(4), 1007-11, table

of contents.

- Esclapez, M., Tillakaratne, N. J., Kaufman, D. L., Tobin, A. J., & Houser, C. R. (1994). Comparative localization of two forms of glutamic acid decarboxylase and their mRNAs in rat brain supports the concept of functional differences between the forms. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 14(3 Pt 2), 1834-1855.
- Fu, F., Chen, X., Feng, Y., Shen, Y., Feng, Z., & Bein, B. (2014). Propofol EC50 for inducing loss of consciousness is lower in the luteal phase of the menstrual cycle. *British Journal of Anaesthesia*, 112(3), 506-513.
- Gepts, E., Camu, F., Cockshott, I. D., & Douglas, E. J. (1987). Disposition of propofol administered as constant rate intravenous infusions in humans. *Anesthesia and Analgesia*, 66(12), 1256-1263.
- Hirose, K., Kuroishi, T., Inoue, M., & Tajima, K. (1992). Changing patterns of cancer mortality among the elderly population in japan. *Japanese Journal of Clinical Oncology*, 22(6), 428-432.
- Hoymork, S. C., & Raeder, J. (2005). Why do women wake up faster than men from propofol anaesthesia? *British Journal of Anaesthesia*, 95(5), 627-633.
- Jawaid, M., Mushtaq, A., Mukhtar, S., & Khan, Z. (2007). Preoperative anxiety before elective surgery. *Neurosciences (Riyadh, Saudi Arabia)*, 12(2), 145-148.
- Johnston, M. (1980). Anxiety in surgical patients. *Psychological Medicine*, 10(1), 145-152.
- Kasa, K., Otsuki, S., Yamamoto, M., Sato, M., Kuroda, H., & Ogawa, N. (1982). Cerebrospinal fluid gamma-aminobutyric acid and homovanillic acid in depressive disorders. *Biological Psychiatry*, 17(8), 877-883.
- Kay, B., & Rolly, G. (1977). I.C.I. 35868, a new intravenous induction agent. *Acta Anaesthesiologica Belgica*, 28(4), 303-316.
- Kay, B., & Stephenson, D. K. (1980). ICI 35868 (diprivan): A new intravenous anaesthetic. A comparison with althesin. *Anaesthesia*, 35(12), 1182-1187.
- Kim, S. Y., Park, J. E., Lee, Y. J., Seo, H. J., Sheen, S. S., Hahn, S., et al. (2013). Testing a tool for assessing the risk of bias for nonrandomized studies showed

- moderate reliability and promising validity. *Journal of Clinical Epidemiology*, 66(4), 408-414.
- Kindler, C. H., Harms, C., Amsler, F., Ihde-Scholl, T., & Scheidegger, D. (2000). The visual analog scale allows effective measurement of preoperative anxiety and detection of patients' anesthetic concerns. *Anesthesia and Analgesia*, 90(3), 706-712.
- 小板橋 俊哉. (2007). 麻酔科研修における tiva/tci 教育の意義 tiva/tci の現状. *日本臨床麻酔学会誌*, 27(7), 652-657.
- 小板橋 俊哉. (2016). プロポフォール麻酔の基本. *日本臨床麻酔学会誌*, 36(4), 441-447.
- Kubo, K., Nishikawa, K., Hardy-Yamada, M., Ishizeki, J., Yanagawa, Y., & Saito, S. (2009). Altered responses to propofol, but not ketamine, in mice deficient in the 65-kilodalton isoform of glutamate decarboxylase. *The Journal of Pharmacology and Experimental Therapeutics*, 329(2), 592-599.
- ロナルド・D.ミラー(編).(2007). 武田純三(監訳) 非オピオイド静脈薬. *Miller's anesthesia*(第 6 版) (pp. 251-258).メディカル・サイエンス・インターナショナル
- Loryan, I., Lindqvist, M., Johansson, I., Hiratsuka, M., van der Heiden, I., van Schaik, R. H., et al. (2012). Influence of sex on propofol metabolism, a pilot study: Implications for propofol anesthesia. *European Journal of Clinical Pharmacology*, 68(4), 397-406.
- Lysakowski, C., Dumont, L., Czarnetzki, C., Bertrand, D., Tassonyi, E., & Tramer, M. R. (2006). The effect of cigarette smoking on the hypnotic efficacy of propofol. *Anaesthesia*, 61(9), 826-831.
- Lysakowski, C., Elia, N., Czarnetzki, C., Dumont, L., Haller, G., Combescure, C., et al. (2009). Bispectral and spectral entropy indices at propofol-induced loss of consciousness in young and elderly patients. *British Journal of Anaesthesia*, 103(3), 387-393.
- Marsh, B., White, M., Morton, N., & Kenny, G. N. (1991). Pharmacokinetic model driven infusion of propofol in children. *British Journal of Anaesthesia*, 67(1), 41-48.
- Masui, K., Upton, R. N., Doufas, A. G., Coetzee, J. F., Kazama, T., Mortier, E. P., et al.

- (2010). The performance of compartmental and physiologically based recirculatory pharmacokinetic models for propofol: A comparison using bolus, continuous, and target-controlled infusion data. *Anesthesia and Analgesia*, 111(2), 368-379.
- Milne, S. E., Troy, A., Irwin, M. G., & Kenny, G. N. (2003). Relationship between bispectral index, auditory evoked potential index and effect-site EC50 for propofol at two clinical end-points. *British Journal of Anaesthesia*, 90(2), 127-131.
- Moerman, N., van Dam, F. S., Muller, M. J., & Oosting, H. (1996). The amsterdam preoperative anxiety and information scale (APAIS). *Anesthesia and Analgesia*, 82(3), 445-451.
- 西山美鈴(編).(2013).静脈麻酔.麻酔科レジデントマニュアル(171-182).ライフリサーチプレス
- Ochs, H. R., Greenblatt, D. J., & Knuchel, M. (1985). Kinetics of diazepam, midazolam, and lorazepam in cigarette smokers. *Chest*, 87(2), 223-226.
- 尾花初美, 杉下美穂子, 松村栄子, 他. 手術を受ける患者の不安の程度と関連要因. 第3回日本看護学会論文集 (成人看護 I) 1999:105-107.
- Petty, F., & Sherman, A. D. (1984). Plasma GABA levels in psychiatric illness. *Journal of Affective Disorders*, 6(2), 131-138.
- Reed, M. D., Yamashita, T. S., Marx, C. M., Myers, C. M., & Blumer, J. L. (1996). A pharmacokinetically based propofol dosing strategy for sedation of the critically ill, mechanically ventilated pediatric patient. *Critical Care Medicine*, 24(9), 1473-1481.
- Rigouzzo, A., Servin, F., & Constant, I. (2010). Pharmacokinetic-pharmacodynamic modeling of propofol in children. *Anesthesiology*, 113(2), 343-352.
- Sahinovic, M. M., Eleveld, D. J., Miyabe-Nishiwaki, T., Struys, M. M. R. F., & Absalom, A. R. (2017). Pharmacokinetics and pharmacodynamics of propofol: Changes in patients with frontal brain tumours. *British Journal of Anaesthesia*, 118(6), 901-909.
- Schnider, T. W., Minto, C. F., Gambus, P. L., Andresen, C., Goodale, D. B., Shafer, S. L., et al. (1998). The influence of method of administration and covariates on the pharmacokinetics of propofol in adult volunteers. *Anesthesiology*, 88(5), 1170-1182.

- Schnider, T. W., Minto, C. F., Gambus, P. L., Andresen, C., Goodale, D. B., Shafer, S. L., et al. (1998). The influence of method of administration and covariates on the pharmacokinetics of propofol in adult volunteers. *Anesthesiology*, 88(5), 1170-1182.
- Schnider, T. W., Minto, C. F., Shafer, S. L., Gambus, P. L., Andresen, C., Goodale, D. B., et al. (1999). The influence of age on propofol pharmacodynamics. *Anesthesiology*, 90(6), 1502-1516.
- Shevde, K., & Panagopoulos, G. (1991). A survey of 800 patients' knowledge, attitudes, and concerns regarding anesthesia. *Anesthesia and Analgesia*, 73(2), 190-198.
- Xu, Z., Liu, F., Yue, Y., Ye, T., Zhang, B., Zuo, M., et al. (2009). C50 for propofol-remifentanyl target-controlled infusion and bispectral index at loss of consciousness and response to painful stimulus in chinese patients: A multicenter clinical trial. *Anesthesia and Analgesia*, 108(2), 478-483.
- Yavuz, L., Eroglu, F., Ceylan, B. G., Ozsoy, H. M., & Ozbasar, D. (2007). High estradiol levels and depth of anaesthesia. *Clinical and Experimental Obstetrics & Gynecology*, 34(1), 31-34.