

- Ahlund, S., Nordgren, B., Wilander, E., Wiklund, I., & Fridén, C. (2013). Is home-based pelvic floor muscle training effective in treatment of urinary incontinence after birth in primiparous women? A randomized controlled trial. *Acta Obstetricia Et Gynecologica Scandinavica*, 92 (8), 909-915. doi:10.1111/aogs.12173.
- Albrich, S., Steetskamp, J., Knoechel, S., Porta, S., Hoffmann, G., & Skala, C. (2016). Assessment of pelvic floor muscle contractility: Digital palpation versus 2D and 3D perineal ultrasound. *Archives of Gynecology and Obstetrics*, 293 (4), 839-843. doi:10.1007/s00404-015-3897-5.
- Antonovsky, A. (1987) . Unraveling the Mystery of Health: How People Manage Stress and Stay Well. San Francisco: Jossey-Bass. 山崎喜比古, 吉井清子. (2001) .健康の謎を解く—ストレス対処と健康保持のメカニズム. 東京:有信堂高文社.
- Ariail, A., Sears, T., & Hampton, E., (2008). Use of transabdominal ultrasound imaging in retraining the pelvic-floor muscles of a woman postpartum. *Physical Therapy*, 88 (10), 1208-1217. doi:10.2522/ptj.20070330.
- Bandura, A. (1977) Social Learning Theory. Prentice Hall.原野広太郎,社会的学習理論 オンデマンド版—人間理解と教育の基礎.東京:金子書房,2012.
- Barton, A., Serrao, C., Thompson, J., & Briffa, K., (2015). Transabdominal ultrasound to assess pelvic floor muscle performance during abdominal curl in exercising women. *International Urogynecology Journal*, 26 (12), 1789-1795. doi:10.1007/s00192-015-2791-9.
- Bø, K. (2004). Pelvic floor muscle training is effective in treatment of

female stress urinary incontinence, but how does it work?.*International Urogynecology Journal*, 15(2), 76-84. doi:10.1007/s00192-004-1125-0.

Bø, K., Lilleas, F., Talseth, T., Hedland, H. (2001a). Dynamic MRI of the Pelvic Floor Muscles in an Upright Sitting Position. *Neurourology and Urodynamics*, 20 (2). 167-174. doi:AID-NAU19>3.0.CO;2-4.

Bø, K., & Finckenhagen, H. B. (2001b). Vaginal palpation of pelvic floor muscle strength: Inter-test reproducibility and comparison between palpation and vaginal squeeze pressure. *Acta Obstetricia Et Gynecologica Scandinavica*, 80 (10), 883-887. doi:10.1034/j.1600-0412.2001.801003.x.

Bø, K., & Sherburn, M. (2005). Evaluation of female pelvic-floor muscle function and strength. *Physical Therapy*, 85 (3), 269-282.

Bø, K., Brækken, I.,H., Majida, M., & Engh, M. E. (2009). Constriction of the levator hiatus during instruction of pelvic floor or transversus abdominis contraction: A 4D ultrasound study. *International Urogynecology Journal*, 20(1), 27-32. doi:<http://dx.doi.org/10.1007/s00192-008-0719-3>.

Bø, K., & Herbert, R. D. (2013). There is not yet strong evidence that exercise regimens other than pelvic floor muscle training can reduce stress urinary incontinence in women: A systematic review. *Journal of Physiotherapy*, 59 (3), 159-168. doi:10.1016/S1836-9553(13)70180-2.

Botelho, S., Pereira, L. C., Marques, J., Lanza, A. H., Amorim, C. F.,P alma, P., & Riccetto, C. (2013). Is there correlation between electromyography and digital palpation as means of measuring

- pelvic floor muscle contractility in nulliparous, pregnant, and postpartum women? *Neurourology and Urodynamics*, 32(5), 420 -423. doi:10.1002/nau.22321.
- Boyle, R., Hay-Smith, E. J. C., Cody, J. D., & Mørkved, S. (2012). Pelvic floor muscle training for prevention and treatment of urinary and faecal incontinence in antenatal and postnatal women. *The Cochrane Database of Systematic Reviews*, 10, CD007471.
- Boyle, R., Hay - Smith, E. J. C., Cody, J. D., & Mørkved, S. (2014). Pelvic floor muscle training for prevention and treatment of urinary and fecal incontinence in antenatal and postnatal women: A short version cochrane review. *Neurourology and Urodynamics*, 33(3), 269-276. doi:10.1002/nau.22402.
- Brækken, I. H., Majida, M., Engh, M. E., & Bø, K. (2009). Test-retest reliability of pelvic floor muscle contraction measured by 4D ultrasound. *Neurourology and Urodynamics*, 28(1), 68-73. doi:10.1002/nau.20618.
- Braekken, I. H., Hoff Braekken, I., Majida, M., Engh, M. E., & Bø, K. (2010). Morphological changes after pelvic floor muscle training measured by 3-dimensional ultrasonography: A randomized controlled trial. *Obstetrics and Gynecology*, 115(2 Pt 1), 317-324. doi:10.1097/AOG.0b013e3181cbd35f.
- Camargo, F., Rodrigues, A., Arruda, R., Ferreira Sartori, M., Girao, M., & Castro, R. (2009). Pelvic floor muscle training in female stress urinary incontinence: Comparison between group training and individual treatment using PERFECT assessment scheme. *International Urogynecology Journal*, 20(12), 1455-1462.
- Chan, S. S. C., Cheung, R. Y. K., Yiu, K. W., Lee, L. L., & Chung, T.

- K. H. (2014). Pelvic floor biometry in chinese primiparous women 1 year after delivery: A prospective observational study. *Ultrasound in Obstetrics & Gynecology*, 43(4), 466-474. doi:10.1002/uog.13249.
- Chehrehrazi, M., Arab, A., Karimi, N., & Zargham, M. (2009). Assessment of pelvic floor muscle contraction in stress urinary incontinent women: Comparison between transabdominal ultrasound and perineometry. *International Urogynecology Journal*, 20 (12), 1491-1496. doi:10.1007/s00192-009-0977-8.
- Chen, S. (2004). The development and testing of the pelvic floor muscle exercise self-efficacy scale. *Journal of Nursing Research*, 12 (4), 257-266. Doi:10.1097/01.JNR.0000387510.52243.c8.
- Chen, S., & Tzeng, Y. (2009). Path analysis for adherence to pelvic floor muscle exercise among women with urinary incontinence. *Journal of Nursing Research*, 17 (2), 83-92. doi:10.1097/JNR.0b013e3181a53e7e.
- Chen, R., Song, Y., Jiang, L., Hong, X., & Ye, P. (2011). The assessment of voluntary pelvic floor muscle contraction by three-dimensional transperineal ultrasonography. *Archives of Gynecology and Obstetrics*, 284 (4), 931-936. doi:10.1007/s00404-010-1795-4.
- Christensen, L. L., Djurhuus,J.C., & Constantinou, C.E., (1995). Imaging of pelvic floor contractions using MRI. *Neurourology and Urodynamics*, 14 (3), 209-216. doi.org/10.1002/nau.1930140302.
- DeLancey, J. O. L., Kearney, R., Chou, Q., Speights, S., & Binno, S.(2003). The appearance of levator ani muscle abnormalities in magnetic resonance images after vaginal delivery. *Obstetrics & Gynecology*, 101 (1), 46-53.

- Dietz, H. P., Wilson, P. D., & Clarke, B. (2001). The use of perineal ultrasound to quantify levator activity and teach pelvic floor muscle exercises. *International Urogynecology Journal*, 12(3), 166-169. doi:10.1007/s001920170059.
- Dietz, H. P., Steensma, A. B., & Vancaillie, T. G. (2003). Levator function in nulliparous women. *International Urogynecology Journal*, 14(1), 24-26. doi:10.1007/s00192-002-1001-8.
- Dietz, H. P., & Lanzarone, V. (2005). Levator trauma after vaginal delivery. *Obstetrics and Gynecology*, 106(4), 707-712. doi:10.1097/01.AOG.0000178779.62181.01.
- Dinc, A., Kizilkaya Beji, N., & Yalcin, O. (2009). Effect of pelvic floor muscle exercises in the treatment of urinary incontinence during pregnancy and the postpartum period. *International Urogynecology Journal*, 20 (10), 1223-1231. doi:10.1007/s00192-009-0929-3.
- Dumoulin, C., Lemieux, M., Bourbonnais, D., Gravel, D., Bravo, G., & Morin, M. (2004). Physiotherapy for persistent postnatal stress urinary incontinence: A randomized controlled trial. *Obstetrics and Gynecology*, 104(3), 504-510. doi:10.1097/01.AOG.0000135274.92416.62.
- Dumoulin, C., Glazener, C., & Jenkinson, D. (2011). Determining the optimal pelvic floor muscle training regimen for women with stress urinary incontinence. *Neurourology and Urodynamics*, 30(5), 746-753. doi:10.1002/nau.21104.
- Dumoulin, C., & Hay-Smith, J. (2014). Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *The Cochrane Database of Systematic Reviews*.

*eviews*, (1), CD005654. doi:10.1002/14651858.CD005654.pub3

- Dumoulin, C., Hay-Smith, J., Frawley, H., McClurg, D., Alewijnse, D., Bo, K., . . . Van Kampen, M. (2015). 2014 consensus statement on improving pelvic floor muscle training adherence: International continence society 2011 State - of - the - Scienceseminar. *Neurourology and Urodynamics*, 34(7), 600-605. doi:10.1002/nau.22796.
- Dumoulin, C., Cacciari, L. P., & Hay-Smith, E. J. C. (2018). Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *The Cochrane Database of Systematic Reviews*, 10, CD005654. doi:10.1002/14651858.CD005654.pub4.
- Eriksson, M., & Mittelmark, M. B. (2017). The sense of coherence and its measurement. *The handbook of salutogenesis*. 97-106.
- Ferreira, C. H. J., Barbosa, P. B., Souza, F. d. O., Antônio, F. I., Franco, M. M., & Bø, K. (2010). Inter-rater reliability study of the modified oxford grading scale and the peritron manometer. *Physiotherapy*, 97(2), 132-138. doi:10.1016/j.physio.2010.06.007.
- Fitz, F. F., Stüpp, L., Costa, T. F., Sartori, M. G. F., Girão, Manoel João Batista Castello, & Castro, R. A. (2016). Correlation between maximum voluntary contraction and endurance measured by digital palpation and manometry: An observational study. *Revista Da Associação Médica Brasileira*, 62(7), 635-640.
- 福田佳治, 八幡裕一郎,& 今井博之. 一目でわかるヘルスプロモーション: 理論と実践ガイドブック.14-35, 埼玉: 国立保健医療科学院.
- Frawley, H. C., Galea, M. P., Phillips, B. A., Sherburn, M., & Bø, K. (2006a). Reliability of pelvic floor muscle strength assessment

- using different test positions and tools. *Neurourology and Urodynamics*, 25(3), 236-242. doi:10.1002/nau.20201.
- Frawley, H. C., Galea, M. P., Phillips, B. A., Sherburn, M., & Bø, K. (2006b). Effect of test position on pelvic floor muscle assessment. *International Urogynecology Journal and Pelvic Floor Dysfunction*, 17(4), 365-371.
- Germain B,C. (1996) / 田中美緒監訳 (2015) . 女性の骨盤 妊娠・出産における身体的変化とエクササイズ. 24-32, 東京:株式会社メディカルプレス.
- Gotoh, M., Homma, Y., Funahashi, Y., Matsukawa, Y., & Kato, M. (2009). Psychometric validation of the Japanese version of the international consultation on incontinence Questionnaire - Short form. *International Journal of Urology*, 16(3), 303-306. doi:10.1111/j.1442-2042.2008.02237.x.
- Guzmán Rojas, R., Wong, V., Shek, K., & Dietz, H. (2014). Impact of levator trauma on pelvic floor muscle function. *International Urogynecology Journal*, 25(3), 375-380. doi:10.1007/s00192-013-2226-4.
- Hagen, S., McClurg, D., Bugge, C., Hay-Smith, J., Dean, S. G., Elders, A., Wilson, L. (2019). Effectiveness and cost-effectiveness of basic versus biofeedback-mediated intensive pelvic floor muscle training for female stress or mixed urinary incontinence: Protocol for the OPAL randomised trial. *BMJ Open*, 9 (2), e024153. doi:10.1136/bmjopen-2018-024153.
- 畠栄一,& 土井由利子. 行動科学 健康づくりのための理論と応用. 25-28, 東京:株式会社南江堂.
- Herderschee, R., Hay-Smith, E. C. J., Herbison, G. P., Roovers, J. P.,

- & Heineman, M. J. (2013). Feedback or biofeedback to augment pelvic floor muscle training for urinary incontinence in women: Shortened version of a cochrane systematic review. *Neuro urology and Urodynamics*, 32(4), 325-329. doi:10.1002/nau.22329.
- Hilde, G., Stær-Jensen, J., Ellström, E. M., Brækken, I., & Bø, K. (2012). Continence and pelvic floor status in nulliparous women at midterm pregnancy. *International Urogynecology Journal*, 23 (9), 1257-1263. doi:10.1007/s00192-012-1716-0.
- 泌尿器科領域の治療標準化に関する研究班 女性尿失禁診療ガイドライン EBMに基づく尿失禁診療ガイドライン. じほう ,2004.
- 本郷祥子, 藤崎章子, 下稻葉美佐, 植村天受,& 嘉村康邦. (2018). 騎乗型フィットネスマシーンは腹圧性尿失禁を改善するか. 日本女性骨盤底医学会誌, 15(1), 103-105.
- 池田真弓,& 森明子 (2016) . 経腔分娩後の骨盤底筋評価－内診による骨盤底筋力測定の有用性－.母性衛生, 56 (4) , 576-582.
- 池田真弓 (2019) . 分娩後の骨盤底筋訓練を助産師が経腔触診で指導することの実行可能性.日本助産学会誌,33 (1) , 14-26.
- 一般社団法人日本創傷・オストミー・失禁管理学会編 (2017) .コンチネンスケアの充実をめざして 排泄ケアガイドブック, 照林社.
- Isherwood,P.J., & Rane, A.(2000). Comparative assessment of pelvic floor strength using a perineometer and digital examination. British Journal of Obstetrics and Gynecology, 107,1007-1011.
- Junginger, B., Vollhaber, H., & Baessler, K. (2018). Submaximal pelvic floor muscle contractions: Similar bladder-neck elevation, longer duration, less intra-abdominal pressure. *International Urogynecology Journal*, 29 (11), 1681-1687.
- 亀崎明子,& 田中満由美.(2015).日本における女性尿失禁患者に対する指導の

実態.母性衛生,56(1),120-127.

- Kelly, M., Tan, B., Thompson, J., Carroll, S., Follington, M., Arndt, A., & Seet, M. (2007). Healthy adults can more easily elevate the pelvic floor in standing than in crook-lying: An experimental study. *The Australian Journal of Physiotherapy*, 53(3), 187. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17725476>.
- Kruger, J., Budgett, D., Goodman, J., & Bø, K. (2019). Can you train the pelvic floor muscles by contracting other related muscles? *Neurourology and Urodynamics*, 38(2), 677-683. doi:10.1002/nau.23890.
- Laycock, J., & Jerwood,D.(2001). Pelvic floor muscle assessment : The PERFECT scheme. *Physiotherapy*, 87(12),631-642. doi:10.1016/S0031-9406 (05) 61108-X.
- Leitner, M., Moser, H., Taeymans, J., Kuhn, A., & Radlinger, L. (2015). Pelvic floor muscle displacement during voluntary and involuntary activation in continent and incontinent women: A systematic review. *International Urogynecology Journal*, 26(11), 1587-1598. doi:10.1007/s00192-015-2700-2
- McClurg, D., Frawley, H., Hay - Smith, J., Dean, S., Chen, S., Chiarelli, P., ... Dumoulin, C. (2015). Scoping review of adherence promotion theories in pelvic floor muscle training – 2011 ics state - of - the - science seminar research paper i of iv. *Neurourology and Urodynamics*, 34 (7) , 606-614. doi:10.1002/nau.22769.
- Marques, J., Botelho, S., Pereira, L. C., Lanza, A. H., Amorim, C. F., Palma, P., & Riccetto, C. (2013). Pelvic floor muscle training program increases muscular contractility during first pregnancy and postpartum: Electromyographic study. *Neurourology and*

*Urodynamics*, 32 (7) , 998-1003. doi:10.1002/nau.22346.

Mercier, J., Morin, M., Zaki, D., Reichetzer, B., Lemieux, M., Khalifé, S., & Dumoulin, C. (2019). Pelvic floor muscle training as a treatment for genitourinary syndrome of menopause: A single-arm feasibility study. *Maturitas*, 125, 57-62. doi:10.1016/j.maturitas.2019.03.002.

Miller, J. M., Ashton - Miller, J. A., & DeLancey, J. O. L. (1998). A pelvic muscle precontraction can reduce Cough - Related urine loss in selected women with mild SUI. *Journal of the American Geriatrics Society*, 46 (7) , 870-874. doi:10.1111/j.1532-5415.1998.tb02721.x.

Moen, M., Noone, M., Vassallo, B., & Elser, D. (2009). Pelvic floor muscle function in women presenting with pelvic floor disorders. *International Urogynecology Journal*, 20 (7) , 843-846. doi:10.1007/s00192-009-0853-6.

Moosdorff-Steinhauser, H. F. A., Albers-Heitner, P., Weemhoff, M., Spaanderman, M. E. A., Nieman, F. H. M., & Berghmans, B. (2015). Factors influencing postpartum women's willingness to participate in a preventive pelvic floor muscle training program: A web-based survey. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 195, 182-187. doi:10.1016/j.ejogrb.2015.10.009.

森恵美,高橋真理,工藤美子,堤治,定月みゆき,坂上明子&大月恵理子,他. (2016) 系統看護学講座 専門分野Ⅱ 母性看護学 2 母性看護学各論.第13版, 東京: 医学書院.

Morin, M., Dumoulin, C., Bourbonnais, D., Gravel, D., & Lemieux, M. (2004). Pelvic floor maximal strength using vaginal digital ass

- essment compared to dynamometric measurements. *Neurourology and Urodynamics*, 23 (4) , 336-341. doi:10.1002/nau.20021.
- Morin, M., Dumoulin, C., Gravel, D., Bourbonnais, D., & Lemieux, M. (2007). Reliability of speed of contraction and endurance dynamometric measurements of the pelvic floor musculature in stress incontinent parous women. *Neurourology and Urodynamics*, 26 (3) , 397-403. doi:10.1002/nau.20334.
- Mørkved, S., & Bø, K. (1997). The effect of postpartum pelvic floor muscle exercise in the prevention and treatment of urinary incontinence. *International Urogynecology Journal*, 8 (4) , 217-222. doi:10.1007/BF02765817.
- Mørkved, S., Salvesen, K. Å, Bø, K., & Eik-nes, S. (2004). Pelvic floor muscle strength and thickness in continent and incontinent nulliparous pregnant women. *International Urogynecology Journal*, 15(6), 38-9; discussion 390. doi://dx.doi.org/10.1007/s00192-004-1194-0.
- Mørkved, S., & Bø, K. (2014). Effect of pelvic floor muscle training during pregnancy and after childbirth on prevention and treatment of urinary incontinence: A systematic review. *British Journal of Sports Medicine*, 48 (4) , 299-310. doi:10.1136/bjsports-2012-091758.
- Moser, H., Luginbuehl, H., Baeyens, J., & Radlinger, L. (2019). Reliability and validity of pelvic floor muscle displacement measurements during voluntary contractions. *International Urogynecology Journal*, 1-8. doi:10.1007/s00192-019-03932-x.
- Neels, H., Wyndaele, J., & Vermandel, A. (2018). Common errors made in attempt to contract the pelvic floor muscles in women earl

y after delivery: A prospective observational study. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, doi:10.1016/J.EJOGRB.2017.11.019.

日本排尿機能学会 女性下部尿路症状診療ガイドライン作成委員会.(2013).女性下部尿路症状診療ガイドライン,リッヂヒルメディカル.National Cancer Institute:Theory at a glance;a guide for health promotion practice. NIH Publication, 2005.

Okamoto, M., Murayama, R., Haruna, M., Matsuzaki, M., Kozuma, S., Nakata, M., & Murashima, S. (2010). Evaluation of pelvic floor function by transabdominal ultrasound in postpartum women. *Journal of Medical Ultrasonics*, 37 (4) , 187-193. doi:10.1007/s10396-010-0271-x.

Peng, Q., Jones. R., Shishido, K., & Constantinou, C. E. (2007). Ultrasound evaluation of dynamic responses of female pelvic floor muscles. *Ultrasound in Medicine & Biology*, 33 (3), 342-352. doi:10.1016/j.ultrasmedbio.2006.08.020.

Riesco, M. L. G., Caroci, A. d. S., Vasconcellos, de Oliveira, Sonia Maria Junqueira, & Lopes, de Moraes Maria Helena Baena. (2010). Perineal muscle strength during pregnancy and postpartum: The correlation between perineometry and digital vaginal palpation. *Revista Latino-Americana De Enfermagem*, 18(6), 1138-1144. doi:10.1590/S0104-11692010000600014.

Sacomori, C., Cardoso, F. L., Porto, I. P., & Negri, N. B. (2013). The development and psychometric evaluation of a self-efficacy scale for practicing pelvic floor exercises. *Brazilian Journal of Physical Therapy*, 17 (4) , 336-342. doi:10.1590/S1413-35552013005000104.

佐々木涼子, 若田真実, 矢花由佳, 百瀬公人, 平林直樹, 山ノ井万里子, & 西澤理. (2018) .尿失禁患者に対する骨盤底筋訓練の効果について－排尿日誌、骨盤底筋群筋力、ICIQ-SFによる検討－. 日本女性骨盤底医学会誌, 15 (1) ,100-102.

Sherburn, M., Murphy, C. A., Carroll, S., Allen, T. J., & Galea, M. P. (2005). Investigation of transabdominal real-time ultrasound to visualise the muscles of the pelvic floor.*Australian Journal of Physiotherapy*, 51 (3) , 167-170. doi:10.1016/S0004-9514(05)70023-4.

重田美和 (2017). 思春期女性の腹圧性尿失禁に対するアプローチ . 日本女性骨盤底医学会誌,14 (1) ,65-68.

高岡智子,谷口珠実, & 小林康江. (2016) .妊娠分娩による肛門挙筋の変化と評価.山梨県母性衛生学会誌,15,16-22.

高岡智子,近藤好枝,小林康江, & 谷口珠実. (2017) .産後の下部尿路症状と QOLとの関連性 包括的尺度 SF-12ver.2 を用いた検討. 日本助産学会誌, 31 (1), 78-87.

竹内京子, & 岡橋優子.(2015).見るみるわかる骨盤ナビ,pp50-57,東京:有限会社ラウンドフラット .

Talasz, H., Himmer-Perschak, G., Marth, E., Fischer-Colbrie, J., Hoefner, E., & Lechleitner, M. (2008). Evaluation of pelvic floor muscle function in a random group of adult women in austria.*International Urogynecology Journal*, 19 (1) , 131-135. doi:10.1007/s00192-007-0404-y.

谷口珠実, & 武田正之編著. (2017).下部尿路機能障害の治療とケア, 東京:MCメディア出版.

Thom, D. H., & Rortveit, G. (2010). Prevalence of postpartum urinary incontinence: A systematic review.*Acta Obstetricia Et Gynecologica Scandinavica*

*gica Scandinavica, 89* (12) , 1511-1522. doi:10.3109/00016349.2010.526188.

Thompson, J., & O'Sullivan, P. (2003). Levator plate movement during voluntary pelvic floor muscle contraction in subjects with incontinence and prolapse: A cross-sectional study and review. *International Urogynecology Journal, 14* (2), 84-88. doi:10.1007/s00192-003-1036-5.

Thompson, J., O'Sullivan, P., Briffa, K., Neumann, P., & Court, S. (2005). Assessment of pelvic floor movement using transabdominal and transperineal ultrasound. *International Urogynecology Journal, 16* (4) , 285-292. doi:10.1007/s00192-005-1308-3.

Thompson, J., O'Sullivan, P., Briffa, N., & Neumann, P. (2006). Assessment of voluntary pelvic floor muscle contraction in continent and incontinent women using transperineal ultrasound, manual muscle testing and vaginal squeeze pressure measurements. *International Urogynecology Journal, 17* (6) , 624-630. doi:10.1007/s00192-006-0081-2.

Thompson, J., O'Sullivan, P., Briffa, N., & Neumann, P. (2007). Comparison of transperineal and transabdominal ultrasound in the assessment of voluntary pelvic floor muscle contractions and functional manoeuvres in continent and incontinent women. *International Urogynecology Journal, 18*(7) , 779-786. doi:10.1007/s00192-006-0225-4.

辻野和美, 星野聰子, & 竹山政美. (2017) .骨盤底筋体操教室による筋収縮感覚の獲得と主体的取り組みへの可能性. 日本女性骨盤底医学会誌, 14 (1) , 74-80.

Timur-Taşhan, S., Beji, N. K., Aslan, E., & Yalçın, Ö. (2012).

- Determining lower urinary tract symptoms and associated risk factors in young women. *International Journal of Gynecology & Obstetrics*, 118 (1), 27-30. doi:10.1016/j.ijgo.2012.01.021
- 東條光彦, & 坂野雄二.(2003).セルフ・エフィカシー尺度.上里一郎監修,Hand book of Phycho logical Assessment 第 2 版, 425-433, 東京: 西村書店.
- Tsai, Y., & Liu, C. (2009). The effectiveness of pelvic floor exercises, digital vaginal palpation and interpersonal support on stress urinary incontinence: An experimental study. *International Journal of Nursing Studies*, 46 (9), 1181-1186. doi:10.1016/j.ijnurstu.2009.03.003
- van Delft, K., Thakar, R., & Sultan, A. H. (2015). Pelvic floor muscle contractility: Digital assessment vs transperineal ultrasound. *Ultrasound in Obstetrics & Gynecology*, 45 (2), 217-222. doi:10.1002/uog.13456.
- van Veelen, G. A., Schweitzer, K. J., & van der Vaart, C. H. (2014). Ultrasound imaging of the pelvic floor: Changes in anatomy during and after first pregnancy. *Ultrasound in Obstetrics & Gynecology*, 44 (4), 476-480. doi:10.1002/uog.13301.
- Vermandel, A., De Wachter, S., Beyltjens, T., D'Hondt, D., Jacquemyn, Y., & Wyndaele, J. (2015). Pelvic floor awareness and the positive effect of verbal instructions in 958 women early postdelivery. *International Urogynecology Journal*, 26 (2), 223-228. doi:10.1007/s00192-014-2483-x.
- Wesnes, S. L., & Lose, G. (2013). Preventing urinary incontinence during pregnancy and postpartum: A review. *International Urogynecology Journal*, 24 (6), 889. doi:10.1007/s00192-012-2017-3.

- Whittaker, J. L., Thompson, J. A., Teyhen, D. S., & Hodges, P. (2007). Rehabilitative ultrasound imaging of pelvic floor muscle function. *The Journal of Orthopaedic and Sports Physical Therapy*, 37(8), 487-498. doi:10.2519/jospt.2007.2548.
- Williams, K. S., Assassa, R. P., Gillies, C. L., Abrams, K. R., Turner, D. A., Shaw, C., ... McGrother, C. W. (2006). A randomized controlled trial of the effectiveness of pelvic floor therapies for urodynamic stress and mixed incontinence. *BJU International*, 98 (5), 1043-1050. doi:10.1111/j.1464-410X.2006.06484.x.
- 山崎喜比古,戸ヶ里泰典, & 坂野純子. (2012) .ストレス対処能力 SOC.東京:有信堂.
- Yoshida, M., Murayama, R., Hotta, K., Higuchi, Y., & Sanada, H. (2017). Differences in motor learning of pelvic floor muscle contraction between women with and without stress urinary incontinence: Evaluation by transabdominal ultrasonography. *Neurology and Urodynamics*, 36 (1), 98-103. doi:10.1002/nau.22867.