

References

Adewuya, A. (2006). Early postpartum mood as a risk factor for postnatal depression in Nigerian women. *American Journal Psychiatry*, 163(8), 1435-1437.

American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Washington, DC: American Psychiatric Publishing; 2013.

Amico, J. A., Tenicela, R. Johnston J. & Robinson, A. G. (1983). A time-dependent peak of oxytocin exists in cerebrospinal fluid but not in plasma of humans. *Journal of Clinical Endocrinology & Metabolism*. 57(5), 947-951.

Amico, J. A., Tenicela, R., Robinson, A. G. (1984). Studies of oxytocin in plasma of women during hypocontractile labor. *Journal Clinical Endocrinology Metabolism* 58, 274-279.

Arrindell WA, Gerlsma C, Vandereycken W, Hageman WJJM, Daeseleire T. (1998). Convergent validity of the dimensions underlying the Parental Bonding Instrument (PBI) and the EMBU. *Personality and Individual Differences*. 24(3), 341–350

Anim-Somuah, M. Smyth, R. M. & Jones, L. (2011). Epidural versus non-epidural or no analgesia in labour. *The Cochrane Database of Systematic Reviews* (12), doi:10.1002/14651858.CD000331.pub3

Anim - Somuah, M., Smyth, R., Cyna A. M., Cuthbert, A. (2018). Epidural versus non - epidural or no analgesia for pain management in labour. *Cochrane Database of Systematic Reviews*, 5. doi: 10.1002/14651858.CD000331.pub4.

Badr, H. A., & Zauszniewski, J. A. (2017). Kangaroo care and postpartum depression: The role of oxytocin. *International Journal of Nursing Sciences*, 4(2), 179-183. doi:10.1016/j.ijnss.2017.01.001

Bagherzadeh, R., Zahmatkeshan, N., Moatamed, N., Khorramroudi, R. & Ganjoo, M. (2009). Prevalence of maternal blues, postpartum depression and their correlation with premenstrual syndrome in women referred to health centers affiliated to Bushehr University of Medical Science. *Iranian Journal of Obstetrics, Gynecology and Infertility*, 12(3), 9-15.

Bell, A. F., Carter, C. S., Steer, C. D., Golding, J., Davis, J. M., Steffen, A. D., . . .

Connelly, J. J. (2015). Interaction between oxytocin receptor DNA methylation and genotype is associated with risk of postpartum depression in women without

depression in pregnancy. *Frontiers in Genetics*, 6, 243.

doi:10.3389/fgene.2015.00243

Bicking K. C. & Hupcey, J. E. (2013). State of the science of maternal-infant bonding:

A principle-based concept analysis. *Midwifery*, 29(12), 1314-1320.

Bienfait, M., Maury, M., Haquet, A., Faillie, J. L., Franc, N., Combes, C., . . .

Cambonie, G. (2011). Pertinence of the self-report mother-to-infant bonding scale

in the neonatal unit of a maternity ward. *Early Human Development*, 87(4), 281-

287. doi:10.1016/j.earlhumdev.2011.01.031

Bozoky, I., & Corwin, E. J. (2002). Fatigue as a predictor of postpartum depression.

Journal Obstetrics Gynecology Neonatal Nursing, 31(4), 436-443.

Brockington, I. (2004). Postpartum psychiatric disorders. *Lancet*, 363(9405), 303-310.

doi:S0140-6736(03)15390-1

Brockington, I. (2011). Maternal rejection of the young child: Present status of the

clinical syndrome. *Psychopathology*, 44(5), 329-336. doi:10.1159/000325058

Buckley, S. J. (2015). Executive summary of hormonal physiology of childbearing:

Evidence and implications for women, babies, and maternity care. *Journal of Perinatal Education*, 24(3), 145-153. doi:10.1891/1058-1243.24.3.145

Buttner, M., Mott, B., Pearlstein, T., Stuart, S., Zlotnick, C., & O'Hara, N. M. (2013).

Examination of premenstrual symptoms as a risk factor for depression in postpartum women. *Archives Womens Mental Health*, 16(3), 219-225.

Carson, D. S., Berquist, S. W., Trujillo, T. H., Garner, J. P., Hannah, S. L., Hyde, S.

A., . . . Parker, K. J. (2015). Cerebrospinal fluid and plasma oxytocin concentrations are positively correlated and negatively predict anxiety in children. *Molecular Psychiatry*, 20(9), 1085-1090. doi:10.1038/mp.2014.132

Choi, D., Minote, N., & Watanuki, S. (2017). Associations between the oxytocin

receptor gene (OXTR) rs53576 polymorphism and emotional processing of social and nonsocial cues: An event-related potential (ERP) study. *Journal Physiological Anthropology*, 36(1), doi:10.1186/s40101-016-0125-3

Christensson, K., Nilsson, B. A., Stock, S., Matthiesen, A. S., & Uvnas-Moberg, K.

(1989). Effect of nipple stimulation on uterine activity and on plasma levels of

oxytocin in full term healthy pregnant women. *Acta Obstetricia Et Gynecologica Scandinavica*, 68(3), 205-210.

Corwin, E. J., Brownstead, J., Barton, N., Heckard, S., & Morin, K. (2005). The impact of fatigue on the development of postpartum depression. *Journal Obstetrics Gynecology Neonatal Nursing*, 34(5), 577-586.

Cox, J., & Holden, J.(2003./ O. Okanao & S. Soda, trans.2006). Postpartum depression guidebook-to use EPDS-. Tokyo: Nanzan-do.

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry*, 150, 782-786.

Dalsant A, Truzzi A, Setoh P, Esposito G.(2015). Maternal bonding in childhood moderates autonomic responses to distress stimuli in adult males. *Behavioral Brain Research*. 292, 428–431.

Daughters, K., Manstead, A. S., Hubble, K., Rees, A., Thapar, A., & van Goozen, S. H.(2015). Salivary oxytocin concentrations in males following intranasal administration of oxytocin: A double-blind cross-over study. *PLoS One*, 10(12), doi:10.1371/journal.pone.0145104

Dawood, M. Y., Ylikorkata, O., Trivedi, D. & Gupta, R. (1980). Oxytocin levels and disappearance rate and plasma follicle-stimulating hormone and luteinizing hormone after oxytocin infusion in men. *Journal Clinical Endocrinology Metabolism*, 50(2), 397-400.

De Dreu, C. K., Greer, L. L., Van Kleef, G. A., Shalvi, S., & Handgraaf, M. J. (2011). Oxytocin promotes human ethnocentrism. *Proceedings of the National Academy of Sciences of the United States of America*, 108(4), 1262-1266.
doi:10.1073/pnas.1015316108

De Dreu, C. K., & Kret, M. E. (2016). Oxytocin conditions intergroup relations through upregulated in-group empathy, cooperation, conformity, and defense. *Biological Psychiatry*, 79(3), 165-173. doi:10.1016/j.biopsych.2015.03.020

De Geest, K., Thiery, M., Piron-Possuyt, G., & Vanden Driessche, R. (1985). Plasma oxytocin in human pregnancy and parturition. *Journal Perinatology Medicine*, (13)1, 3-13.

Ding T., Wang D. X, Qu Y., Chen, Q. & Zhu, S. N. (2014). Epidural labor analgesia is associated with a decreased risk of postpartum: A prospective cohort study. *Anesthesia & Analgesia*, 119(2), 383-392.

- Ditzen, B., Nater, U. M., Schaer, M., La Marca, R., Bodenmann, G., Ehlert, U., & Heinrichs, M. (2013). Sex-specific effects of intranasal oxytocin on autonomic nervous system and emotional responses to couple conflict. *Social Cognitive and Affective Neuroscience*, 8(8), 897-902. doi:10.1093/scan/nss083
- Dubber, S., Reck, C., Muller, M. & Gawlik, S. (2015). Postpartum bonding: The role of perinatal depression, anxiety and maternal-fetal bonding during pregnancy. *Archives of Women's Mental Health*, 18(2), 187-195. doi:10.1007/s00737-014-0445-4
- Edhborg, M., Matthiesen, A. S., Lundh, W., & Widström, A. M. (2005). Some early indicators for depressive symptoms and bonding 2 months postpartum: A study of new mothers and fathers. *Archives of Women's Mental Health*, 8, 221-231.
- Edhborg, M., Nasreen, H. E., & Kabir, Z. N. (2011). Impact of postpartum depressive and anxiety symptoms on mothers' emotional tie to their infants 2-3 months postpartum: A population-based study from rural Bangladesh. *Archives of Women's Mental Health*, 14, 307-316.
- Edhborg, M., Matthiesen, A. S., Lundh, W., & Widström, A. M. (2005). Some early indicators for depressive symptoms and bonding 2 months postpartum: A study of

and paternal care are associated with systematic changes in oxytocin following parent-infant contact. *Psychoneuroendocrinology*, 35(8), 1133-1141.

doi:10.1016/j.psyneuen.2010.01.013

Feldman, R., Eidelman, A. I., Sirota, L. & Weller, A. (2002). Comparison of skin-to-skin (kangaroo) and traditional care: Parenting outcomes and preterm infant development. *Pediatrics*, 110, 16-26.

Feldman, R., Gordon, I., & Zagoory-Sharon, O. (2011). Maternal and paternal plasma, salivary, and urinary oxytocin and parent-infant synchrony: Considering stress and affiliation components of human bonding. *Developmental Science*, 14(4), 752-761.
doi:10.1111/j.1467-7687.2010.01021.x

Feldman, R., Gordon, I. & Zagoory-Sharon, O. (2010). The cross-generation transmission of oxytocin in humans. *Hormones and Behavior*, 58(4), 669-676.

Feldman, R., Weller, A., Zagoory-Sharon, O. & Levine, A. (2007). Evidence for a neuroendocrinological foundation of human affiliation: Plasma oxytocin levels across pregnancy and the postpartum period predict mother-infant bonding. *Psychological Science*, 18(11), 965-970. doi:PSCI2010

- Feldman, R., Zagoory-Sharon, O., Weisman, O., Schneiderman, I., Gordon, I., Maoz, R., . . . Ebstein, R. P. (2012). Sensitive parenting is associated with plasma oxytocin and polymorphisms in the OXTR and CD38 genes. *Biological Psychiatry*, 72(3), 175-181.
- Figueiredo, B., Costa, R., Pacheco, A., & Pais, A. (2009). Mother-to-infant emotional involvement at birth. *Maternal and Child Health Journal*, 13, 539-549.
- Fuchs, A. R., Behrens, O., & Liu, H. C. (1992). Correlation of nocturnal increase in plasma oxytocin with a decrease in plasma estradiol/progesterone ratio in late pregnancy. *American Journal of Obstetrics and Gynecology*, 167(6), 1559-1563.
- Fuchs, A.R, & Fuchs, F. (1984). Endocrinology of human parturition: A review. *Br J Obstetrics and Gynecology*, 91(10), 948-967.
- Fuchs, A. R., Romero, R., Keefe, D., Parra, M., Oyarzun, E., & Behnke, E. (1991). Oxytocin secretion and human parturition: Pulse frequency and duration increase during spontaneous labor in women. *American Journal of Obstetrics and Gynecology*, 165(5 Pt 1), 1515-1523.
- Furman, L., & Kennell, J. (2000). Breastmilk and skin-to-skin kangaroo care for

premature infants: Avoiding bonding failure. *Acta Paeditrica*, 86, 1280-1283.

Giallo, R., Gartland, D., Woolhouse, H., & Brown, S. (2015). Differentiating maternal fatigue and depressive symptoms at six months and four years post partum: Considerations for assessment, diagnosis and intervention. *Midwifery*, 31(2), 316-322. doi:10.1016/j.midw.2014.09.005

Goebel-Stengel, M., Stengel, A., Tache, Y., & Reeve, J. R.. (2011). The importance of using the optimal plasticware and glassware in studies involving peptides. *Analytical Biochemistry*, 414(1), 38-46. doi:10.1016/j.ab.2011.02.009

Gomenz, A., Martinez O., Torrecillas J.(2015).Retrospective study of the association between epidural analgesia during labour and complications for the newborn.*Midwifery*, 31, 613-616.8

Gonidakis, F., Rabavilas, A.D., Varsou, E., Kreatsas, G., Christodoulou, GN.(2007). Maternity blues in Athens, Greece: a study during the first 3 days after delivery.*Journal of Affective Disorders*, 99(1-3), 107-115.

Gordon I., Zagoory-Sharon, I., Leckman., Feldman, R.(2010).Oxytocin and the Development of Parenting in Humans.*Biol Psychiatry*, 68(4), 377-382.

Gordon Parker., Hilary Tupling., L. B. Brown. (1979). A parental bonding instrument.

British Journal of Medical Psychology, 52, 1-10.

G*POWER <http://www.gpower.hhu.de/> [access 2018/08/31]

Grewen, K. M., Davenport, R. E., & Light, K. C. (2010). An investigation of plasma and salivary oxytocin responses in breast- and formula-feeding mothers of infants.

Psychophysiology, 47(4), 625-632. doi:10.1111/j.1469-8986.2009.00968.x

Gu, V., Feeley, N., Gold, I., Hayton, B., Robins, S., Mackinnon, A., . . . Zelkowitz, P.

(2016). Intrapartum synthetic oxytocin and its effects on maternal well-being at 2 months postpartum. *Birth (Berkeley, Calif.)*, 43(1), 28-35. doi:10.1111/birt.12198

Guyton, A. C., & Hall, J. E. (2010). ガイトン生理学 [Textbook of medical physiology/ ガイトン生理学] (御手洗玄洋[総監訳], 小川徳雄, 永坂鉄夫, 伊藤嘉房, 松井信夫 & 間野忠明[監訳]). (11th ed.). 東京: エルゼビア・ジャパン.

Guzman, Y. F., Tronson, N. C., Jovasevic, V., Sato, K., Guedea, A. L., Mizukami,

H., . . . Radulovic, J. (2013). Fear-enhancing effects of septal oxytocin receptors.

Nature Neuroscience, 16(9), 1185-1187. doi:10.1038/nn.3465

Hall, R. A. S., Hoffenkamp, H. N., Tooten, A., Braeken, J., Vingerhoets, A. J., & Van

Bakel, H. J. A. (2014). Child-rearing history and emotional bonding in parents of preterm and full-term infants. *Journal of Child and Family Studies*, published online DOI 10.1007/s10826-014-9975-7

Hau F, Levy V. (2003). The maternity blues and Hong Kong Chinese women: an exploratory study. *Journal of Affect Disorders*, 75(2), 197-203.

Heinrichs, M., Baumgartner, T., Kirschbaum, C., & Ehlert, U. (2003). Social support and oxytocin interact to suppress cortisol and subjective responses to psychosocial stress. doi:[https://doi.org/10.1016/S0006-3223\(03\)00465-7](https://doi.org/10.1016/S0006-3223(03)00465-7)

Henderson, J., Alderdice, F., & Redshaw, M. (2019). Factors associated with maternal postpartum fatigue: An observational study. *BMJ Open*, 9(7), e025927-2018-025927. doi:10.1136/bmjopen-2018-025927

Hendrick, V., Altshuler, L., Cohen, L, Stowe, Z. (1998). Evaluation of mental health and depression during pregnancy: position paper. *Psychopharmacol Bull*, 34(3), 297-299.

Hsieh, C. H., Chen, C. L., Han, T. J., Lin, P. J., & Chiu, H. C. (2018). Factors influencing postpartum fatigue in vaginal-birth women: Testing a path model. *The*

Journal of Nursing Research : JNR, 26(5), 332-339.

doi:10.1097/jnr.0000000000000249

Herrera-Gomez, A., Garcia-Martinez, O., Ramos-Torrecillas, J., De Luna-Bertos, E.,

Ruiz, C., & Ocana-Peinado, F. M. (2015). Retrospective study of the association between epidural analgesia during labour and complications for the newborn.

Midwifery, 31(6), 613-616. doi:10.1016/j.midw.2015.02.013

Hoffman, E. R., Brownley, K. A., Hamer, R. M., & Bulik, C. M. (2012). Plasma,

salivary, and urinary oxytocin in anorexia nervosa: A pilot study. *Eating*

Behaviors, 13(3), 256-259. doi:10.1016/j.eatbeh.2012.02.004

Holt-Lunstad, J., Birmingham, W., & Light, K. C. (2011). The influence of depressive

symptomatology and perceived stress on plasma and salivary oxytocin before,

during and after a support enhancement intervention. *Psychoneuroendocrinology*,

36(8), 1249-1256. doi:10.1016/j.psyneuen.2011.03.007

Iida, M., Horiuchi, S., Nagamori, K. (2014). A comparison of midwife-led care versus

obstetrician-led care for low-risk women in Japan. *Women and Birth*, 27, 202-207.

Ishikawa, N.,Goto, S.,Murase, S.,Knai, A.,Masuda, T.,Aleksic, B.,Usui, H.,Ozaki,

N.(2011).Prospective study of maternal depressive symptomatology among

Japanese women. *Journal of Psychosomatic Research*,.71, 264-269.

Iwata, H.,Mori, E.,Sakajo, A.,Aoki, K.,Maehara, K.,Tamakoshi, K.(2016).Prevalence of

postpartum depressive symptoms during the first 6 months postpartum: Association

with maternal age and parity. *Journal of Affective disorders*, 203, 227-232.

Japan Society for obstetric Anesthesia and Perinatology

<https://www.jsoup.com/about/overview> [access 2020/01/05]

Javor, A., Riedl, R., Kindermann, H., Brandstatter, W., Ransmayr, G., & Gabriel, M.

(2014). Correlation of plasma and salivary oxytocin in healthy young men -

experimental evidence. *Neuro Endocrinology Letters*, 35(6), 470-473.

doi:NEL350614A04

Jobst, A.,Krause, D.,Maiwald, C.,Hartl, K.,Myint, A.,Kastner, R.(2016).Oxytocin

course over pregnancy and postpartum period and the association with postpartum

depressive symptoms. *Arch Womens Mental Health*, 19, 571-579.

Jomeen, J.(2004). The importance of assessing psychological status during

pregnancy, childbirth and the postnatal period as a multidimensional construct :

A literature review. *Clinical Effectiveness in Nursing*, 8, 143-155.

Jonas, K.,Johansson, LM.,Nissen, E.,Ransjo-Arvidson, AB.,Uvnas-Moberg, K.(2009).

Effects of intrapartum oxytocin administration and epidural analgesia on the

concentration of plasma oxytocin and prolactin, in response to suckling during

the second day postpartum. *Breastfeeding Medicine*, 4(2), 71-82.

Jong, T. R., Menon, R., Bludau, A., Grund, T., Biermeier, V., Klampfl, S. M., . . .

Neumann, I. D. (2015). Salivary oxytocin concentrations in response to running,

sexual self-stimulation, breastfeeding and the TSST: The regensburg oxytocin

challenge (ROC) study. *Psychoneuroendocrinology*, 62, 81-388.

doi:10.1016/j.psyneuen.2015.08.027

Kendall K., Cong Z., Hale T. (2015). Birth Interventions Related to Lower Rates of

Exclusive Breastfeeding and Increased Risk of Postpartum Depression in a Large

Sample. *clinical Lactation*, (3), 87-97.

Kitamura, T., Ohashi, Y., Kita, S., Haruna, M., Kubo, R. (2013). Depressive mood,

bonding failure, and abusive parenting among mothers with three-month-old

babies in a Japanese community. *Open Journal of Psychiatry*, 3, 1-7.

北村俊則, 南谷真理子, 大橋優紀子.(2017).産後うつ病・児童虐待・嬰兒殺に関

する意見. www.institute-of-mental-health.jp/pdf/201403.pdf

[access 2018/08/01]

Kitamura, T., Yoshida, K., Okano, T., Kinoshita, K., Hayashi, M., Toyoda, N., Ito, M.,

Kudo, N., Tada, K., Kanazawa, K., Sakumoto, K., Satoh, S., Furukawa, T., &

Nakano, H. (2006). Multicentre prospective study of perinatal depression in Japan:

Incidence and correlates. *Archives of Women's Mental Health*, 9, 121-130.

Kosaka, H., Okamoto, Y., Munesue, T., Yamasue, H., Inohara, K., Fujioka, T., . . .

Wada, Y. (2016). Oxytocin efficacy is modulated by dosage and oxytocin receptor

genotype in young adults with high-functioning autism: A 24-week randomized

clinical trial. *Translational Psychiatry*, 6(8), e872. doi:10.1038/tp.2016.152

Kokubu M., Okano, T., & Sugiyama, T. (2012). Postnatal depression, maternal

bonding failure, and negative attitudes towards pregnancy: a longitudinal study

of pregnant women in Japan. *Archives of Women's Mental Health*, 15, 211–

216.

Kumar, R.C.(1997).”Anybody’s child”: severe disorders of mother-to-infant bonding.*Br*

J Psychiatry, 171,175-181.

Kumsta, R., Hummel, E., Chen, F. S., & Heinrichs, M. (2013). Epigenetic regulation of the oxytocin receptor gene: Implications for behavioral neuroscience. *Frontiers in Neuroscience*, 7, 83. doi:10.3389/fnins.2013.00083

Krehbiel, D., Poindron, P., Levy, F., & Prud'Homme, M. J. (1987). Peridural anesthesia disturbs maternal behavior in primiparous and multiparous parturient ewes. *Physiology & Behavior*, 40(4), 463-472. doi:0031-9384(87)90031-X

Lee, H. J., Macbeth, A. H., Pagani, J. H., & Young, W. S. (2009). Oxytocin: The great facilitator of life. *Progress in Neurobiology*, 88(2), 127-151. doi:10.1016/j.pneurobio.2009.04.001

Lee, Y., Wook, S., Ju, D., Kim I. (2015). Correlation between postpartum depression and premenstrual dysphoric disorder: Single center study. *Obstetrics & Gynecology Science*, 58(5). 353-358.

Levy, F., Kendrick, K. M., Keverne, E. B., Piketty, V., & Poindron, P. (1992). Intracerebral oxytocin is important for the onset of maternal behavior in inexperienced ewes delivered under peridural anesthesia. *Behavioral Neuroscience*, 106(2), 427-432.

Light K C., Grewen K M., & Amico J A. (2005). More frequent partner hugs and higher oxytocin levels are linked to lower blood pressure and heart rate in premenopausal women. *Biological Psychology*, 69(1), 5-21.

Lindow, S.W, Hendricks, Van. (1996). The 24-hour rhythm of oxytocin and B-endorphin secretion in human pregnancy. *Clinical Endocrinology*, 45(4), 443-446.

升谷泰裕, 藤澤隆史, 浅野みずき. (2013). 血液検体と唾液検体によるオキシトシン濃度測定値の相関性について. 日本精神医学会総会抄録集54th.

丸山陽子, 川崎佳代子, 竹尾恵子, 金城壽子, 弓削美鈴. (2012). 産褥期うつスクリーニングと背景要因の検討. 佐久大学看護研究雑誌4巻1号, 15-27.

Meades, R., & Ayers, S. (2011). Anxiety measures validated in perinatal populations: A systematic review. *Journal of Affective Disorders*, 133(1-2), 1-15.
doi:10.1016/j.jad.2010.10.009

Mileva-Seitz, V., Steiner, M., Atkinson, L., Meaney, M. J., Levitan, R., Kennedy, J. L.; Sokolowski, M. B.; Fleming, A. S. (2013). Interaction between oxytocin genotypes and early experience predicts quality of mothering and postpartum mood. *PLoS One*, 8(4), e61443.

Morhenn V., Beavin L. E., Zak P.(2012). Massage increases oxytocin and reduce adrenocorticotropin hormone in humans. *ALTERNATIVE THERAPIES*, 18(6), 11-18.

Muchanga S,Yasumitsu K,Eitoku M,Mbelambela E,Ninomiya H,Komori K,Tozin R.(2017).Preconception gynecological risk factors of postpartum depression among Japanese women: The Japan Environment and Children's Study.*Journal of affective Disorders*, 217. 34-41.

Muzik, M., Bocknek, E. L., Broderick, A., Richardson, P., Rosenblum, K. L., Thelen, K., & Seng, J. S. (2013). Mother-infant bonding impairment across the first 6 months postpartum: The primacy of psychopathology in women with childhood abuse and neglect histories. *Archives of Women's Mental Health*, 16, 29-38.

Uvnäs-Moberg, K., Ekström-Bergström, A., Berg, M., Buckley, S., Pajalic, Z., Hadjigeorgiou, E., ... Dencker, A. (2019). Maternal plasma levels of oxytocin during physiological childbirth - a systematic review with implications for uterine contractions and central actions of oxytocin. *BMC pregnancy and childbirth*, 19(1), 285. doi:10.1186/s12884-019-2365-9

Matta, P., Turner, J., Flatley, C., & Kumar, S. (2019). Prolonged second stage of labour increases maternal morbidity but not neonatal morbidity. *The Australian & New Zealand Journal of Obstetrics & Gynaecology*, 59(4), 555-560.
doi:10.1111/ajo.12935

Mörelus, E., Theodorsson, E., & Nelson, N. (2005). Salivary cortisol and mood and pain profiles during skin-to-skin care for an unselected group of mothers and infants in neonatal intensive care. *Pediatrics*, 116, 1105-1113.

Mori, E., Tsuchiya, M., Maehara, K., Iwata, H., Sakajo, A., & Tamakoshi, K. (2017). Fatigue, depression, maternal confidence, and maternal satisfaction during the first month postpartum: A comparison of Japanese mothers by age and parity. *International Journal of Nursing Practice*, 23(1), 10.1111/ijn.12508. Epub 2017 Jan 8. doi:10.1111/ijn.12508

中里克治, 水口公信. (1982). 新しい不安尺度 STAI 日本語版の作成. *心身医*, 22, 107-112.

Nitta, M., Narita, T., Umeda, K., Hattori, M., Naitoh, H., & Iwata, N. (2008). Influence of negative cognition on the parental bonding instrument (PBI) in patients with

major depression. *Journal of Nervous and Mental Disease*, 196(3), 244-

246. <https://doi.org/10.1097/NMD.0b013e318165c116>

Niwayama, R., Nishitani, S., Takamura, T., Shinohara, K., Honda, S., Miyamura, T., . . .

Araki-Nagahashi, M. (2017). Oxytocin mediates a calming effect on postpartum

mood in primiparous mothers. *Breastfeeding Medicine : The Official Journal of the*

Academy of Breastfeeding Medicine, 12, 103-109. doi:10.1089/bfm.2016.0052

Norwitz, E. R., Mahendroo, M., & Lye, S. J. (2014). Biology of parturition. In R. K.

Creasy, Creasy and resnik's maternal-fetal medicine : Principles and practice (7th

ed.,pp. 66-79) Saunders/Elsevier.

E. Nissen, P. Gustavsson, A-M. Widström & K. Uvnäs-Moberg (1998) Oxytocin,

prolactin, milk production and their relationship with personality traits in women

after vaginal delivery or Cesarean section, *Journal of Psychosomatic Obstetrics &*

Gynecology, 19(1), 49-58, DOI: [10.3109/01674829809044221](https://doi.org/10.3109/01674829809044221)

Ohashi, Y., Kitamura, T., Sakanashi, K., & Tanaka, T. (2015). Psychometric properties

of the Puerperal Bonding Questionnaire in Japanese mothers of infants.

O'Hara, Schlechte J, Lewis D, Wright E. (1991). Prospective study of postpartum blues.

Biologic and psychosocial factors. *Archive of General Psychiatry*, 48(9), 801-806.

O'Hara, M. W., & McCabe, J. E. (2013). Postpartum depression: Current status and future directions. *Annual Review of Clinical Psychology*, 9, 379-407. doi:10.1146/annurev-clinpsy-050212-185612

岡野禎治, 野村純一, 越川法子.(1991). Maternity Bluesと産後うつ病の比較文化的研究. *精神医学*, 33, 1051-1058.

岡野禎治, 村田真理子, 増地総子.(1996).日本版エジンバラ産後うつ病評価票(EPDS)の信頼性と妥当性. *精神科診断学*, 7(4), 525-533.

小川 雅美. (1991). PBI (parental bonding instrument)日本版の信頼性, 妥当性に関する研究. *精神科治療学*, 6(10), 1193-1201.

尾仲 達史, 吉田 匡秀, & 高柳 友紀. (2015). 【メンタルストレスの基礎と臨床】不安・恐怖とオキシトシン. *アンチ・エイジング医学*, 11(1), 024-033.

Parker, G., Tupling, H., & Brown, L. B. (1979). A parental bonding instrument. *British Journal of Medical Psychology*. 52(1), 1-10.

Prevost, M., Zelkowitz, P., Tulandi, T., Hayton, B., Feeley, N., Carter, C. S., . . . Gold,

I. (2014). Oxytocin in pregnancy and the postpartum: Relations to labor and its management. *Frontiers in Public Health*, 2, 1. doi:10.3389/fpubh.2014.00001

Radloff, L. S. (1977). The CES-D scale: A self report depression scale for research in the general population. *Applied Psychological Measurements*, 1, 385-401.

Rahm, V. A., Hallgren, A., Hogberg, H., Hurtig, I., & Odland, V. (2002). Plasma oxytocin levels in women during labor with or without epidural analgesia: A prospective study. *Acta Obstetrica Et Gynecologica Scandinavica*, 81(11), 1033-1039. doi:aog811107

Rapaport, M. H., Schettler, P., & Bresee, C. (2012). A preliminary study of the effects of repeated massage on hypothalamic-pituitary-adrenal and immune function in healthy individuals: A study of mechanisms of action and dosage. *Journal of Alternative and Complementary Medicine (New York, N.Y.)*, 18(8), 789-797. doi:10.1089/acm.2011.0071

Reck C, Stehle E, Reinig K, Mundt C. (2009). Maternity blues as a predictor of DSM-IV depression and anxiety disorders in the first three months postpartum. *Journal of Affective Disorders*, 113, 77-87.

Rikhye K, Tyrka AR, Kelly MM, Gagne GG, Carpenter LL. (2008).Interplay between childhood maltreatment, parental bonding, and gender effects: impact on quality of life. *Child Abuse and Neglect.* 32, 19–34.

Rychnovsky, J. D. (2007). Postpartum fatigue in the active-duty military woman. *Journal of Obstetric, Gynecologic, and Neonatal Nursing : JOGNN*, 36(1), 38-46. doi:S0884-2175(15)33649-2

Safari K., Saeed AA., Hasan SS., Moghaddam-Banaem L.(2018). The effect of mother and newborn early skin-to-skin contact on initiation of breastfeeding, newborn temperature and duration of third stage of labor. *International Breastfeeding Journal*, 13(32). PMID: 30026787

Sakumoto, K.,Masamoto, H.,Kanazawa K.(2002).Post-partum maternity ‘blues’ as a reflection of newborn nursing care in Japan.*International Journal of Gynecology and Obstetrics*, 78, 25-30.

Salimetrics LCC. saliva collection and handling advice. (2015).

Scatliffe,N, Sharon Casavant, Dorothy Vittner, Xiaomei Cong.(2019).Oxytocin and early parent-infant interactions: A systematic review, *International Journal of Nursing Sciences*,6(4),445-453.

穴戸 恵理. (2016). 一出産体験における痛みと疲労の予測と現実のギャップー

無痛分娩と非無痛分娩の比較. 聖路加国際大学大学院修士論文

Shishido, E., Shuo, T., Takahata, K., & Horiuchi, S. (2019). Changes in salivary oxytocin levels and bonding disorder in women from late pregnancy to early postpartum: A pilot study. *PloS One*, 14(9), e0221821.

doi:10.1371/journal.pone.0221821

島 悟, 鹿野 達男, & 北村 俊則. (1985). 新しい抑うつ性自己評価尺度について. *精神医学*, 27(6), 717-723.

周産期メンタルヘルス コンセンサスガイド(2017). 日本周産期メンタルヘルス学会.

Skrundz, M., Bolten, M., Nast, I., Hellhammer, D. H., & Meinlschmidt, G. (2011).

Plasma oxytocin concentration during pregnancy is associated with development of postpartum depression. *Neuropsychopharmacology : Official Publication of the American College of Neuropsychopharmacology*, 36(9), 1886-1893.

doi:10.1038/npp.2011.74

Sockol, L. E., Battle, C. L., Howard, M., & Davis, T. (2014). Correlates of impaired mother-infant bonding in a partial hospital program for perinatal women. *Archives of Women's Mental Health*, 17, 465-469.

園田希,小川真世,田所由利子,高畑香織,周尾卓也,堀内成子.(2018).妊婦を対象としたふれて・感じる「Mama's Touch プログラム」の実行可能性ーオキシトシン・コルチゾールによる評価；予備研究ー. *日本助産学会誌*,32(1),pp.60-72.

Spielberger C.,Gorsuch L.,Lushene,R.E. (1970). Manual for the state trait anxiety inventory(Self-evaluation question-aire). *Consulting Psychologists*, 2-24.

Stein, G. S. (1980). The pattern of mental change and body weight change in the first post-partum week. *Journal of Psychosomatic Research*, 24(3-4), 165-171.
doi:0022-3999(80)90038-0

Stevens L. F., Wiesman O., Feldman R., Hurley A. R., & Taber, H K.(2013). Oxytocin and Behavior : Evidence for Effects in the Brain. *Journal of Neuropsychiatry and Clinical Neurosciences*, 25(2).

Stock S.,Bremme K.,Moberg K.(1991). Plasma levels of oxytocin during the menstrual cycle, pregnancy and following treatment with HMG. *Humen Reproduction*,6(8), 1056-1062.

Stuebe A.,Grewen K, Meltzer-Brody S.(2013). Association between maternal mood and oxytocin response to breastfeeding. *Journal of Women's Health*,22(4), 352-361.

Tadokoro, Y.(2016) .Effects of Clary Sage and Lavender or Jasmine Essential Oils Infused in a Foot Bath for Labor Induction on Oxytocin Level. (Graduate School of St. Luke's International University).

高橋徳.(2014) .人は愛することで健康になれる (愛のホルモン・オキシトシン).知道出版.

Tadokoro Y,Horiuchi S,Takahata K,Shuo T,Sawano E,Shinohara K.(2017).Changes in salivary oxytocin after inhalation of clary sage essential oil scent in term pregnant women : a feasibility pilot study. *BMC Research Notes*,10,717.

Takahashi Y,Tamakoshi K.(2014).Factors Associated with early Postpartum Maternity Blues and Depression Tendency Among Japanese Mothers with Full-term Healty ,Infants. *Nagoya Journal of Medical Science*,76, 129-138.

Takahata, K. (2016). Oxytocin levels in low-risk primiparas following breast stimulation for spontaneous onset of labor. (Unpublished Graduate School of St. Luke's International University).

Takahata K,Horiuchi S,Tadokoro Y,Shuo T,Sawano E,Shinohara K.(2018).Effects of breast stimulation for spontaneous onset of labor on salivary oxytocin levels in low-risk pregnant women : A feasibility study.*PLoS One*,13(2), e0192757

Takacs, L., Seidlerova, J. M., Sterbova, Z., Cepicky, P., & Havlicek, J. (2019). The effects of intrapartum synthetic oxytocin on maternal postpartum mood: Findings from a prospective observational study. *Archives of Women's Mental Health*, 22(4), 485-491. doi:10.1007/s00737-018-0913-3

Takeda, S., Kuwabara, Y., & Mizuno, M. (1989). Metabolic clearance rate of oxytocin in human subjects. *Nihon Naibunpi Gakkai Zasshi*, 65(3), 182-189.

Taylor, A., Atkins, R., Kumar, R., Adams, D., & Glover, V. (2005). A new mother-to-infant bonding scale: Links with early maternal mood. *Archives of Women's Mental Health*, 8(1), 45-51. doi:10.1007/s00737-005-0074-z

Terada, M., Matsuda, Y., Ogawa, M., Matsui, H., & Satoh, S. (2013). Effects of maternal factors on birth weight in japan. *Journal of Pregnancy*, 172395. doi:10.1155/2013/172395

Terkawi, A. S., Jackson, W. M., Thiet, M., Hansoti, S., Tabassum, R., & Flood, P.

(2012). Oxytocin and catechol-O-methyltransferase receptor genotype predict the length of the first stage of labor. *American Journal of Obstetrics and Gynecology*, 207(3), 184-184.

Tharner, A., Luijk, M. P., Raat, H., Ijzendoorn, M. H., Bakermans-Kranenburg, M. J., Moll, H.

A., Jaddoe, V. W., Hofman, A., Verhulst, F. C., Tiemeier, H. (2012). Breastfeeding and its relation to maternal sensitivity and infant attachment. *Journal of developmental and behavioral pediatrics : JDBP*, 5, 396-404.

Thornton, S., Davison, J. M., & Baylis, P. H. (1990). Effect of human pregnancy on metabolic clearance rate of oxytocin. *The American Journal of Physiology*, 259(1 Pt 2), 21-24.

Tooten, A., Hall, R. A., Hoffenkamp, H. N., Braeken, J., Vingerhoets, A. J., & van

Bakel, H. J. (2014). Maternal and paternal infant representations: A comparison

between parents of term and preterm infants. *Infant Behavior & Development*,

37(3), 366-379. doi:10.1016/j.infbeh.2014.05.004

Traut R, Watanabe K, Nazarloo P, Schwertz D, Bell A, Carter C. (2009). Detection of

salivary oxytocin levels in lactating women. *Developmental Psychobiology*. 51(4),

367-373.

Troy, N. W. (2003). Is the significance of postpartum fatigue being overlooked in the lives of women? *The American Journal of Maternal Child Nursing*, 28(4), 252-257.

Unternaehrer, E., Luers, P., Mill, J., Dempster, E., Meyer, A. H., Staehli, S., . . .

Meinlschmidt, G. (2012). Dynamic changes in DNA methylation of stress-associated genes (OXTR, BDNF) after acute psychosocial stress. *Translational Psychiatry*, 2, e150. doi:10.1038/tp.2012.77

Van Bussel, J. C. H., Spitz, B., & Demyttenaere, K. (2010). Three self-report questionnaires of the early mother-to-infant bond: Reliability and validity of the Dutch version of the MPAS, PBQ and MIBS. *Archives of Women's Mental Health*, 13, 373-384.

Vittner, D., McGrath, J., Robinson, J., Lawhon, G., Cusson, R., Eisenfeld, L., . . . Cong, X. (2018). Increase in oxytocin from skin-to-skin contact enhances development of parent-infant relationship. *Biological Research for Nursing*, 20(1), 54-62. doi:10.1177/1099800417735633

Waldman-Levi, A., Finzi-Dottan, R., & Weintaub, N. (2015). Attachment security and parental perception of competency among abused women in the shadow of PTSD

and childhood exposure to domestic violence. *Journal of Child and Family Studies*, 24, 57-65.

Watanabe, M., Wada, K., Sakata, Y., Aratake, Y., Kato, N., Ohta, H., & Tanaka, K.

(2008). Maternity blues as predictor of postpartum depression: A prospective cohort study among Japanese women. *Journal of Psychosomatic Obstetrics and Gynaecology*, 29(3), 206-212. doi:10.1080/01674820801990577

Wilson, N., Lee, J. J., & Bei, B. (2019). Postpartum fatigue and depression: A systematic review and meta-analysis. *Journal of Affective Disorders*, 246, 224-233. doi:S0165-0327(18)31947-5

Wilson, N., Wynter, K., Anderson, C., Rajaratnam, S. M. W., Fisher, J., & Bei, B. (2019). Postpartum fatigue, daytime sleepiness, and psychomotor vigilance are modifiable through a brief residential early parenting program. *Sleep Medicine*, 59, 33-41. doi:S1389-9457(19)30012-7

Witteveen, A. B., Stramrood, C. A. I., Henrichs, J., Flanagan, J. C., van Pampus, M. G., & Olff, M. (2019). The oxytocinergic system in PTSD following traumatic childbirth: Endogenous and exogenous oxytocin in the peripartum period. *Archives of Women's Mental Health*, doi:10.1007/s00737-019-00994-0

Working Group for Occupational Fatigue. How to use

“Jikaku-sho shirabe”. <http://square.umin.ac.jp/of/service.html> (in Japanese).

Xu, M. K., Morin, A., Marsh, H. W., Richards, M., & Jones, P. B. (2018). Psychometric

Validation of the Parental Bonding Instrument in a U.K. Population-Based Sample:

Role of Gender and Association With Mental Health in Mid-Late

Life. *Assessment*, 25(6), 716–728. doi:10.1177/1073191116660813

山崎圭子, 東園子, 増田知実, 永嶋輝子.(2015).感覚距離等分法、SF-36、唾液コルチゾール

値を用いた産褥早期の母親の疲労感の評価.東邦看護学会誌,12, 7-12.

Yamashita H,Yoshida K,Nakano K,Tashiro N.(2000).Postnatal depression in Japanese

women detecting the early onset of postnatal depression by closely monitoring the

postpartum mood.*Journal of Affective Disorders*, 58, 145-154.

Yoshida, K., Yamashita, H., Conroy, S., Marks, M., & Kumar, C. (2012). A japanese

version of mother-to-infant bonding scale: Factor structure, longitudinal changes

and links with maternal mood during the early postnatal period in japanese

mothers. *Archives of Women's Mental Health*, 15(5), 343-352. doi:10.1007/s00737-

012-0291-1

Young, S. N., & Anderson, G. N. (2010). Bioanalytical inaccuracy: A threat to the integrity and efficiency of research. *Journal of Psychiatry & Neuroscience : JPN*, 35(1), 3-6.

Zanardo, V., Volpe, F., de Luca, F., Giliberti, L., Giustardi, A., Parotto, M., . . . Soldera, G. (2019). Maternity blues: A risk factor for anhedonia, anxiety, and depression components of edinburgh postnatal depression scale. *The Journal of Maternal-Fetal & Neonatal Medicine : The Official Journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstetricians*, , 1-7. doi:10.1080/14767058.2019.1593363

Zelkowitz P.,Gold I.,Feely N.,Hayton B.,Carter S.,Tulandi T.,Abenhaim H.,Levin P.(2014). Psychosocial stress moderates the relationships between oxytocin, perinatal depression, and maternal behavior. *Hormones and Behavior*, 66(2), 351-360.