

5. References

- 1) Kawasaki, T. Febrile oculo-oro-cutaneo-acrodesquamatus syndrome with or without acute non-suppurative cervical lymphadenitis in infancy and childhood: Clinical observations, of 50 cases. Jpn J Allergy 1967; 16: 178–222. [Japanese]
- 2) <https://www.jichi.ac.jp/dph/wp-dph/wp-content/uploads/2022/04/a19b047d4b9e6fbb84b6b187236779c8.pdf> [Japanese]
- 3) McCrindle BW, et al. Diagnosis, Treatment, and Long-Term Management of Kawasaki Disease: A Scientific Statement for Health Professionals From the American Heart Association. Circulation 2017; 135: e927–e999.
- 4) JCS/JSCS 2020 Guideline on Diagnosis and Management of Cardiovascular Sequelae in Kawasaki disease(Published in March 13th 2020)
- 5) Suzuki A, et al. Remodeling of coronary artery lesions due to Kawasaki disease: comparison of arteriographic and immunohistochemical findings. Jpn Heart J 2000; 41: 245–256.
- 6) Fukazawa R, Kobayashi T, Mikami M, et al. Nationwide Survey of Patients With Giant Coronary Aneurysm Secondary to Kawasaki Disease 1999-2010 in Japan. Circ J 2017; 82: 239–246.
- 7) Suda K, et al. Long-term prognosis of patients with Kawasaki disease complicated by giant coronary aneurysms: a single-institution experience. Circulation 2011; 123: 1836–1842.
- 8) Mitani Y, et al. Impaired endothelial function in epicardial coronary arteries after Kawasaki disease. Circulation 1997;96: 454–461.

- 9) Yamakawa R, et al. Coronary endothelial dysfunction after Kawasaki disease: evaluation by intracoronary injection of acetylcholine. *J Am Coll Cardiol* 1998; 31: 1074–1080.
- 10) Tobayama H, et al. Analysis of arterial function in adults with a history of Kawasaki disease. *J Cardiol* 2013;61: 330–335.
- 11) Tacke CE, et al. Cardiac magnetic resonance imaging for noninvasive assessment of cardiovascular disease during the follow-up of patients with Kawasaki disease. *Circ Cardiovasc Imaging* 2011; 4: 712–720.
- 12) Furuyama H, et al. Assessment of coronary function in children with a history of Kawasaki disease using ^{15}O -water positron emission tomography. *Circulation* 2002; 105: 2878–2884.
- 13) Muzik O, et al. Quantification of myocardial blood flow and flow reserve in children with a history of Kawasaki disease and normal coronary arteries using positron emission tomography. *J Am Coll Cardiol* 1996; 28: 757–762.
- 14) Matsuda H, et al. Long-term follow-up of a patient with Kawasaki disease and coronary aneurysm associated with asymptomatic thrombosis: a case report. *J Cardiol* 2005; 46: 113–118.
- 15) Kobayashi T, et al. Z Score Project Investigators. A New Z Score Curve of the Coronary Arterial Internal Diameter Using the Lambda-Mu-Sigma Method in a Pediatric Population. *J Am Soc Echocardiogr* 2016; 29: 794–801.e29.

- 16) Friedman KG, et al. Coronary Artery Aneurysms in Kawasaki Disease: Risk Factors for Progressive Disease and Adverse Cardiac Events in the US Population. *J Am Heart Assoc* 2016; 5: e003289.
- 17) Miura M, et al. The Z-score Project 2nd Stage Study Group. Association of Severity of Coronary Artery Aneurysms in Patients With Kawasaki Disease and Risk of Later Coronary Events. *JAMA Pediatr* 2018; 172: e180030.
- 18) Suzuki A, et al. Natural history of coronary arterial lesions in Kawasaki disease. *Prog Pediatr Cardiol* 1997; 6:211–218.
- 19) Singhal M, Singh S, Gupta P, et al. Computed Tomography Coronary Angiography for Evaluation of Children With Kawasaki Disease. *Curr Probl Diagn Radiol* 2018; 47: 238–244.
- 20) Mavrogeni S, et al. Magnetic resonance angiography, function and viability evaluation in patients with Kawasaki disease. *J Cardiovasc Magn Reson* 2006; 8: 493–498.
- 21) Kondo C. Myocardial perfusion imaging in pediatric cardiology. *Ann Nucl Med* 2004; 18: 551–561.
- 22) http://jsnm.sakura.ne.jp/wp_jsnm/wp-content/themes/theme_jsnm/doc/PediatricNuclMed-Guideline1-2-3.pdf (Japanese)
- 23) Prabhu AS, Singh TP, Morrow WR, et al. Safety and efficacy of intravenous adenosine for pharmacologic stress testing in children with aortic valve disease or Kawasaki disease. *Am J Cardiol* 1999; 83: 284–286.

- 24) Hoshina M, Shiraishi H, Igarashi H, et al. Efficacy of iodine-123-15-(p-iodophenyl)-3-R, S-methylpentadecanoic acid single photon emission computed tomography imaging in detecting myocardial ischemia in children with Kawasaki disease. *Circ J* 2003; 67: 663–666.
- 25) Zhao C, Shuke N, Yamamoto W, et al. Impaired cardiac sympathetic nerve function in patients with Kawasaki disease: comparison with myocardial perfusion. *Pediatr Res* 2005; 57: 744–748.
- 26) Suda K, et al. Statin reduces persistent coronary arterial inflammation evaluated by serial 18fluorodeoxyglucose positron emission tomography imaging long after Kawasaki disease. *Int J Cardiol* 2015; 179: 61–62.
- 27) Sudo D, Nakamura Y. Nationwide surveys show that the incidence of recurrent Kawasaki disease in Japan has hardly changed over the last 30 years. *Acta Paediatr.* 2017;106:796–800.