

5. References

1. Hajiro T, Nishimura K, Tsukino M, Ikeda A, Koyama H, Izumi T. Comparison of discriminative properties among disease-specific questionnaires for measuring health-related quality of life in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. Mar 1998;157(3 Pt 1):785-90.
2. Nishimura K, Izumi T, Tsukino M, Oga T. Dyspnea is a better predictor of 5-year survival than airway obstruction in patients with COPD. *Chest*. May 2002;121(5):1434-40.
3. Criner GJ, Bourbeau J, Diekemper RL, et al. Executive summary: prevention of acute exacerbation of COPD: American College of Chest Physicians and Canadian Thoracic Society Guideline. *Chest*. Apr 2015;147(4):883-893.
4. Casaburi R, Patessio A, Ioli F, Zanaboni S, Donner CF, Wasserman K. Reductions in exercise lactic acidosis and ventilation as a result of exercise training in patients with obstructive lung disease. *Am Rev Respir Dis*. Jan 1991;143(1):9-18.
5. Gosselink R, Troosters T, Decramer M. Exercise training in COPD patients: the basic questions. *Eur Respir J*. Dec 1997;10(12):2884-91.
6. Petrof BJ, Calderini E, Gottfried SB. Effect of CPAP on respiratory effort and dyspnea during exercise in severe COPD. *J Appl Physiol (1985)*. Jul 1990;69(1):179-88.
7. Koch R, Augusto TRL, Ramos AG, Müller PT. Inspiratory Muscle Training Potentiates the Beneficial Effects of Proportional Assisted Ventilation on Exertional Dyspnea and Exercise Tolerance in COPD: A Proof-of-Concept Randomized and Controlled Trial. *COPD*. Aug 2020;17(4):384-391.
8. O'Donnell DE, Sanii R, Giesbrecht G, Younes M. Effect of continuous positive airway pressure on respiratory sensation in patients with chronic obstructive pulmonary disease during submaximal exercise. *Am Rev Respir Dis*. Nov 1988;138(5):1185-91.
9. O'Donnell DE, Sanii R, Younes M. Improvement in exercise endurance in patients with chronic airflow limitation using continuous positive airway pressure. *Am Rev Respir Dis*. Dec 1988;138(6):1510-4.
10. Kataoka J, Kuriyama A, Norisue Y, Fujitani S. Proportional modes versus pressure support ventilation: a systematic review and meta-analysis. *Ann Intensive Care*. Dec 10 2018;8(1):123.
11. Dolmage TE, Goldstein RS. Proportional assist ventilation and exercise tolerance in subjects with COPD. *Chest*. Apr 1997;111(4):948-54.

12. Bianchi L, Foglio K, Pagani M, Vitacca M, Rossi A, Ambrosino N. Effects of proportional assist ventilation on exercise tolerance in COPD patients with chronic hypercapnia. *Eur Respir J*. Feb 1998;11(2):422-7.
13. Bianchi L, Foglio K, Porta R, Baiardi R, Vitacca M, Ambrosino N. Lack of additional effect of adjunct of assisted ventilation to pulmonary rehabilitation in mild COPD patients. *Respir Med*. May 2002;96(5):359-67.
14. Hawkins P, Johnson LC, Nikolettou D, et al. Proportional assist ventilation as an aid to exercise training in severe chronic obstructive pulmonary disease. *Thorax*. Oct 2002;57(10):853-9.
15. Poggi R, Appendini L, Polese G, Colombo R, Donner CF, Rossi A. Noninvasive proportional assist ventilation and pressure support ventilation during arm elevation in patients with chronic respiratory failure. A preliminary, physiologic study. *Respir Med*. Jun 2006;100(6):972-9.
16. Borghi-Silva A, Oliveira CC, Carrascosa C, et al. Respiratory muscle unloading improves leg muscle oxygenation during exercise in patients with COPD. *Thorax*. Oct 2008;63(10):910-5.
17. Carrascosa CR, Oliveira CC, Borghi-Silva A, et al. Haemodynamic effects of proportional assist ventilation during high-intensity exercise in patients with chronic obstructive pulmonary disease. *Respirology*. Nov 2010;15(8):1185-91.
18. Koch R, Rapello GVG, de Tarso Müller P. Is Inspiratory Muscle Weakness a Determinant of Endurance Exercise Tolerance During NIV-Supported Exercise in Patients With COPD? *J Cardiopulm Rehabil Prev*. Nov 2018;38(6):E9-E11.
19. Higgins J, Thomas J. Cochrane Handbook for Systematic Reviews of Interventions. v6.4 2023
20. Ries AL, Bauldoff GS, Carlin BW, Casaburi R, Emery CF, Mahler DA, et al. Pulmonary Rehabilitation: Joint ACCP/AACVPR Evidence-Based Clinical Practice Guidelines. *Chest*. 2007;131(5 Suppl):4S-42S.
21. O'Donnell DE, D'Arsigny C, Fitzpatrick M, Webb KA. Exercise hypercapnia in advanced chronic obstructive pulmonary disease: the role of lung hyperinflation. *Am J Respir Crit Care Med*. 2002;166(5):663-8.
22. Killian KJ, Leblanc P, Martin DH, Summers E, Jones NL, Campbell EJ. Exercise capacity and ventilatory, circulatory, and symptom limitation in patients with chronic airflow limitation. *Am Rev Respir Dis*. 1992;146(4):935-40.
23. Belman MJ. Factors limiting exercise performance in lung disease. Ventilatory insufficiency. *Chest*. 1992;101(5 Suppl):253S-4S.
24. McCurdy BR. Hospital-at-home programs for patients with acute exacerbations of

chronic obstructive pulmonary disease (COPD): an evidence-based analysis. Ont Health Technol Assess Ser. 2012;12(10):1-65.

25. Akoumianaki E, Dousse N, Lyazidi A, Lefebvre JC, Graf S, Cordoli RL, et al. Can proportional ventilation modes facilitate exercise in critically ill patients? A physiological cross-over study : Pressure support versus proportional ventilation during lower limb exercise in ventilated critically ill patients. Ann Intensive Care. 2017;7(1):64.
26. Younes M. Proportional assist ventilation, a new approach to ventilatory support. Theory. Am Rev Respir Dis. 1992;145(1):114-20.
27. Zhang J, Luo Q, Chen R. Patient-Ventilator Interaction With Noninvasive Proportional Assist Ventilation in Subjects With COPD. Respir Care. 2020;65(1):45-52.
28. Kataoka J. Proportional assist ventilation. INTENSIVIST. 2018;10(3):677-689
29. Ayukawa K. Proportional assist ventilation /proportional pressure support. 人工呼吸. 2004;21(2):125-132
30. Japan Intensive Care Education and Research Association. "Brief Survey No. 20: Ventilator Settings and Weaning (Conducted in October 2012)." 2012.