

引用文献

- Ahmed, J., Khan, S., Gatt, M., et al. (2010). Compliance with enhanced recovery programmes in elective colorectal surgery. *The British Journal of Surgery*, 97(5), 754-758.
- Allvin, R., Berg, K., Idvall, E., et al. (2007). Postoperative recovery: A concept analysis. *Journal of Advanced Nursing*, 57(5), 552-558.
- Allvin, R., Ehnfors, M., Rawal, N., et al. (2008). Experiences of the postoperative recovery process: An interview study. *The Open Nursing Journal*, 2, 1-7.
- Allvin, R., Ehnfors, M., Rawal, N., et al. (2009). Development of a questionnaire to measure patient-reported postoperative recovery: Content validity and intra-patient reliability. *Journal of Evaluation in Clinical Practice*, 15(3), 411-419.
- Allvin, R., Svensson, E., Rawal, N., et al. (2011). The postoperative recovery profile (PRP) - a multidimensional questionnaire for evaluation of recovery profiles. *Journal of Evaluation in Clinical Practice*, 17(2), 236-243.
- Anderson, A. D., McNaught, C. E., MacFie, J., et al. (2003). Randomized clinical trial of multimodal optimization and standard perioperative surgical care. *The British Journal of Surgery*, 90(12), 1497-1504.
- Baker, C. A. (1989). Recovery: A phenomenon extending beyond discharge. *Scholarly Inquiry for Nursing Practice*, 3(3), 181-94.
- Bardram, L., Funch-Jensen, P., Jensen, P., et al. (1995). Recovery after laparoscopic colonic surgery with epidural analgesia, and early oral nutrition and mobilisation. *Lancet*, 345(8952), 763-764.
- Brieger, G. H. (1983). Early ambulation. A study in the history of surgery. *Annals of Surgery*, 197(4), 443-449.
- Canon, W. B. (1932). からだの知恵—この不思議なはたらき. 館鄰他訳 (1981). 講談社.
- 茶園美香 (2007). 第5章治療・処置を受けている患者の看護 4.術後患者の看護. 岩井郁子編, 系統看護学講座別巻17 臨床看護総論 (pp280-285). 医学書院.
- Crouter, S. E., Schneider, P. L., & Bassett, D. R., Jr. (2005). Spring-levered versus piezo-electric pedometer accuracy in overweight and obese adults. *Medicine and Science in Sports and Exercise*, 37(10), 1673-1679.

- Delaney, C. P., Zutshi, M., Senagore, A. J., et al. (2003). Prospective, randomized, controlled trial between a pathway of controlled rehabilitation with early ambulation and diet and traditional postoperative care after laparotomy and intestinal resection. *Diseases of the Colon and Rectum*, 46(7), 851-859.
- Eskicioglu, C., Forbes, S. S., Aarts, M. A., et al. (2009). Enhanced recovery after surgery (ERAS) programs for patients having colorectal surgery: A meta-analysis of randomized trials. *Journal of Gastrointestinal Surgery*, 13(12), 2321-2329.
- Fearon, K. C. (2010). ERAS: an update on evidence and protocols. *ESPEN Congress 2010 Highlight Topics*. <http://www.jspen.jp/doc6/sec7.html> [2012-01-03]
- Fearon, K. C., Ljungqvist, O., Von Meyenfeldt, M., et al. (2005). Enhanced recovery after surgery: A consensus review of clinical care for patients undergoing colonic resection. *Clinical Nutrition*, 24(3), 466-477.
- Gatt, M., Anderson, A. D., Reddy, B. S., et al. (2005). Randomized clinical trial of multimodal optimization of surgical care in patients undergoing major colonic resection. *The British Journal of Surgery*, 92(11), 1354-1362.
- Gouvas, N., Tan, E., Windsor, A., et al. (2009). Fast-track vs standard care in colorectal surgery: A meta-analysis update. *International Journal of Colorectal Disease*, 24(10), 1119-1131.
- Gustafsson, U. O., Hausel, J., Thorell, A., et al. (2011). Adherence to the enhanced recovery after surgery protocol and outcomes after colorectal cancer surgery. *Archives of Surgery*, 146(5), 571-577.
- Harper, C. M., & Lyles, Y. M. (1988). Physiology and complications of bed rest. *Journal of the American Geriatrics Society*, 36(11), 1047-1054.
- Henderson, V. (1969). 看護の基本となるもの. 湯槻ます, 小玉香津子訳 (1995). 日本看護協会出版会.
- Kaplan, R. M. (2002). Quality of life: An outcomes perspective. *Archives of Physical Medicine and Rehabilitation*, 83(12 Suppl 2), S44-50.
- 五十嵐英子, 片桐優子, 若山ユキ子他. (2003). 全身麻酔下における開腹手術患者の離床プログラムの開発と有効性. *日本看護学会論文集: 看護総合*, (34), 108-110.
- Inoue, Y., Kimura, T., Fujita, S., et al. (2003). A new parameter for assessing

- postoperative recovery of physical activity using an accelerometer. *Surgery Today*, 33(9), 645-650.
- Kamolz, T., & Pointner, R. (2002). Expectations of patients with gastroesophageal reflux disease for the outcome of laparoscopic antireflux surgery. *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques*, 12(6), 389-392.
- Kaplan, R. M. (2002). Quality of life: An outcomes perspective. *Archives of Physical Medicine and Rehabilitation*, 83(12 Suppl 2), S44-50.
- Kehlet, H. (1997). Multimodal approach to control postoperative pathophysiology and rehabilitation. *British Journal of Anaesthesia*, 78(5), 606-617.
- Kehlet, H., & Wilmore, D. W. (2002). Multimodal strategies to improve surgical outcome. *American Journal of Surgery*, 183(6), 630-641.
- Khan, S., Wilson, T., Ahmed, J., et al . (2010). Quality of life and patient satisfaction with enhanced recovery protocols. *Colorectal Disease*, 12(12), 1175-1182.
- Khoo, C. K., Vickery, C. J., Forsyth, N., et al. (2007). A prospective randomized controlled trial of multimodal perioperative management protocol in patients undergoing elective colorectal resection for cancer. *Annals of Surgery*, 245(6), 867-872.
- Korolija, D., Wood-Dauphinee, S., & Pointner, R. (2007). Patient-reported outcomes. how important are they? *Surgical Endoscopy*, 21(4), 503-507.
- Kumahara, H., Ishii, K., & Tanaka, H. (2006). Physical activity monitoring for health management: Practical techniques and methodological issue. *International Journal of Sport and Health Science*, 4, 380-393.
- Lassen, K., Soop, M., Nygren, J., et al. (2009). Consensus review of optimal perioperative care in colorectal surgery: Enhanced recovery after surgery (ERAS) group recommendations. *Archives of Surgery*, 144(10), 961-969.
- Leithauser, D. J., & Bergo, H. L. (1941). Early rising and ambulatory activity after operation. *Archives of Surgery*, 42(6), 1086-1093.
- Lin, J. H., Whelan, R. L., Sakellarios, N. E., et al. (2009). Prospective study of ambulation after open and laparoscopic colorectal resection. *Surgical Innovation*, 16(1), 16-20.

Melanson, E. L., Knoll, J. R., Bell, M. L., et al. (2004). Commercially available pedometers: Considerations for accurate step counting. *Preventive Medicine*, 39(2), 361-368.

宮田剛 (2009). Fast-track Surgery. 外科と代謝・栄養, 43(5), 109-112.

宮澤寛子, 高橋哲也, 熊丸めぐみ他. (2008). 心臓外科手術後の離床に対する応用行動分析学的アプローチ 段階パス導入についての紹介. 心臓リハビリテーション, 13(1), 100-104.

Moore, F. D. (1959). Metabolic care of the surgical patient. Philadelphia: W. B. Saunders.

Muller, S., Zalunardo, M. P., Hubner, M., et al. (2009). A fast-track program reduces complications and length of hospital stay after open colonic surgery. *Gastroenterology*, 136(3), 842-847.

日本看護科学学会看護学学術用語検討委員会編 (2005). 看護行為用語分類 看護行為の言語化と用語体系の構築. 日本看護協会出版会. 185-186.

大平雄一, 西田宗幹, 植松光俊. (2005). アクティブラーディングによる基本動作の種類を判別するための評価 各種基本動作における加速度領域分布特性. 専門リハビリテーション, 4, 28-31.

大平雄一, 西田宗幹, 大西和弘他. (2008). 自宅退院する入院患者における退院前後での身体活動量の比較検討. 理学療法科学, 23(2), 313-317.

雄西智恵美, 小島操子, 数間恵子. (1983). 術後老人患者の早期離床促進のための看護プログラム作成に関する研究(2) 初回離床時の呼吸・循環・筋力の経時的变化の検討. 日本看護科学会誌, 3(1), 31-38.

Polle, S. W., Wind, J., Fuhring, J. W., et al. (2007). Implementation of a fast-track perioperative care program: What are the difficulties? *Digestive Surgery*, 24(6), 441-449.

笹川寿美, 福録恵子, 藤田淳子他. (2005). 健常者における姿勢変換に伴う血圧・心拍応答と主観的指標との関連 起立性低血圧の発現に注目して. 京都府立医科大学看護学科紀要, 14, 29-35.

佐藤知子, 茂司恵美子 (2004). 看護の成果を追い求めて 新日鐵広畠病院におけるアウトカム・マネジメント導入の試み 術後の早期離床に向けての取り組み. 看護, 56(13),

96-98.

Schneider, P. L., Crouter, S. E., Lukajic, O., et al. (2003). Accuracy and reliability of 10 pedometers for measuring steps over a 400-m walk. Medicine and Science in Sports and Exercise, 35(10), 1779-1784.

Schneider, P. L., Crouter, S. E., & Bassett, D. R. (2004). Pedometer measures of free-living physical activity: Comparison of 13 models. Medicine and Science in Sports and Exercise, 36(2), 331-335.

瀬下明良. (1986). 早期離床と術後肺機能の関連についての臨床的検討. 東京女子医科大学雑誌, 56(12), 1093-1103.

篠澤洋太郎 (2007). 外科的侵襲の病態生理. 松野正紀監修, 標準外科学 (pp.11-16). 医学書院.

Spanjersberg, W. R., Reurings, J., Keus, F., et al. (2011). Fast track surgery versus conventional recovery strategies for colorectal surgery. Cochrane Database of Systematic Reviews (Online). (2)(2), doi: 10.1002/14651858.CD007635.pub2.

[2012-01-03]

谷口英喜 (2008). ERAS(enhaned recovery after surgery). 栄養－評価と治療, 25(6), 528-532.

Thompson, J. B., Maclean, K. F. & Coller, F. A. (1949). Role of the transverse abdominal incision and early ambulation in the reduction of postoperative complications. Archives of Surgery, 59(6), 1267-1277.

角田明良, 中尾健太郎, 平塚研之他 (2005). 科学的根拠に基づく結腸癌クリニカルパスの成績. 日本大腸肛門病会誌, 58, 107-111.

Varadhan, K. K., Neal, K. R., Dejong, C. H., et al. (2010). The enhanced recovery after surgery (ERAS) pathway for patients undergoing major elective open colorectal surgery: A meta-analysis of randomized controlled trials. Clinical Nutrition, 29(4), 434-440.

Vlug, M. S., Bartels, S. A., Wind, J., et al. (2011). Which fast track elements predict early recovery after colon cancer surgery? Colorectal Disease. [Epub ahead of print]. doi: 10.1111/j.1463-1318.2011.02854.x. [2012-01-03]

若本恵子 (2007). 第9章 術後リハビリテーション. 渡邊五朗, 宗村美江子編, 新体系看護

学全書別巻1 臨床外科看護学 I (pp279-287). メディカルフレンド社.

Walter, C. J., Collin, J., Dumville, J. C., et al. (2009). Enhanced recovery in colorectal resections: A systematic review and meta-analysis. *Colorectal Disease*, 11(4), 344-353.

渡邊ふみ子, 丸山和子 (2002). IV手術と看護の方法. 井上幸子, 平山朝子, 金子道子編, 看護学大系第9巻看護の方法[4] (pp104-183). 日本看護協会出版会.

Wilmore, D. W., & Kehlet, H. (2001). Management of patients in fast track surgery. *BMJ (Clinical Research Ed.)*, 322(7284), 473-476.

Wind, J., Polle, S. W., Fung Kon Jin, P. H., et al. (2006). Systematic review of enhanced recovery programmes in colonic surgery. *The British Journal of Surgery*, 93(7), 800-809.

Zorowitz, R. D., & Tietjen, G. E. (1999). Medical complications after stroke. *Journal of Stroke and Cerebrovascular Diseases*, 8(3), 192-196.