

参考／引用文献

- 阿彦忠之(2008). わが国の結核対策の現状と課題(4)「結核対策における都道府県、保健所の役割と課題」. 日本公衆衛生雑誌, 55(12), 848-852.
- Arcencio,R. A., Oliveira, M. F., Cardozo-Gonzales, R. I, et al(2008). City tuberculosis control coordinators' perspectives of patient adherence to DOT in Sao Paulo State, Brazil, 2005. International Journal of Tuberculosis and Lung Disease, 12(5), 527-531.
- Aye, R., Wyss, K., Abdualimova, H. & et al. (2010). Illness costs to households are a key barrier to access diagnostic and treatment services for tuberculosis in Tajikistan. BioMedCentral Research Notes, 3. 340-346. Retrieved September 20, 2013, <http://www.biomedcentral.com/>
- Awaisu, A., Mohamed, M. HN., Noordin, N. M. & et al. (2012). Impact of connecting tuberculosis directly observed therapy short-course with smoking cessation on health-related quality of life. Tabacco Induced Diseases, 10(2). Retrieved

September 20, 2013, from <http://www.tabaccoinduceddiseases.com/content/10/1>

- Bagchi, S., Ambe, G. & Sathiakumar, N. (2010). Determinants of Poor Adherence to Anti-Tuberculosis Treatment in Mumbai, India. International Journal of Preventive Medicine, 1(4), 223-232.
- Bam, T. S., Gunneberg, C., Chamroonsawasdi, K. & et al. (2006). Factors affecting patient adherent to DOTS in urban Kathmandu, Nepal. International Journal of Tuberculosis and Lung Diseases, 10(3). 270-276.
- Bissell, P., May, C. R., & Noyce, P. R. (2004). From compliance to concordance: barriers to accomplishing a re-framed model of health care interactions. Social Science & Medicine, 58, 851-862.
- Bissonnette, J. M. (2008). Adherence: a concept analysis. Journal of Advanced Nursing, 63(6), 634-43.
- Calhoun, LG., Tedeschi,RG. (2010). 心的外傷後成長ハンドブック～耐え難い体験が人の心にもたらすもの～. 宅香菜子, 清水研訳 (2013) . 医学書院.
- Cann, A., Calhoun, L. G., Tedeschi, R. G. & et al. (2010). A short form of the posttraumatic Growth Inventory. Anxiety, Stress & Coping, 23(2), 127-137.
- Chamla D. (2004). The assessment of patients' health related quality of life during tuberculosis treatment in Wuhan, China. International Journal of Tuberculosis and Lung Diseases, 8(9). 1100-1106.
- Chang, B., Wu, A. B., Hansel, N. N., & et al. (2004). Quality of life in tuberculosis: A review of the English language literature. Quality of Life Research, 13, 1633-1642.
- Chaulk, C. P. & Kazandjian, V. A. (1998). Directly observed therapy for treatment completion of pulmonary tuberculosis: Consensus Statement of the Public Health Tuberculosis Guidelines Panel. JAMA, 279, 943-948.
- Cox H. S., Morrow, M., & Deutschmann, P. W. (2008). Long term efficacy of DOTS regimens for tuberculosis: systematic review. BMJ. 1(336).
- Centers for Disease Control and Prevention(1990). Screening for Tuberculosis and Tuberculosis Infection in High-Risk Populations Recommendations of the Advisory Committee for Elimination of Tuberculosis. 39(RR-8);1-7.
- Corless,I. B., Nicholas, P. K., Wantland, D., et al(2006). The impact of meaning in life

- and life goals on adherence to a tuberculosis medication regimen in South Africa. The International Journal of Tuberculosis and Lung Disease, 10(10), 1159-1165.
- Creswell, J. W. (2003). Research Design. Qualitative, Quantitative and Mixed Method Approaches 2nd editions. London, SAGE Publication Inc.
- Creswell, J. W., Clark, VL. (2011). 大谷順子 (2012). 人間科学のための混合研究法. 北大路書房.
- Davidson, H., Schluger, N. W., Feldman, P. H., & et al(2000). The effects of increasing incentives on adherence to tuberculosis directly observed therapy. International Journal of Tuberculosis and Lung disease, 4(9), 860-865.
- Demissie,M., Getahun, H., & Lindtjorn B. (2003). Community tuberculosis care through "TB clubs" in rural North Ethiopia. Social Science & Medicine, 56(19), 2009-2018.
- Dimitrova, B., Balabanova, D., Atun, R. & et al. (2006). Health service providers' perceptions of barriers to tuberculosis care in Russia. Oxford University Press, 265-274. Retrieved September 20, 2013, from <http://heapol.oxfordjournals.org/>
- Duria, M., Nandini, S., Nalender, P. S. & et al. (2009). A study of the Impact of Tuberculosis on the Quality of Life and the Effect After Treatment with DOTS. Asia-Pacific Journal of Public Health, 21(3). 312-320.
- Frieden TR., et al. (2003). Tuberculosis. Lancet. 362. 887-889.
- Guo N., Marra F., Marra C.A. (2009). Measuring health-related quality of life in tuberculosis: a systematic review. Health and Quality of Life Outcomes. 7(14).
- 橋本容子, 野村繁雄, 和田圭司(2009). 地域 DOTS の推進—服薬支援計画票を活用して—。結核, 84 (1), 165-172.
- Hasker, E., Khodjikhanov, M., Sayfiddinova, S. & et al. (2010). Why do tuberculosis patients default in Tashkent City, Uzbekistan? A qualitative study. International Journal of Tuberculosis and Lung Diseases, 14(9). 1132-1139.
- Horne, R., Weinman, J., Barber, N., & et al. (2005). Concordance, adherence and compliance in medicine taking.
http://www.nets.nihr.ac.uk/_data/assets/pdf_file/0009/64494/FR-08-1412-076.pdf. [2013-10-24].
- 星野斎之, 大森正子, 内村和宏, 他(2008). 高齢者結核の疫学的検討. 結核, 83(5), 423-

- 堀田信之, 宮沢直幹, 吉山崇, 他(2013). 結核患者の生命予後. 結核, 88(6), 565-570.
- 石川信克(2005). 世界の結核問題にいかに関わるべきか—国際協力の課題と実践—. 結核, 80(2), 89-94.
- 石川信克(2008). わが国の結核対策の現状と課題(3)「世界, 日本の結核の疫学と課題」. 日本公衆衛生雑誌, 55(11), 791-794.
- 伊藤邦彦, 吉山崇, 永田容子, 他(2008). 結核治療中断を防ぐために何が必要か?. 結核, 83(9), 621-628.
- 稻垣孝行, 中川拓, 前田伸司, 他(2013). 初回肺結核治療中において短期間に多剤耐性化した結核菌の遺伝子解析—臨床分離株を用いて. 結核, 88(7), 595-604.
- Jarvis, M. (2010). Tuberculosis 2: exploring methods of diagnosis, treatment regimens and concordance. Nursing Times, 106(2), 22-24.
- 神楽岡澄, 大森正子, 高尾良子, 他(2008). 新宿区保健所における結核対策—DOTS 事業の推進と成果—. 結核, 83(9), 611-620.
- Kaler ME, Erbes CR, Tedeschi RG, et al. (2011). Factor structure and concurrent validity of the Posttraumatic Growth Inventory-Short Form among veterans from the Iraq War. J Trauma Stress, 24(2), 200-207.
- 加藤由希子, 有本梓, 島村珠枝, 村嶋幸代(2012). 自覚症状のある肺結核患者の受診の遅れとその特徴. 日本公衆衛生雑誌, 59(4), 251-258.
- Kurbatova, E V., Taylor, A., Gammino, V. M., et al (2012). Predictors of poor outcomes among patients treated for multidrug-resistant tuberculosis at DOTS-plus projects. Tuberculosis, 92, 397-403.
- 結核予防会(2014). 結核の統計 2013. 東京, 財団法人結核予防会.
- Khan, M., Walley, J. D., Witter, S. N. & et al. (2002). Costs and cost-effectiveness of different DOT strategies for the treatment of tuberculosis in Pakistan. Health Policy and Planning, 17(2), 178-186.
- Khan, M., Walley, J. D., Witter, S. N. & et al. (2005). Tuberculosis patient adherence to direct observation: results of a social study in Pakistan. Oxford University Press, 354-365. Retrieved September 20, 2013, from <http://heapol.oxfordjournals.org/>
- 木村もりよ(2004). わが国における DOTS の費用対効果分析—大阪市住所不定者を 1 モデ

- ル集団として—. 厚生の指標, 51, 17-21.
- 木下節子, 大森正子, 塚本和秀, 他(2007). 駅周辺の不特定多数利用施設を中心とした結核感染—都市結核問題の観点より. 結核, 82(11), 749-757.
- 小林弘美, 小柳孝太郎, 加藤収, 他(2013). 血液透析施設における超多剤耐性結核の集団感染. 結核, 88(5), 477-484.
- 厚生労働省. http://www.mhlw.go.jp/seisakunitsuite/bunya/kenkou_iryou/kenkou/ [2013-4-25]
- Long Q. (2011). Patient medical costs for tuberculosis treatment and impact on adherence in China: a systematic review. BMC Public Health. 11.393-402.
- Lorraine, S E. (1999). Compliance: A Concept Analysis. Nursing Forum. 34(1). 5-11.
- MacIntyre, C. R., Goebel, K., Brown, G. V. & et al. (2003). A randomized controlled clinical trial of the efficacy of family-based direct observation of anti-tuberculosis treatment in an urban, developed-country setting. International Journal of Tuberculosis and Lung disease, 7(9). 848-854.
- Macq J., Torfoss T., Getahun H. (2007). Patient empowerment in tuberculosis control: reflection on past documented experiences. Tropical Medicine and International Health. 12(7). 873-885.
- Maher D., et al. (1997). Treatment of tuberculosis: guidelines for national programmes, second edn. Geneva: WHO.
- Martins, N., Grace, J. & Kelly, P. M. (2008). An ethnographic study of barriers to and enabling factors for tuberculosis treatment adherence in Timor Leste. International Journal of Tuberculosis and Lung Diseases, 12(5). 532-537.
- Maslow, A. H. (1970). Motivation and personality. New York: Harper & Row Publication.
(小口忠彦, 他訳(2004). 人間性の心理学. 東京:産業能率大学出版部.)
- 松本健二, 福永淑江, 門林順子, 他(2009). 受診の遅れに関する検討. 結核, 84(7), 523-529.
- 森亨(2004). 新たな結核対策の技術と展望. 結核, 79(10), 587-604.
- McDonnell M., Turner J., Weaver T. M. (2001). Antecedents of Adherence to Antituberculosis Therapy. Public Health Nursing. 18(6). 392-400.
- Morisky DE, Green LW, Levine DM. (1986). Concurrent and predictive validity of a self-

- reported measure of medication adherence. Medicine and Care. 24(1), 67-74.
- Munro S. A., Lewin S. A., et al. (2007). Patient Adherence to Tuberculosis Treatment: A systematic Review of Qualitative Research. PLOS Medicine. 4(7), 1230-1245.
- M'Imunya, J. M., Kredo, T., & Volmink, J.(2012). Patient education and counseling for promoting adherence to treatment for tuberculosis (Review). Cochrane Database of Systematic Reviews.
- 長弘佳恵, 小林小百合, 村嶋幸代(2007). 不安定就労・生活者にとっての Directly Observed Treatment Short-course (DOTS)受療の意味. 日本公衆衛生雑誌, 54(12), 857-866.
- Naidoo P. (2013). Predictors of tuberculosis (TB) and antiretroviral (ARV) medication non-adherence in public primary care patients in South Africa: a cross sectional study. BMC Public Health. 13, 396-406.
- 中西好子, 大山泰雄, 高橋光良, 他(1997). サウナでの結核多発の分子疫学的解明 大都市のホームレスの結核問題に関連して. 日本公衆衛生雑誌. 44(10), 769-768.
- 成田友代, 小林典子 (2009). 地域 DOTS の展開. 結核, 84(4), 187-201.
- Niazi, A. D. & Al-Delaimi, A. M. (2003). Impact of community participation on treatment outcomes and compliance of DOTS patients in Iraq. Eastern Mediterranean Health Journal, 9(4), 709-717.
- 日本語版 EuroQol 開発委員会 (1998). 日本語版 EuroQol の開発. 医療と社会, 8(1), 109-123.
- 沼田久美子, 藤田利治(2002). 新宿区登録患者における治療中断の関連要因と Directly Observed Therapy の意義. 日本公衆衛生雑誌. 49(1), 58-63.
- Ohkado A., et al. (2009). Transmission of specific genotype streptomycin resistant strains of *Myco-bacterium tuberculosis* in the Tokyo Metropolitan Area in Japan. BMC Infectious Disease. 9, 138.
- 大森正子, 下内昭, 伊藤邦彦, 他 (2012). 結核サーベイランス情報からみた薬剤耐性結核患者の背景. 結核, 87(4), 357-365.
- Ohshima N., et.al. (2013). Chronological Decrease of Tuberculosis Incidence Rates by Organ Classification Based on Birth Cohort Study in Japan, 1975-2005. Kekkaku. 88(1), 1-7.
- 奥村昌夫, 吉山崇, 尾形英雄, 他(2011). 当院における多剤耐性結核症(MDR-TB), 超多剤

耐性結核症(XDR-TB)の耐性化の過程についての検討. 結核, 86(12), 863-868.

小塩真司 (2011). SPSS と Amos による心理・調査データ解析[第2版]—因子分析・今日分散構造分析まで. 東京図書株式会社.

O'boyle, S. J., Power, J. J. & Watson, J. P. (2002). Factors affecting patient compliance with anti-tuberculosis chemotherapy using the directly observed treatment, short-course strategy (DOTS). International Journal of Tuberculosis and Lung disease, 6(4). 307-312.

Prochaska J.O., Velicer W.F. The transtheoretical model of health behavior change. American Journal of Health Promotion 12(1), p38 - 48, 1997.

Prochaska J.O., Redding C.A., Evers K.E. The transtheoretical model and stages of change.

In K. Glanz, B.K. Rimer, K. Viswanath (eds.), Health behavior and health education : theory, research, and practice. (4th ed), Jossey - Bass, p97 - 121, 2008.

Raviglione MC., et al. (1995). Global epidemiology of tuberculosis: mobility and mortality of a worldwide epidemic. JAMA. 273.220-226.

Rusen, I. D., Ait-Khaled, N., Alarcon, E. & et al. (2007). Cochrane systematic review of directly observed therapy for treating tuberculosis: good analysis of the wrong outcome. International Journal of Tuberculosis and Lung Disease, 11(2), 120-121.

Sagbakken, M., Bjune, G. A. & Frich, J. C.(2011). Humiliation or care? A qualitative study of patients' and health professionals' experiences with tuberculosis treatment in Norway. Scandinavian Journal of Caring Sciences, 26. 313-323.

島尾忠男(2003). Living with TB for fifty years 結核と歩んで五十年. 東京, 財団法人結核予防会.

Shimamura, T, Taguchi, A, Kobayashi, S & et al. (2012). The starategy of Japanese public health nurses in medication support for high-risk tuberculosis patients. Public Health Nursing, 30(4), 370-378.

Snowden, A., Martin, C., Mathers, B. & et al. (2013). Concordance: a concept analysis. Journal of Advanced Nursing 00(00), 000-000.doi: 10.1111/jan.12147 [2013-10-24]

Stevenson, F. A., Cox, K., Britten, N., & et al. (2004). A systematic review of the research on communication between patients and health care professionals about

- medicines: the consequences for concordance. Blackwell Publishing Ltd, Health Expectations, 7, 235-245.
- Storla DG, et al. 2008., A systematic review of delay in the diagnosis and treatment of tuberculosis. BMC Public Health. 8, 15-24.
- Styblo K.(1989). Overview and Epidemiologic Assessment of the Current Global Tuberculosis Situation with an Emphasis on Control in Developing Countries. Clinical Infectious Diseases, 11(2), 339-346.
- 多田有希, 大森正子, 伊藤邦彦, 他 (2004). 川崎市の結核対策. 結核, 79(1), 17-24.
- Taku, K., Lawrence, G., Calhoun, R. G., & et al(2007). Examining posttraumatic growth among Japanese university students. Anxiety, Stress, & Coping, 20(4), 353-367.
- Tedschi, R. G., & Calhoun, L. G. (1996). The posttraumatic Growth Inventory : Measuring the positive legacy of trauma. Journal of Traumatic Stress, 9, 455-471.
- Tedschi, R. G., & Calhoun, L. G. (2004). Posttraumatic Growth : Conceptual foundations and empirical evidence. Psychological Inquiry, 15, 1-18.
- Terra, M F. & Bertolozzi, M. R. (2008). Does directly observed treatment ("DOTS") contribute to tuberculosis treatment compliance?. Rev Latino-am Enfermagem, 16(4). 659-664.
- 豊田誠(2012). 若年者を中心に複数の経路で拡大した結核集団感染. 結核, 87(12), 757-763.
- Trinity, L. I. (2005). Compliance: A Concept Analysis. Nursing Forum. 44(3). 189-194.
- Tuberculosis Research Committee (2007). Drug-resistant *Mycobacterium tuberculosis* in Japan: a nationwide survey, 2002. International Journal of Tuberculosis and Lung Diseases, 11(10), 1129-1135.
- 土屋有紀, 長谷川敏彦, 西村周三, 他 (1998). 日本語版 EuroQol 検証の課題. 医療と社会, 8(1), 67-77.
- Volmink, J. & Garner, P.(1997). Systematic review of randomized controlled trials of strategies to promote adherence to tuberculosis treatment. BMJ, 315, 1403-1406.
- Volmink, J. & Garner, P. (2012). Directly observed therapy for treating tuberculosis. Cochrane Database of Systematic Reviews.
- Walley, J. D., Khan, M. A., Newell, J. N. & et al. (2001). Effectiveness of the direct observation component of DOTS for tuberculosis: a randomized controlled trial in

- Pakistan. *The Lancet*, 357(9257). 664-669.
- Wandalo, E., Makundi, E., Hasler, T. & et al. (2006). Acceptability of community and health facility-based directly observed treatment of tuberculosis in Tanzanian urban setting. *Health Policy*, 78. 284-294. Retrieved September 20, 2013, <http://www.elsevier.com/locate/healthpol>
- Ward HA., et al. (2004). Extent of pulmonary tuberculosis in patients diagnosed by active compared to passive case finding. *International Journal of Tuberculosis and Lung Diseases*. 8(5). 593-597.
- World Health Organization(1994). Framework for effective tuberculosis control. WHO Global Tuberculosis Programme 1994. WHO/TB, 94, 179.
- World Health Organization(1999). What is DOTS? A Guide to Understanding the WHO-recommended TB Control Strategy Known as DOTS. WHO, Geneva,1999.
- World Health Organization. (2003). Adherence to Long-Term Therapies: Evidence for Action. Geneva, WHO.
- World Health Organization. (2011). The Global Plan to Stop TB 2011-2015.
- World Health Organization. (2012). Global Tuberculosis Report 2012.
- Xu, L., Gai, R., Liu, Z. & et al. (2010). Socio-economic factors affecting the success of tuberculosis treatment in six conties of Shandong Province, China. *International Journal of Tuberculosis and Lung Diseases*, 14(4). 440-446.
- 吉山崇(1998). 多剤耐性結核の疫学. 結核, 73(10), 665-672.
- 吉山崇, 尾形英雄, 和田雅子 (2005). 多剤耐性結核の治療成績. 結核, 80(10), 687-693.
- 山本弘庫, 真島千寿子, 菅野美穂(2003). DOTS 対象者の治療成績および背景の検討. 保健師・看護師の結核展望, 41(2), 53-57.
- Yin, X., Tu, X., Tong Y. et al. (2012). Development and Validation of a Tuberculosis Medication Adherence Scale. *PLOS ONE*, 7(12), 1-6.