

文 献

- 有澤正義, 福庭一人, 関隆, 持丸文雄 (1996). 生体インピーダンス法による妊産婦浮腫の評価. 日本産科婦人科学会雑誌, 48(1), 25-31.
- Armstrong, L.E. (2007). Assessing hydration status: The elusive gold standard. *Journal of the American College of Nutrition*, 26(5), 575S-584S.
- Bieniarz, J., Burd, L., Motew, M. & Scommegna, A. (1971). Inhibition of uterine contractility in labor. *American Journal of Obstetrics and Gynecology*, 111(7), 874-879.
- Blackburn, S.T. (2007). *Maternal, fetal, & neonatal physiology A clinical perspective 3rd ed.* St. Lois, Missouri: Saunders Elsevier.
- Centers for Disease Control and Prevention, National Center for Health Statistics (2009). *National Health and Nutrition Examination Survey, 2009-2010 Dietary Interview Component Includes 24-Hour Dietary Recall Interview and Post-Recall Questionnaires.* Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, http://www.cdc.gov/nchs/data/nhanes/nhanes_09_10/dietaryrecall_e.pdf [Accessed on: 6/30/10]
- Centers for Disease Control and Prevention, National Center for Health Statistics (2010). *National Health and Nutrition Examination Survey, 2009-2010 Measuring Guides for the Dietary Recall Interview.* Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. http://www.cdc.gov/nchs/nhanes/measuring_guides_dri/measuringguides.htm [Accessed on: 6/30/10]
- Chelmos, D., Baker, E. R. & Jones, L. (1996). Maternal intravenous hydration and amniotic fluid index in patients with preterm ruptured membranes. *Journal of the Society for Gynecologic Investigation*, 3(3), 127-130.
- Chumlea, W.C. & Sun, S.S. (2005). Bioelectrical impedance analysis. In S.B. Heymsfield, T.G. Lohman, Z. Wang & S.B. Going (Eds.), *Human Body Composition 2nd ed.* (pp.79-88). Champaign, IL: Human Kinetics.

- Cohen, J. (1992). A Power Primer. *Psychological Bulletin*, 112(1), 155-159.
- Doi, S., Osada, H., Seki, K. & Sekiya, S. (1998). Effect of maternal hydration on oligohydramnios: A comparison of three volume expansion methods. *Obstetrics and Gynecology*, 92(4 Pt 1), 525-529.
- Gleichauf, C. N. & Roe, D. A. (1989). The menstrual cycle's effect on the reliability of bioimpedance measurements for assessing body composition. *The American Journal of Clinical Nutrition*, 50(5), 903-907.
- Gonzales, G.F., Steenland, K. & Tapia, V. (2009). Maternal hemoglobin level and fetal outcome at low and high altitudes. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 297, R1477-R1485.
- Goodlin, R.C., Quaife, M.A. & Dirksen, J.W. (1981). The significance, diagnosis, and treatment of maternal hypovolemia as associated with fetal/maternal illness. *Seminars in Perinatology*, 5(2), 163-174.
- Grandjean, A. C. & Grandjean, N. R. (2007). Dehydration and cognitive performance. *Journal of the American College of Nutrition*, 26(5 Suppl), 549S-554S.
- Grosso, L. M., Triche, E. W., Belanger, K., Benowitz, N. L., Holford, T. R. & Bracken, M. B. (2006). Caffeine metabolites in umbilical cord blood, cytochrome P-450 1A2 activity, and intrauterine growth restriction. *American Journal of Epidemiology*, 163(11), 1035-1041.
- Grosso, L. M., Triche, E., Benowitz, N. L. & Bracken, M. B. (2008). Prenatal caffeine assessment: Fetal and maternal biomarkers or self-reported intake? *Annals of Epidemiology*, 18(3), 172-178.
- Guinn, D. A., Goepfert, A. R., Owen, J., Brumfield, C. & Hauth, J. C. (1997). Management options in women with preterm uterine contractions: A randomized clinical trial. *American Journal of Obstetrics and Gynecology*, 177(4), 814-818.
- Helfgott, A. W., Willis, D. C. & Blanco, J. D. (1994). Is hydration and sedation beneficial in the treatment of threatened preterm labor? A preliminary report. *Journal of Maternal-Fetal Medicine*, 3(1), 37-42.
- Hofmeyr, G. J. & Gulmezoglu, A. M. (2002). Maternal hydration for increasing amniotic fluid volume in oligohydramnios and normal amniotic fluid volume. *Cochrane*

- database of systematic reviews (Online), (Issue 1). CD000134.
- Institute of Medicine (2004). Dietary reference intakes for water, potassium, sodium, chloride, and sulfate. Washington DC: The National Academies Press.
- 石川雅一, 品川長夫(1991). Bioelectrical impedance analysis による栄養評価. 日本臨床 49, 87-90.
- 金山尚裕, 河村隆一, 照井克生 (2008). D. 産科疾患の診断・治療・管理 16. 帝王切開の麻酔. 日本産科婦人科学会雑誌, 60(5), N-104-N-110.
- Kavouras, S. A. (2002). Assessing hydration status. *Current Opinion in Clinical Nutrition and Metabolic Care*, 5(5), 519-524.
- 木部博文, 浅井光興, 斉藤徹, 梶浦弘明, 二石かおり, 野口昌良 (2002). [周産期医療のトピックス]妊婦水泳による体内水分分布の変化 生体インピーダンス法による水分動態の解析. 産婦人科の世界, 54(10), 1055-1059.
- 小宮秀一, 中尾武平 (2002). 健康行動の科学 身体組成学 ー栄養・運動・健康ー. 東京: 技報堂出版株式会社.
- 厚生労働統計協会(2011). 第2章人口動態, 国民衛生の動向 2011/2012 厚生の指標 増刊, 58(9). 45-69.
- 久保隆彦 (2010). クリニカルカンファレンス 2 産科異常出血の管理 1) 分娩時異常出血量の新しい考え方, 日本産科婦人科学会雑誌, 62(9), N-121-N-125.
- Larciprete, G., Valensise, H., Vasapollo, B., Altomare, F., Sorge, R., Casalino, B., De Lorenzo, A. & Arduini, D. (2003a). Body composition during normal pregnancy: reference ranges. *Acta Diabetologica*, 40, S225-S232.
- Larciprete, G., Valensise, H., Vasapollo, B., Pierro, G.D., Menghini, S., Magnani, F., De Lorenzo, A. & Arduini, D. (2003b). Maternal body composition at term gestation and birth weight: is there a link? *Acta Diabetologica*, 40, S222-S224.
- Lof, M. & Forsum, E. (2004). Evaluation of bioimpedance spectroscopy for measurements of body water distribution in healthy women before, during, and after pregnancy. *Journal of Applied Physiology*, 96, 967-973.
- Lukaski, H.C., Siders, W.A., Nilsen, E.J. & Hall, C.B. (1994). Total body water in pregnancy: assessment by using bioelectrical impedance. *American Journal of Clinical Nutrition*, 59, 578-585.

- Lukaski, H.C., Hall, C.B. & Siders, W.A. (2007). Assessment of change in hydration in women during pregnancy and postpartum with bioelectrical impedance vectors. *Nutrition*, 23, 543-550.
- Malhotra, B. & Deka, D. (2002). Effect of maternal oral hydration on amniotic fluid index in women with pregnancy-induced hypertension. *The Journal of Obstetrics and Gynaecology Research*, 28(4), 194-198.
- Meis, P.J., Michielutte, R., Peters, T.J., Wells, H.B., Sands, R.E., Coles, E.C. & Johns, K.A. (1995). Factors associated with preterm birth in Cardiff, Wales. I. Univariable and multivariable analysis. *American Journal of Obstetrics and Gynaecology*, 173(2), 590-596.
- Montgomery, K. S. (2002). An Update on Water Needs during Pregnancy and Beyond. *The Journal of Perinatal Education*, 11(3), 40-42.
- Morita, H., Takeuchi, K., Funakoshi, Mizutori, M. & Maruo. T. (1999). Potential use of bioelectrical impedance analysis in the assessment of edema in pregnancy. *Clinical and Experimental Obstetrics & Gynecology*, 26(3-4), 151-154.
- 中田かおり(2010). 妊婦の体水分バランスと予後に関する文献検討. *日本助産学会誌*, 24(2), 196-204.
- 中田かおり(2013). 生体インピーダンス法による体水分と妊娠・分娩異常との関連：パイロット・スタディ, *日本助産学会誌*, 27(1), 掲載予定.
- 日本高血圧学会. 特定健診・特定保険指導実施に対する日本高血圧学会よりの提言（平成20年4月）. from <http://www.jpnh.org>. Retrieved June 30, 2010.
- 西澤美幸, 佐藤等, 池田義雄 (2007). 自己健康管理ツールとしての BIA 体組成計. *臨床病理レビュー特集第 138 号別冊*, 158-164.
- 阪本要一, 船間敬子, 池田義雄, Heymsfield, S.B., 大槻尚子, 西澤美幸 (1995). インピーダンス法による体組成の検討. 第 16 回日本肥満学会記録, 312-313.
- 阪本要一, 佐藤等, 池田義雄 (2003). 体脂肪量の測定法 電気伝導度法, インピーダンス法. *日本臨床*, 61(増 6), 368-373.
- Sanin Aguirre, L.H., Reza-Lopez, S. & Levario-Carrillo, M. (2004). Relation between Maternal Body Composition and Birth Weight. *Biology of the Neonate*, 86(1), 55-62.
- Sarvazyan, A., Tatarinov, A. & Sarvazyan, N. (2005). Ultrasonic assessment of tissue

hydration status. *Ultrasonic*, 43, 661-671.

佐々木敏 (2009). 簡易型自記式食事歴法質問票(BDHQ)1-1 2009 01 版. from

<http://www.ebn-japan.org> Retrieved June 30, 2010.

Schoeller, D.A. (2005). Hydrometry. In S.B. Heymsfield, T.G. Lohman, Z. Wang & S.B. Going (Eds.), *Human Body Composition 2nd ed.* (pp.35-49). Champaign, IL: Human Kinetics.

Selberg, O. & Selberg, D. (2002). Norms and correlates of bioimpedance phase angle in healthy human subjects, hospitalized patients and patients with liver cirrhosis. *European Journal of Applied Physiology*, 86, 509-516.

Shanholtzer, B.A. & Patterson, S.M. (2003). Use of bioelectrical impedance in hydration status assessment: reliability of a new tool in psychophysiology research. *International Journal of Psychophysiology*, 49, 217-226.

Stan, C.M., Boulvain, M., Pfister, R. & Hirsbrunner-Almagbaly, P. (2002). Hydration for treatment of preterm labour. *Cochrane Database of Systematic Reviews (Online)*, (Issue 2), CD003096.

Steer, P., Alam, M.A., Wadsworth, J. & Welch, A. (1995). Relation between maternal haemoglobin concentration and birth weight in different ethnic groups, *BMJ*, 310, 489-491.

Sun, S.S. & Chumlea, W.C. (2005). Statistical Methods. In S.B. Heymsfield, T.G. Lohman, Z. Wang & S.B. Going (Eds.), *Human Body Composition 2nd ed.* (pp.151-160). Champaign, IL: Human Kinetics.

武内享介, 喜吉賢二, 安田立子, 中後聡, 船越徹, 本山寛, 丸尾猛 (2001). 生体インピーダンス法による妊娠中毒症の水分動態分析とその重症化の予知. *日本中毒症学会雑誌*, 9, 97-100.

植田充治, 谷口幸一, 山村博三 (1987). 妊娠母体循環血液量の測定方法ーインドシアニングリーンを使った希釈法ー. *産婦人科の進歩*, 39(2), 193-194.

上田康夫, 丸尾原義, 足高善彦 (2002). 生体インピーダンス法を用いた妊婦体組成の検討と体水分量・体脂肪量比の妊娠浮腫評価への応用. *日本産科婦人科学会雑誌*, 54(5), 799-808.

上田康夫, 丸尾原義, 足高善彦, 深山知子, 本田由佳, 中林正雄 (2004). 正常および妊娠中

毒症妊婦での母体体脂肪・体水分の動態分析－胎児部分重量補正を加えた生体インピーダンス法による検討－. 産婦人科の進歩, 56(3), 275-286.

Umber, A. & Chohan, M. A. (2007). Intravenous maternal hydration in third trimester oligohydramnios: Effect on amniotic fluid volume. *Journal of the College of Physicians and Surgeons*, 17(6), 336-339.

Valensise, H., Andreoli, A., Lello, S., Magnani, F., Romanini, C. & De Lorenzo, A. (2000). Multifrequency bioelectrical impedance analysis in women with a normal and hypertensive pregnancy. *American Journal of Clinical Nutrition*, 72, 780-783.

Valensise, H., Vasapollo, B., Novelli, G.P., Larciprete, G., Andreoli, A., Altomare, F., DiPierro, G., Galante, A., Arduini, D. & De Lorenzo, A. (2004). Total body water estimation and maternal cardiac systolic function assessment in normal and gestational hypertensive pregnant women. *Medical Science Monitor*, 10(9), CR530-534.

Valenzuela, G., Cline, S. & Hayashi, R. H. (1983). Follow-up of hydration and sedation in the pretherapy of premature labor. *American Journal of Obstetrics and Gynecology*, 147(4), 396-398.

山下祐子, 茅原幸子, 甲元ゆみ子(2000). 妊婦の体内脂肪計による測定結果の検討. 母性衛生, 41(2), 242-247.

Yan-Rosenberg, L., Burt, B., Bombard, A. T., Callado-Khoury, F., Sharett, L., Julliard, K. & Weiner, Z. (2007). A randomized clinical trial comparing the effect of maternal intravenous hydration and placebo on the amniotic fluid index in oligohydramnios. *The Journal of Maternal-Fetal & Neonatal Medicine*, 20(10), 715-718.

Yasuda, R., Takeuchi, K., Funakoshi, T. & Maruo, T. (2003). Bioelectrical impedance analysis in the clinical management of preeclampsia women with edema. *Journal of Perinatal Medicine*, 31, 275-280.

Yip, R. (2000). Significance of an abnormally low or high hemoglobin concentration during pregnancy: special consideration of iron nutrition. *American Journal of Clinical Nutrition*, 72(1 Suppl), 272S-279S.

Zhang, Q., Ananth, C.V., Li, Z. & Smulian, J.C. (2009). Maternal anaemia and preterm birth: a prospective cohort study. *International Journal of Epidemiology*, 38,

1380-1389.