

## 文献

1. Aissaoui, Y., Zeggwagh, A. A., Zekraoui, A., Abidi, K., Abouqal, R. (2005). Validation of behavioral pain scale in critically ill, sedated, and mechanically ventilated patients. *Anesthesia and Analgesia*, 101, 1470-1476.
2. Ambuel, B., Hamlett, K. W., Marx, C. M., & Blumer, J. L. (1992). Assessing distress in pediatric intensive care environments: The COMFORT scale. *Journal of Pediatric Psychology*, 17(1), 95-109.
3. Andrews, C. M., & Chrzanowski, M. (1990). Maternal position, labor, and comfort. *Applied Nursing Research : ANR*, 3(1), 7-13.
4. Ashkenazy, S., & Dekeyser-Ganz, F. (2011). Assessment of the reliability and validity of the comfort scale for adult intensive care patients. *Heart & Lung : The Journal of Critical Care*, 40(3), e44-51.
5. Bader MK, Burn M, Campbell M, Chulay M, Hill C, Delongh M, et al. (2004). Consensus conference on sedation assessment: a collaborative, American Association of Critical-Care Nurses, and Saint Thomas health system. *Crit Care Nurse*, 24, 33-41.
6. Chulay, M. (2004). Sedation assessment: Easier said than done! *Critical Care Nursing Clinics of North America*, 16(3), 359-64.
7. Carnevale, F. A., & Razacks, S. (2002). An item analysis of COMFORT scale in a pediatric intensive care unit. *Pediatric Critical Care Medicine*, 117, 192-196.
8. De Jonghe, B., Cook, D., Appere-De-Vecchi, C., Guyatt, G., Meade, M., & Outin, H. (2000). Using and understanding sedation scoring systems: A systematic review. *Intensive Care Medicine*, 26(3), 275-285.
9. Ely EW, Inouye SK, Bernard GR, et al. (2001a). Delirium in mechanically ventilated patients: validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU). *JAMA*, 286(21), 2703-10.
10. Ely EW, Margolin R, Francis J, et al. (2001b). Evaluation of delirium in critically ill patients: validation of the Confusion Assessment Method for the Intensive Care Unit (CAM-ICU). *Critical Care Medicine*. 29(7), 1370-9.
11. Fischer J, Mathieson C. (2001). The history of the Glasgow Coma Scale: implications for practice. *Crit Care Nurse*, 23, 52-8.
12. Gabbe BJ, Cameron PA, Finch CF. (2003). The status of the Glasgow Coma Scale. *Emerg Med*, 15, 353-60.
13. Gift AG. (1989). Validation of a vertical visual analogue scale as a measure of clinical dyspnea. *Rehabil Nurs*, 14(3), 23-5.

14. Hogan-Miller, E., Rustad, D., Sendelbach, S., & Goldenberg, I. (1995). Effects of three methods of femoral site immobilization on bleeding and comfort after coronary angiogram. *American Journal of Critical Care : An Official Publication, American Association of Critical-Care Nurses*, 4(2), 143-148.
15. Hogarth, D. K., & Hall, J. (2004). Management of sedation in mechanically ventilated patients. *Current Opinion in Critical Care*, 10(1), 40-46.
16. Hurley, A. C., Volicer, B. J., Hanrahan, P. A., Houde, S., & Volicer, L. (1992). Assessment of discomfort in advanced alzheimer patients. *Research in Nursing & Health*, 15(5), 369-377.
17. Inouye SK, van Dyck CH, Alessi CA, et al.(1990). Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med*,113(12), 941-8.
18. Kolcaba, K. (2003).*コルカバ コンフォート理論 理論の開発過程と実践への適用*.太田喜久子監訳(2008).医学書院.
19. Kolcaba, K. Y. (1992). Holistic comfort: Operationalizing the construct as a nurse-sensitive outcome. *ANS.Advances in Nursing Science*, 15(1), 1-10.
20. Lora A.Bear.& Peggy. Ward-Smith.(2006).Interrater Reliability og the COMFORT Scale.*Pediatric Nursing*,32(5), 427-434.
21. Marx, C. M., Smith, P. G., Lowrie, L. H., Hamlett, K. W., Ambuel, B., Yamashita, T. S., et al. (1994). Optimal sedation of mechanically ventilated pediatric critical care patients. *Critical Care Medicine*, 22(1), 163-170.
22. McGaffigan, P. A. (2002). Advancing sedation assessment to promote patient comfort. *Critical Care Nurse*, Suppl, 29-36.
23. Novak, B., Kolcaba, K., Steiner, R., & Dowd, T. (2001). Measuring comfort in caregivers and patients during late end-of-life care. *The American Journal of Hospice & Palliative Care*, 18(3), 170-180.
24. Payen, J., Chanques, G., Mantz, J., Hercuel, C., Auriant, I., Leguillou, J., Binhas, M., Genty, C., Rolland, C., Bosson, J.(2007).Current practices in sedation and analgesia for mechanically ventilated critically ill patients: a prospective multicenter patient based study. *Anesthesiology*,107,687-695.
25. Polit,D. F., Beck, C. T. (2004).*看護研究 原理と方法 第2版*. 近藤潤子監訳(2010). 医学書院. p238.
26. Puntillo, K., Miaskowski, C., Kehrle,K., Stannard, D., Gleeson, S., Nye, P.(1997). Relationship between behavioral and physiological indicators of pain, critical care patients' self-reports of pain, and opioid administration. *Critical Care Medicine*, 25,

1159-1166.

27. Sessler CN, Gosnell MS, Grap MJ, Brophy GM, O'Neal V, Keane KA, et al.(2002) The Richmond Agitation-Sedation Scale. Validity and reliability in adult intensive care unit patients. *Am J Respir Crit Care Med*, 166, 1338-44.
28. Teasdale G, Jennett B.(1974). Assessment of coma and impaired consciousness. *Lancet*,13, 81-3.
29. van Dijk, M., de Boer, J. B., Koot, H. M., Tibboel, D., Passchier, J., & Duivenvoorden, H. J. (2000). The reliability and validity of the COMFORT scale as a postoperative pain instrument in 0 to 3-year-old infants. *Pain*, 84(2-3), 367-377.
30. Wielenga,L.M.,De Vos,R.,de Leeuw,R.,& De Haan,R.J.(2004).COMFORT Scale:A reliable and valid method to measure the amount stress of ventilated preterm infants. *Nonatal Network; The Journal of Neonatal Nursing*,23,39-44.
31. Wild, D., Grove, A., Martin, M., Eremenco, S., McElroy, S., Verjee-Lorenz, A., et al. (2005). Principles of good practice for the translation and cultural adaptation process for patient-reported outcomes (PRO) measures: Report of the ISPOR task force for translation and cultural adaptation. *Value in Health : The Journal of the International Society for Pharmacoeconomics and Outcomes Research*, 8(2), 94-104.
32. Young, J., Siffleet, J., Nikoletti, S., Shaw, T.(2006).Use of behavioral Pain Scale to assess pain in ventilated, unconscious and/or sedated patients. *Intensive & critical care Nursing*,22,32-39.
33. 佐居由美. (2004). 看護実践場面における「安楽」という用語の意味するもの. 聖路加看護大学紀要, (30), 1-9.
34. 佐居由美. (2005). 和文献にみる「安楽」と英文献にみる「comfort」の比較 Rodgers の概念分析の方法を用いている日米 2 つの看護文献レビューから. 聖路加看護大学紀要, (31), 1-7.
35. 縄秀志. (2006). 看護実践における"Comfort"の概念分析. 聖路加看護学会誌, 10(1), 11-22.
36. 井上智子.(2007).クリティカルケア看護概論. 寺町優子編,クリティカルケア看護 理論と臨床への応用.(pp.1-15).東京:日本看護協会出版会.
37. 日本集中治療医学会. <http://www.jsicm.org/> [2011-04-15]
38. 厚生労働省. <http://www.mhlw.go.jp/toukei/saikin/hw/iryosd/08/index.html> [2011-04-20]