

## References

- Arai, Y. & Takahashi, M. (2006). Shusankino kazokukinouto ninsanjokuhuno mentaruherusu [Family processes in the perinatal period and pregnant women's mental health]. *Kitasato International Journal of Nursing Science*, 8(1), 30-37. (in Japanese).
- Ayabe, T. (2011). Abnormal labor. In T. Okai & T. Ayabe (Eds.), *Hyoujun sanko hujinkagaku [Standards of obstetric and gynecology (4th ed)]*. (pp. 469-505). Tokyo: Igakushoin. (in Japanese).
- Baby-Friendly Hospital Initiative (1989). Retrieved December 4, 2011, from <http://www.unicef.org/programme/breastfeeding/baby.htm>
- Beck, C. (2001). Predictors of postpartum depression. *Nursing Research*, 50(5), 275-285.
- Biro, M., Waldenstrom, U. & Pannifex, J. (2000). Team midwifery care in a tertiary level obstetric service: A randomized controlled trial. *BIRTH*, 27(3), 168-173.
- Biro, M., Waldenstrom, U., Brown S. & Pannifex J. (2003). Satisfaction with team midwifery care for low- and high-risk women: A randomized controlled trial. *BIRTH*, 30(1), 1-10.
- Bondas, T. (2002). Finnish women's experience of antenatal care. *Midwifery*, 18(1), 61-71.
- Bottorff, J., Balneaves, L., Sent, L., Grewal, S. & Browne, A. J. (2001). Cervical cancer screening in ethno cultural groups: Case studies in women-centered care. *Women Health*, 33(3-4), 29-46.
- Brown, S., Davey, M. & Bruinsma, F. (2000). Women's views and experiences of postnatal hospital care in the Victorian Survey of Recent Mothers 2000. *Midwifery*, 21(2), 109-126.
- Cox, J., Holden, J. & Sagovsky, R. (1987). Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry*, 150, 782-786.

- Davey, M., Brown, S. & Bruinsma, F. (2005). What is it about antenatal continuity of caregiver that matters to women? *BIRTH*, 32(4), 262-271.
- Department of Health (1993). Changing Childbirth. Part I: Report of the Expert Maternity Group. London: HMSO.
- Endo, K., Waki, M., Yamakawa, Y., Komatsu, R., Hori, M., Maki, K., Misawa, T. & Kawasaki, K. (2008). Sanjokusoukiniokeru sangoutubyouhasshouno yosokuinshi [Predictors of postpartum depression in the early postpartum stage]. *Yamagata Journal of Health Science*, 11, 1-8. (in Japanese).
- Escriba-Aguir, V. & Artazcoz, L. (2011). Gender differences in postpartum depression: A longitudinal cohort study. *Journal of Epidemiology and Community Health*, 65, 320-326.
- Evans, J., Heron, J., Francomb, H. & Oke, S. (2001). Cohort study of depressed mood during pregnancy after childbirth. *BMJ*, 323, 257-260.
- Gepshtein, Y., Horiuchi, S. & Eto, H. (2007). Independent Japanese midwives: A qualitative study of their practice and beliefs. *Japan Journal of Nursing Science*, 4(2), 85-93.
- Giarratano, G. (2003). Woman-centered maternity nursing education and practice. *Journal of Perinatal Education*, 12(1), 18-28.
- Hasegawa, H. & Murakami, A. (2005). Shussansurujoseiga manzokudekiru osan-josaninno shussantaikenno-tokarano bunseki [Satisfactory childbirth: Analysis of women's notes from a midwifery institute]. *Japanese Journal of Maternal Health*, 45(4), 12(1), 18-28. (in Japanese).
- Hatem, M., Sandall, J., Devane, D., Soltani, H. & Gates, S. (2008). Midwife-led versus other models of care for childbearing women. *Cochrane Database System of Systematic Reviews*, Issue 4. Art. No.: CD004667. DOI: 10.1002/14651858.CD00667.pub2.
- Henley-Einion, A. (2008). Midwifery basics: Caring for women with medical conditions (2). Respiratory disorders in pregnancy. *The Practicing Midwife*, 11(10), 48-53.
- Hicks, C., Spurgeon, P. & Barwell, F. (2003). Changing childbirth: A pilot project. *Journal of Advanced Nursing*, 42(6), 617-628.

- Higuma, H., Yoshikawa, Y., Takeda, Y. & Terao, A. (2000). Josaninsentakujoseino shussantaikenno bunseki [Analysis of women's experience of childbirth at maternity centers]. *Annual reports of School of Health Sciences Faculty of Medicine, Kyoto University, 20*, 45-53. (in Japanese).
- Hildingsson, I. & Thomas, J. (2007). Women's perspectives on maternity services in Sweden: Processes, problems, and solutions. *Journal of Midwifery & Women's Health, 52*(2), 126-133.
- Hills, M. & Mullett, J. (2002). Women-centred care: Working collaboratively to develop gender inclusive health policy. *Health Care for Women International, 23*(1), 84-97.
- Hills, M. & Mullett, J. (2005). Primary health care: A preferred health service delivery option for women. *Health Care for Women International, 26*(4), 325-339.
- Hindley, C., Hinsliff, S. & Thomson, A. (2006). English midwives' views and experiences of intrapartum fetal heart rate monitoring in women at low obstetric risk: Conflicts and compromises. *Journal of Midwifery & Women's Health, 51*(5), 354-360.
- Hiruta, Y., Saito, S. & Suehara, K. (2002). Byouinto josanjoniokeru ninsanpukeano jittai (ge) [The actual status of care provided at hospitals and birth centres]. *The Japanese Journal for Midwives, 56*(5), 421-424.
- Hodnett, E. (2000). Continuity of caregivers for care during pregnancy and childbirth. *Cochrane Database of Systematic Reviews*, Issue 1. Art. No.: CD000062. DOI: 10.1002/14651858.CD000062.
- Holzemer, W. (1994). The impact of nursing care in Latin America and the Caribbean: A focus on outcomes. *Journal of Advanced Nursing, 20*, 5-12.
- Homer, C., Davis, G., Cooke, M. & Barclay, L. (2002). Women's experiences of continuity of midwifery care in a randomized controlled trial in Australia. *Midwifery, 18*(2), 102-112.
- Horiuchi, S., Shimada, K., Suzuki, M., Mori, T., Taniguchi, M., Taga, Y. & Miyasato, K. (1997). Shussanwoshitajoseiga hyoukasuru ninsanjokukino keanoshitu [Women's evaluation of quality of care during pregnancy, delivery, and postpartum period]. *Journal of Japan Academy of Midwifery, 11*(1), 9-16. (in Japanese).

- Horiuchi, S., Kataoka, Y., Eto, H., Oguro, M. & Mori, T. (2006). The applicability of women-centered care: Two case studies of capacity-building for maternal health through international collaboration. *Japan Journal of Nursing Science*, 3, 143-150.
- Horiuchi, S., Yaju, Y., Kataoka, Y., Eto, H. G. & Matsumoto, N. (2009). Development of an evidence-based domestic violence guideline: Supporting perinatal women-centred care in Japan. *Midwifery*, 25, 72-78.
- Iida, M. (2009a). Toujishaniyoru ninshinchuno joseiwo chushintoshitakeano ninshiki [Women's perception of receiving 'women-centered care' during pregnancy]. Unpublished master's thesis, St. Luke's College of Nursing, Tokyo-Japan. (in Japanese).
- Iida, M. (2009b). Women-centred care no gainen bunseki ['Women-centred care'- a concept analysis]. Unpublished manuscript, St. Luke's College of Nursing Tokyo, Japan. (in Japanese)
- Iida, M. (2010). Joseiwo chushintoshitakea-ninshinkishakudono kaihatuto sonodatouseino kentou [Developing and testing of an instrument to assess women-centered care during pregnancy]. *Journal of Japan Academy of Midwifery*, 24(2), 284-293. (in Japanese).
- Iida, M. (2011). A review of the methodological features of randomized controlled trials in midwifery-led care- what are the threats to validity? Manuscript submitted for publication.
- Iida, M., Horiuchi, S. & Porter, S. (2011). Women's perceptions of receiving women-centred care during pregnancy: A comparison among birth centres, clinics, and hospitals in Japan. *Midwifery*, doi:10.1016/j.midw.2011.07.002
- Ishii, H. (2007). Toukeibunsekino kokogashiritai- hoken, kango, shinri, kyouikukeikenkyuno matomekata [Learning statistical analysis. 1<sup>st</sup> ed]. Tokyo: Bunkodo. (in Japanese).
- Iwatani, S., Naruse, E., Yoshikawa, T. & Yamakawa, M. (2000). Sangomikkameto ikkagetumeno jokuhuno seishinjoutaino judanteki chousa [A longitudinal research of women's psychological condition at three days postpartum and one month postpartum]. *Japanese Journal of Maternal Health*, 41(2), 347-355. (in Japanese).

- Jackson, K. & Toda, R. (1998). Changing childbirth, Keynote speech from the 24<sup>th</sup> ICM Conference, Oslo. *The Japanese Journal for Midwives*, 52(4), 298-303. (in Japanese).
- Japan's government statistics: e-statics (2011). Seihutoukeino sougoumadoguchi [Government statistics]. Retrieved March 2, 2012, from <http://www.e-stat.go.jp/SG1/estat/List.do?lid=000001082326> (in Japanese).
- Japan Ministry of Health, Labour and Welfare (2006). Heisei17nendo nyuyouji eiyouchousakekkano gaiyou [Survey of the nutrition of infants and preschool children, 2005]. Retrieved December 2, 2011, from <http://www.mhlw.go.jp/houdou/2006/06/h0629-1.html> (in Japanese).
- Japan Ministry of Health, Labour and Welfare (2008). Anshintou kibouno iryoukakuho bijon [A vision to secure peace and hope]. Retrieved April 5, 2011, from <http://www.mhlw.go.jp/shingi/2008/06/dl/s0618-8a.pdf> (in Japanese).
- Japan Ministry of Health, Labour and Welfare (2009). Iryoushitsu (seitai/doutai) chousa, byouinhoukokuno gaiyou [Conspectus of health facilities (static/dynamic) and hospitals report]. Retrieved April 5, 2011, from <http://www.mhlw.go.jp/toukei/saikin/hw/iryosd/08/index.html> (in Japanese).
- Japanese Nursing Association (2009). Heisei20nendo josansenta-no sechisuishin purojekutotoushin [Report of the project of promoting inside midwifery units (2008)]. Retrieved October 27, 2011, from <http://www.nurse.or.jp/home/innaijyosan/pdf/toshin.pdf> (in Japanese).
- Kataoka, Y. (2005). Joseinitaisuru bouryoku sukuri-ningushakudono kaihatu [Development of the violence against women screen]. *Journal of Japan Academy of Nursing Science*, 25(3), 51-60. (in Japanese).
- Kennedy, H. (1995). The essence of nurse-midwifery care-the woman's story. *Journal of Nurse-Midwifery*, 40(5), 410-417.

- Kitamura, T., Okamura, S., Takeda, N., Hujita, J., Kamisato, T., Sugiyama, T., Kusaka, H., Sato, A., Hukushima, K., Ohoba, T., Matui, K., Kanazawa, K., Okano, T., Yoshida, K., Yamashita, H., Arai, Y. & Nakano, H. (2005). Josanshiniyoru ninshinkikanchuno shinrishienga sangoutubyouno jushoudoni ataeru yoboukoukani kansuru kenkyu [The effect of psychological support from midwives in the antenatal period that impacts the prevention of the severity of postpartum depression]. *2005 Summary Shared research bulletin. Health and Labour Sciences Research Grant. Japan Child and Family Research Institute Bulletin*, 2-11. (in Japanese).
- Lenihan, J. P. (1984). Relationship of antepartum pelvic examinations to premature rupture of the membranes. *Obstetrics & Gynecology*, 83(1), 33-37.
- Luyben, A. & Fleming, V. (2005). Women's needs from antenatal care in three European countries. *Midwifery*, 21, 212-223.
- Lyberg, A. & Severinsson, E. (2010). Fear of childbirth: Mothers' experiences of team-midwifery care - a follow-up study. *Journal of Nursing Management*, 18, 383-390.
- McCourt, C & Pearce, A. (2000). Does continuity of carer matter to women from minority ethnic groups? *Midwifery*, 16(2), 145-154.
- Morgan, M., Fenwick, N., McKenzie, C. & Wolfe, C.D. (1998). Quality of midwifery led care: assessing the effects of different models of continuity for women's satisfaction. *Quality in Health Care*, 7(2), 77-82.
- Mothers' & Children's Health Organization (2011). Boshihokenno omonarutoukei [Maternal and Child Health Statistics of Japan]. 131-133. Tokyo. (in Japanese)
- Meadows, L., Thurston, W., Quantz, D. & Bobey, M. (2006). Women's health resources: Facilitating a community of care for midlife women. *Promotion & Education*, 13(1), 42-46.
- Meddings, F., Phipps, F., Haith-Cooper, M. & Haigh, J. (2007). Vaginal birth after caesarean section (VBAC): Exploring women's perceptions. *Journal of Clinical Nursing*, 16(1), 160-167.

- Misago, C., Umenai, T., Noguchi, M., Mori, T. & Mori, T. (2000). Satisfying birth experiences in Japan. *Lancet*, 355, 2256.
- Misago, C. (2008). Joseino seikatushukanto kenkouni kansuru ekigakukenkyu-Tokyotono josanjoniokeru ninpuwo taishoutoshita judankenkyuyori [An epidemiology research of women's lifestyle habit and health: From a longitudinal study including women receiving care at birth centres in Tokyo]. *The report in 2008 from International medical aid research expenses for commission*. (in Japanese).
- Misago, C., Takehara, K., Matumoto, A., Moriya, M. & Minoura, S. (2009). Ninshin, shussanzengono joseino seikatushukannohenka josanjode shussanwosuru joseiwo taishoutoshita tusekichousayori [Women's lifestyle change during pregnancy and before and after delivery: A follow-up research of women who delivered at birth centres]. *Japanese Journal of Maternal Health*, 50(3), 237. (in Japanese).
- Nakada, K. (2008). Bonyuikujino keizokuni eikyosuruyouinto hahayano seruhuefikashi-tono kanren [An analysis of maternal self-efficacy and breastfeeding continuation]. *Journal of Japan Academy of Midwifery*, 22(2), 208-221. (in Japanese).
- Nakamura, S. (2011). Ninpuno hieshoga motarasu ijoubunbennokaimeikeikousukoaniyoru kourakuinshinochousei [Explication of abnormal delivery caused by pregnant women's sensitivity to cold (*hiesho*) - Adjustment of confounding factors by using propensity scores]. Unpublished doctoral dissertation, St. Luke's College of Nursing, Tokyo-Japan. (in Japanese).
- Nakano, Y. (1994). Ninsanpuwotorimaku shoyouinto boshinokenkouni kansurukenkyu [Research of causes surrounding parturient and mother and child health]. *Ministry of Health & Welfare, Psychosomatic disorder research*, 7-10. (in Japanese).
- National Institute for Clinical Excellence (NICE) (2007). Intrapartum care: Care of the healthy women and their babies during childbirth. Retrieved December 10, 2011, from <http://www.nice.org.uk/nicemedia/pdf/CG55FullGuideline.pdf>

- National Institute for Clinical Excellence (NICE) (2008). Antenatal care: Routine care for the healthy pregnant woman. Retrieved September 23, 2011, from <http://www.nice.org.uk/nicemedia/live/11947/40145/40145.pdf>
- Okano, T., Nomura, J., Makita, I., Makita, A. & Yamaguchi, T. (1989). Maternity bluesno rinshounaibunpituteki kenkyu [Clinical endocrinology research of maternity blues]. *Clinical Psychiatry*, *31*(7), 725-783. (in Japanese).
- Okano, T., Nomura, J., Koshikawa, N., Doi, M. & Tatumuma, T., (1991). Maternity bluesto sangoutubyouno hikakubunkateki kenkyu [Cultural research comparing maternity blues and postpartum depression]. *Clinical Psychiatry*, *33*(10), 1051-1058. (in Japanese).
- Okano, T. (1993). Honpouniokeru sangoseishinshougaikenkyuno jittai [Status of Japan's postpartum depression research]. *Perinatal Medicine (Tokyo)*, *23*(10), 1397-1404. (in Japanese).
- Okano, T., Murata, M., Masuji, S., Tamaki, R., Nomura, J., Miyaoka, H. & Kitamura, T. (1996). Nihonban Ejinbarasangoutubyou jikohyoukahyou(EPDS)no shinraiseito datousei [Testing reliability and validity of the Japanese version of Edinburgh Postnatal Depression Scale]. *Archives of Psychiatric Diagnostics and Clinical Evaluation*, *7*(4), 525-533. (in Japanese).
- Osbourne, A. (2005). Happy midwives, happy mothers. Interview by Lynne Pearce. *Nursing Standard*, *19*(47), 20-21.
- Page, L. (1996). *Umarekawaru josanputachi* [Effective group practice in midwifery: Working with women] (Sano, N., Aono, Y., Sato, K. & Takeuchi, M., Trans.). Tokyo: Igakushoin. (Original work published 1995) (in Japanese).
- Pope, R., Granham, L. & Patel, S. (2001). Women-centered care. *International Journal of Nursing Studies*, *38*, 227-238.
- Sato, S., Sato, R., Sato, K., Katakura, M., Abe, E. & Sasaki, M. (2002). Jokuhunohuanbunbenyoushikibetuni kangaeru [The change in the mental condition of puerperal women: the effect of delivery style]. *Bulletin of School of Health Sciences Tohoku University*, *11*(2), 195-205. (in Japanese).



- Sato, K. & Sato, S. (2010). Ninshinkikarano keizokushita shinritekishienga shusankijoseino huan, yokuutuni oyobosukouka [Effect of continuous psychological support during pregnancy on anxiety and depression among women during the perinatal period]. *Japanese Journal of Maternal Health*, 51(1), 215-225. (in Japanese).
- Shields, N., Turnbull, D., Reid, M., Holmes, A., McGinley, M. & Smith, L. N. (1998). Satisfaction with midwife-managed care in different time periods: A randomized controlled trial of 1299 women. *Midwifery*, 14, 85-93.
- Stein, G. (1980). The pattern of mental change and body weight change in the first post-partum week. *Journal of Psychosomatic Research*, 24, 165-171.
- Suzuki, K., Omachi, H., Mizutani, S. & Matsuo, T. (2003). What women desire for birth: A survey from birth centres. *Japanese Journal of Maternal Health*, 44(1), 98-104.
- Takehara, K., Kitamura, N., Misago, C. & Minoura, S. (2008). 'Keizokukea'toha donoyouna keananoka [What is 'continuity of care'?]. *The Japanese Journal for Midwives*, 62(5), 443-446. (in Japanese).
- Takehara, K., Okamoto, N., Yoshiasa, K., Misago, C., Osanai, Y., Okamoto, K. & Minoura, S. (2009). Josanjode ninpunitaishite jishhisareteiru keanikansuru shitutekikenkyu- josanjono keanohonshitutoha douiumonoka [A qualitative study of care provided for pregnant women at birth houses in Japan: What is the essence of midwifery care at birth houses?]. *Japanese Journal of Maternal Health*, 50(1), 190-198. (in Japanese).
- Turnbull, D., Holmes, A., Shields, N., Cheyne, H., Twaddle, S., Gilmour, W. H., McGinley, M., Reid, M., Johnstone, I., Geer, I., McIlwaine, G. & Lunan, C. B. (1996). Randomized controlled trial of efficacy of midwife-managed care. *Lancet*, 348, 213-218.
- Waldenstrom, U., Brown, S., McLachlan, H., Forster, D. & Brennecke, S. (2000). Does team midwife care increase satisfaction with antenatal, intrapartum, and postpartum care? A randomized controlled trial. *BIRTH*, 27(3), 156-167.

- Waldenstrom, U., McLachlan, H., Forster, D., Brennecke, S. & Brown, S. (2001). Team midwife care: Maternal and infant outcomes. *The Australian & New Zealand Journal of Obstetrics & Gynaecology*, 41(3), 257-64.
- Yamashita, H. (1994). Mateniti-buru-zuno honpouniokerujittaito sonotaisaku-Jittaichousakekkato sukuri-ningushakudono kentou, Ninsanpuwotorimaku shoyouinto boshinokenkouni kansurukenkyu [Japan's actual status of maternity blues and countermeasure: Structuring the diagnostic criteria and the screening system, Ministry of Health & Welfare Psychosomatic disorder research]. *Research of causes surrounding parturient and mother and child health*, 26-32. (in Japanese).
- Yamashita, H. & Yoshida, K. (2002). Sanjokukino seishinshougaito seishinmenshien [Psychiatric disorders and mental support during the postpartum period]. *Perinatal Medicine (Tokyo)*, 32(1), 67-72. (in Japanese).
- Yuan, K. & Bentler, P. (2007). Structural Equation Modeling. In Rao, C.R. & Sinharay, S. (Eds.), *Handbook of Statistics*, Vol. 26 (pp. 297-358). Amsterdam: Elsevier.