

引用文献

- Adams, K. F., Jr, Fonarow, G. C., Emerman, C. L., LeJemtel, T. H., Costanzo, M. R., Abraham, W. T., . . . ADHERE Scientific Advisory Committee and Investigators. (2005). Characteristics and outcomes of patients hospitalized for heart failure in the united states: Rationale, design, and preliminary observations from the first 100,000 cases in the acute decompensated heart failure national registry (ADHERE). *American Heart Journal*, 149(2), 209-216. doi:S0002870304004995 [pii]
- 安達 仁. (2016). 【変わりつつある高齢者心不全の考え方と治療法】 フレイルと心臓リハビリテーション. *Geriatric Medicine*, 54(7), 663-667.
- Albert, N. M., Levy, P., Langlois, E., Nutter, B., Yang, D., Kumar, V. A., . . . Nykun, E. (2014). Heart failure beliefs and self-care adherence while being treated in an emergency department. *The Journal of Emergency Medicine*, 46(1), 122-129. doi:10.1016/j.jemermed.2013.04.060 [doi]
- Anderson, E. S., Wojcik, J. R., Winett, R. A., & Williams, D. M. (2006). Social-cognitive determinants of physical activity: The influence of social support, self-efficacy, outcome expectations, and self-regulation among participants in a church-based health promotion study. *Health Psychology : Official Journal of the Division of Health Psychology, American Psychological Association*, 25(4), 510-520. doi:2006-08842-008 [pii]
- Angus, D. C., Musthafa, A. A., Clermont, G., Griffin, M. F., Linde-Zwirble, W. T., Dremsizov, T. T., & Pinsky, M. R. (2001). Quality-adjusted survival in the first year after the acute respiratory distress syndrome. *American Journal of Respiratory and Critical Care Medicine*, 163(6), 1389-1394.
- 荒井 秀典.(2014).フレイルの意義. *日本老年医学会雑誌*, 51(6), 497-501.

- 麻野井 英次. (1997). 慢性心不全患者の運動制限要因:Physical deconditioning の重要性. *日本老年医学会雑誌*, 34(10), 845.
- 麻野井 英次.(2010).総論自律神経系による循環調節.井上博(編), 循環器疾患と自律神経機能(第2版)(pp32).医学書院.2010.
- 東 あかね, 八城 博子, 清田 啓介, & 他. (1996). 消化器内科外来における hospital anxiety and depression scale (HAD 尺度)日本語版の信頼性と妥当性の検討. *日本消化器病学会雑誌*, 93(12), 884-892.
- Baratz, D. M., Westbrook, P. R., Shah, P. K., & Mohsenifar, Z. (1992). Effect of nasal continuous positive airway pressure on cardiac output and oxygen delivery in patients with congestive heart failure. *Chest*, 102(5), 1397-1401.
- Bandura, A. (2004). Health promotion by social cognitive means. *Health Education & Behavior : The Official Publication of the Society for Public Health Education*, 31(2), 143-164. doi:10.1177/1090198104263660 [doi]
- Bloomfield, S. A. (1997). Changes in musculoskeletal structure and function with prolonged bed rest. *Medicine and Science in Sports and Exercise*, 29(2), 197-206.
- Brummel, N. E., Jackson, J. C., Girard, T. D., Pandharipande, P. P., Schiro, E., Work, B., . . . Ely, E. W. (2012). A combined early cognitive and physical rehabilitation program for people who are critically ill: The activity and cognitive therapy in the intensive care unit (ACT-ICU) trial. *Physical Therapy*, 92(12), 1580-1592. doi:10.2522/ptj.20110414 [doi]
- Chen, J., Ross, J. S., Carlson, M. D., Lin, Z., Normand, S. L., Bernheim, S. M., . . . Krumholz, H. M. (2012). Skilled nursing facility referral and hospital readmission rates after heart failure or myocardial infarction. *The American Journal of Medicine*, 125(1), 100.e1-100.e9. doi:10.1016/j.amjmed.2011.06.011 [doi]

- Clark, A. L., Poole-Wilson, P. A., & Coats, A. J. (1996). Exercise limitation in chronic heart failure: Central role of the periphery. *Journal of the American College of Cardiology*, 28(5), 1092-1102. doi:S0735-1097(96)00323-3 [pii]
- Cleland, J. G., Swedberg, K., Follath, F., Komajda, M., Cohen-Solal, A., Aguilar, J. C., . . . Study Group on Diagnosis of the Working Group on Heart Failure of the European Society of Cardiology. (2003). The EuroHeart failure survey programme - a survey on the quality of care among patients with heart failure in europe. part 1: Patient characteristics and diagnosis. *European Heart Journal*, 24(5), 442-463. doi:S0195668X02008230 [pii]
- Collin, C., Wade, D. T., Davies, S., & Horne, V. (1988). The barthel ADL index: A reliability study. *International Disability Studies*, 10(2), 61-63.
- De Jonghe, B., Bastuji-Garin, S., Sharshar, T., Outin, H., & Brochard, L. (2004). Does ICU-acquired paresis lengthen weaning from mechanical ventilation? *Intensive Care Medicine*, 30(6), 1117-1121. doi:10.1007/s00134-004-2174-z [doi]
- Ferrando, A. A., Paddon-Jones, D., & Wolfe, R. R. (2006). Bed rest and myopathies. *Current Opinion in Clinical Nutrition and Metabolic Care*, 9(4), 410-415. doi:10.1097/01.mco.0000232901.59168.e9 [doi]
- Franciosa, J. A., Park, M., & Levine, T. B. (1981). Lack of correlation between exercise capacity and indexes of resting left ventricular performance in heart failure. *The American Journal of Cardiology*, 47(1), 33-39.
- Galper, D. I., Trivedi, M. H., Barlow, C. E., Dunn, A. L., & Kampert, J. B. (2006). Inverse association between physical inactivity and mental health in men and women. *Medicine and Science in Sports and Exercise*, 38(1), 173-178. doi:00005768-200601000-00027 [pii]

- Georgiadou, P., & Adamopoulos, S. (2012). Skeletal muscle abnormalities in chronic heart failure. *Current Heart Failure Reports*, 9(2), 128-132. doi:10.1007/s11897-012-0090-z; 10.1007/s11897-012-0090-z
- 後藤 杏里, 西 将則, & 猪飼 哲夫. (2007). 【内部疾患のリハビリテーション 効果と実際】 廃用症候群について. *Medical Rehabilitation*, (82), 72-78.
- 後藤 葉一. (2009). 心臓リハビリテーション エビデンスと展望. *日本心臓病学会誌*, 3(3), 195-215.
- Granger, C. V., Greer, D. S., Liset, E., Coulombe, J., & O'Brien, E. (1975). Measurement of outcomes of care for stroke patients. *Stroke; a Journal of Cerebral Circulation*, 6(1), 34-41.
- Granger, C. V., Albrecht, G. L., & Hamilton, B. B. (1979). Outcome of comprehensive medical rehabilitation: Measurement by PULSES profile and the barthel index. *Archives of Physical Medicine and Rehabilitation*, 60(4), 145-154.
- Guck, T. P., Elsasser, G. N., Kavan, M. G., & Barone, E. J. (2003). Depression and congestive heart failure. *Congestive Heart Failure (Greenwich, Conn.)*, 9(3), 163-169.
- 廃用症候群.(2015).医学大辞典 (第 20 版) (pp1941-1942).南山堂.
- 原 清江, 三井 彩子, & 内田 緑. (2014). 入退院を繰り返す高齢独居心不全患者の退院支援の有効性. *信州大学医学部附属病院看護研究集録*, 42(1), 19-23.
- 春名 由一郎, 李 廷秀, & 川久保 清. (1999). 【在宅高齢者のケア】 長期臥床による影響. *保健の科学*, 41(11), 824-828.
- Herridge, M. S., Cheung, A. M., Tansey, C. M., Matte-Martyn, A., Diaz-Granados, N., Al-Saidi, F., . . . Canadian Critical Care Trials Group. (2003). One-year outcomes

in survivors of the acute respiratory distress syndrome. *The New England Journal of Medicine*, 348(8), 683-693. doi:10.1056/NEJMoa022450 [doi]

平木 幸治, 井澤 和大, 渡辺 敏, 森尾 裕志, 笠原 西介, 長田 尚彦, ... 飯島 節. (2011). 糖尿病を合併した急性心筋梗塞患者の運動耐容能低下の関連要因. *理学療法学*, 38(5), 343-350.

Holzemer William L. (1988). 評価研究の実施過程 研究の方法論 標本の抽出と測定用具. *看護研究*, 22(1), 61-67.

Horacio J. Adrogué & Donald E. Wesson (1998). 心不全-病態生理からマネジメントまで-. 207. 後藤葉一監訳. 東京: 医学書院 207.

石川 朗, 宮坂 智哉, 長谷 陽子, & 高橋 尚明. (2004). 【脱・安静の弊害 一般病棟で実践したい早期回復援助の技術】 "安静の弊害"を検証する. *看護*, 56(10), 40-43.

伊東 春樹. (2010). 運動負荷試験で診る慢性心不全の病態生理. *心臓リハビリテーション*, 15(2), 214-217.

Izawa Kazuhiro P., Watanabe Satoshi, & Oka Koichiro. (2015). Relationship of thresholds of physical performance to nutritional status in older hospitalized male cardiac patients. *Geriatrics & Gerontology International*, 15(2), 189-195.

井澤 和大, 渡辺 敏, 岡 浩一朗, 平木 幸治, 森尾 裕志, 笠原 西介, ... 三宅 良彦. (2010). 慢性心不全患者における重症度別の身体活動と運動能力指標. *心臓リハビリテーション*, 15(1), 134-138.

Junger, J., Schellberg, D., Muller-Tasch, T., Raupp, G., Zugck, C., Haunstetter, A., ... Haass, M. (2005). Depression increasingly predicts mortality in the course of congestive heart failure. *European Journal of Heart Failure*, 7(2), 261-267. doi:S1388-9842(04)00174-6 [pii]

絹川 真太郎, & 沖田 孝一. (2011). 【サルコペニアの診かた】 各種疾患と骨格筋障害
心不全と骨格筋障害. *Modern Physician*, 31(11), 1341-1346.

北村俊則.(2013).第 14 章欠損値に対応しよう.多重代入法とその他の代入法.臨床で働きな
がら研究をしよう(pp.25-51).北村メンタルヘルス研究所.

http://www.procomu.jp/jspd2016/pdf/yougo_kaisetu04.pdf#search=%27%E5%A4%9A%E9%87%8D%E4%BB%A3%E5%85%A5%E6%B3%95%27

Ko, J. K., & McKelvie, R. S. (2005). The role of exercise training for patients with heart failure. *Europa Medicophysica*, 41(1), 35-47.

Kortebein, P., Ferrando, A., Lombeida, J., Wolfe, R., & Evans, W. J. (2007). Effect of 10 days of bed rest on skeletal muscle in healthy older adults. *JAMA : The Journal of the American Medical Association*, 297(16), 1772-1774. doi:297/16/1772-a [pii]

Levine, T. B., Francis, G. S., Goldsmith, S. R., & Cohn, J. N. (1983). The neurohumoral and hemodynamic response to orthostatic tilt in patients with congestive heart failure. *Circulation*, 67(5), 1070-1075.

Liesching, T., Kwok, H., & Hill, N. S. (2003). Acute applications of noninvasive positive pressure ventilation. *Chest*, 124(2), 699-713.

Loewen, S. C., & Anderson, B. A. (1988). Reliability of the modified motor assessment scale and the barthel index. *Physical Therapy*, 68(7), 1077-1081.

Loretsen, E., Bay, G., Grendahl, H., & Sivertssen, E. (1967). The hemodynamic effect of changes in posture in cardiac patients. *Acta Medica Scandinavica*, 182(2), 175-182.

Macauley K. (2012). Physical therapy management of two patients with stage d heart failure in the cardiac medical intensive care unit. *Cardiopulm.Phys.Ther.J.*, 23(3), 37-45.

- MAHONEY, F. I., & BARTHEL, D. W. (1965). Functional evaluation: The barthel index. *Maryland State Medical Journal*, 14, 61-65.
- Mancini, D. M., Eisen, H., Kussmaul, W., Mull, R., Edmunds, L. H., Jr, & Wilson, J. R. (1991). Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation*, 83(3), 778-786.
- 慢性心不全治療ガイドライン(2010 年改訂版).http://www.j-circ.or.jp/guideline/pdf/JCS2010_matsuzaki_h.pdf#search=%27%E6%85%A2%E6%80%A7%E5%BF%83%E4%B8%8D%E5%85%A8+%E3%82%AC%E3%82%A4%E3%83%89%E3%83%A9%E3%82%A4%E3%83%B3%27
- 松村 美佳, 改發 麻優, 中野 ひとみ, 小林 希望, & 原田 友香. (2014). 心不全患者に対するセルフケア確立に向けての指導効果の検証. *愛仁会医学研究誌*, 45, 96-99.
- Muller, E. A. (1970). Influence of training and of inactivity on muscle strength. *Archives of Physical Medicine and Rehabilitation*, 51(8), 449-462.
- Murberg, T. A., Bru, E., Svebak, S., Tvetaras, R., & Aarsland, T. (1999). Depressed mood and subjective health symptoms as predictors of mortality in patients with congestive heart failure: A two-years follow-up study. *International Journal of Psychiatry in Medicine*, 29(3), 311-326.
- 村山航(2011).欠損データ分析(missing data analysis)-完全情報最尤推定法と多重代入法.1-18.
- 森 啓至.(2014). 生理学から見た廃用症候群. 奈良勲ら(編), 理学療法から診る廃用症候群 (pp.12-14).文光堂.
- 長山 雅俊. (2012). 【ICU 患者を動かそう!】 心臓リハビリテーションの現況. *Icu と ccu*, 36(6), 415-422.

Nieminen, M. S., Bohm, M., Cowie, M. R., Drexler, H., Filippatos, G. S., Jondeau, G., . . . ESC Committee for Practice Guideline (CPG). (2005). Executive summary of the guidelines on the diagnosis and treatment of acute heart failure: The task force on acute heart failure of the European Society of Cardiology. *European Heart Journal*, 26(4), 384-416. doi:ehi044 [pii]

Nieminen, M. S., Brutsaert, D., Dickstein, K., Drexler, H., Follath, F., Harjola, V. P., . . . Heart Failure Association, European Society of Cardiology. (2006). EuroHeart failure survey II (EHFS II): A survey on hospitalized acute heart failure patients: Description of population. *European Heart Journal*, 27(22), 2725-2736. doi:ehl193 [pii]

日本循環器学会(2013).急性心不全治療ガイドライン(2011 年改訂版).

[http://www.j-](http://www.j-circ.or.jp/guideline/pdf/JCS2011_izumi_h.pdf#search='%E6%80%A5%E6%80%A7%E5%BF%83%E4%B8%8D%E5%85%A8%E3%82%AC%E3%82%A4%E3%83%89%E3%83%A9%E3%82%A4%E3%83%B32011')

[circ.or.jp/guideline/pdf/JCS2011_izumi_h.pdf#search='%E6%80%A5%E6%80%A7%E5%BF%83%E4%B8%8D%E5%85%A8%E3%82%AC%E3%82%A4%E3%83%89%E3%83%A9%E3%82%A4%E3%83%B32011'](http://www.j-circ.or.jp/guideline/pdf/JCS2011_izumi_h.pdf#search='%E6%80%A5%E6%80%A7%E5%BF%83%E4%B8%8D%E5%85%A8%E3%82%AC%E3%82%A4%E3%83%89%E3%83%A9%E3%82%A4%E3%83%B32011') [2015-7-2]

O'Connor, C. M., Whellan, D. J., Lee, K. L., Keteyian, S. J., Cooper, L. S., Ellis, S. J., . . . HF-ACTION Investigators. (2009). Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA : The Journal of the American Medical Association*, 301(14), 1439-1450. doi:10.1001/jama.2009.454; 10.1001/jama.2009.454

大西勝也.(2014).心不全患者の心行動態.新しい心行動態入門(第 1 版)(pp100).メディカ版.

Ouslander, J. G., Diaz, S., Hain, D., & Tappen, R. (2011). Frequency and diagnoses associated with 7- and 30-day readmission of skilled nursing facility patients to a nonteaching community hospital. *Journal of the American Medical Directors Association*, 12(3), 195-203. doi:10.1016/j.jamda.2010.02.015 [doi]

- Paffenbarger, R. S., Jr, Hyde, R. T., Wing, A. L., Lee, I. M., Jung, D. L., & Kampert, J. B. (1993). The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *The New England Journal of Medicine*, 328(8), 538-545. doi:10.1056/NEJM199302253280804 [doi]
- Peterson, M. J., Giuliani, C., Morey, M. C., Pieper, C. F., Evenson, K. R., Mercer, V., . . . Health, Aging and Body Composition Study Research Group. (2009). Physical activity as a preventative factor for frailty: The health, aging, and body composition study. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 64(1), 61-68. doi:10.1093/gerona/gln001 [doi]
- Piepoli, M. F., Davos, C., Francis, D. P., Coats, A. J., & ExTraMATCH Collaborative. (2004). Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ (Clinical Research Ed.)*, 328(7433), 189. doi:10.1136/bmj.37938.645220.EE [doi]
- Pina, I. L., Apstein, C. S., Balady, G. J., Belardinelli, R., Chaitman, B. R., Duscha, B. D., . . . American Heart Association Committee on exercise, rehabilitation, and prevention. (2003). Exercise and heart failure: A statement from the american heart association committee on exercise, rehabilitation, and prevention. *Circulation*, 107(8), 1210-1225.
- Rapaport, E., Wong, M., Escobar, E. E., & Martinez, G. (1966). The effect of upright posture on right ventricular volumes in patients with and without heart failure. *American Heart Journal*, 71(2), 146-152. doi:0002-8703(66)90177-3 [pii]
- Reeder, K. M., Ercole, P. M., Peek, G. M., & Smith, C. E. (2015). Symptom perceptions and self-care behaviors in patients who self-manage heart failure. *The Journal of Cardiovascular Nursing*, 30(1), E1-7. doi:10.1097/JCN.0000000000000117 [doi]
- Rutledge, T., Reis, V. A., Linke, S. E., Greenberg, B. H., & Mills, P. J. (2006). Depression in heart failure a meta-analytic review of prevalence, intervention

- effects, and associations with clinical outcomes. *Journal of the American College of Cardiology*, 48(8), 1527-1537. doi:S0735-1097(06)01905-X [pii]
- Sahebi, A., Mohammad-Aliha, J., Ansari-Ramandi, M., & Naderi, N. (2015). Investigation the relationship between self-care and readmission in patients with chronic heart failure. *Research in Cardiovascular Medicine*, 4(1), e25472. doi:10.5812/cardiovascmed.25472 [doi]
- Saitoh, M., Rodrigues Dos Santos, M., & von Haehling, S. (2016). Muscle wasting in heart failure : The role of nutrition. *Wiener Klinische Wochenschrift*, 128(Suppl 7), 455-465. doi:10.1007/s00508-016-1100-z [doi]
- 齊藤 正和, 堀 健太郎, 岡村 大介, 坂本 純子, 鈴木 秀俊, 中澤 麻里子, . . . 小澤 哲也. (2015). 多施設共同研究による高齢心不全患者の退院時日常生活動作(ADL)に関連する因子の検討. *理学療法学*, 42(2), 81-89.
- Saltin, B., Blomqvist, G., Mitchell, J. H., Johnson, R. L., Jr, Wildenthal, K., & Chapman, C. B. (1968). Response to exercise after bed rest and after training. *Circulation*, 38(5 Suppl), VII1-78.
- Sato, N., Kajimoto, K., Keida, T., Mizuno, M., Minami, Y., Yumino, D., . . . TEND Investigators. (2013). Clinical features and outcome in hospitalized heart failure in japan (from the ATTEND registry). *Circulation Journal : Official Journal of the Japanese Circulation Society*, 77(4), 944-951. doi:DN/JST.JSTAGE/circj/CJ-13-0187 [pii]
- Scrutinio, D., Passantino, A., Catanzaro, R., Farinola, G., Lagioia, R., Mastropasqua, F., . . . Santoro, D. (2012). Inpatient cardiac rehabilitation soon after hospitalization for acute decompensated heart failure: A propensity score study. *Journal of Cardiopulmonary Rehabilitation and Prevention*, 32(2), 71-77. doi:10.1097/HCR.0b013e31823be124; 10.1097/HCR.0b013e31823be124

- 志賀 剛. (2004). 【集中治療における心房細動の臨床】 心不全. *Icu と ccu*, 28(8), 559-566.
- 島田 裕之, & 吉田 大輔. (2011). 【高齢者の虚弱 評価と対策】 虚弱とサルコペニア(概念の相違). *Geriatric Medicine*, 49(3), 291-295.
- 清水 優子, 河野 裕治, 小林 聖典, & 山田 純生. (2011). 臓器合併症に対する心臓リハビリ療法士としてどう立ち向かうか 心不全に合併する抑うつ. *心臓リハビリテーション*, 16(1), 63-67.
- 関根 由紀, & 小松 浩子. (2010). 人工呼吸器離脱過程における心不全患者の取り組みの構造化. *日本クリティカルケア看護学会誌*, 6(1), 16-25.
- 関根 由紀. (2015). 急性心不全患者の人工呼吸器離脱過程における看護実践の構造化. *日本循環器看護学会誌*, 11(1), 55-63.
- 園田 茂, 椿原 彰夫, 田尻 寿子, & 他. (1992). FIM を用いた脳血管障害患者の機能評価 barthel index (BI)との比較およびコミュニケーションと社会的認知能力の関与. *リハビリテーション医学*, 29(3), 217-222.
- 園田 茂, & 千野 直一. (1993). 能力低下の評価法について. *リハビリテーション医学*, 30(7), 491-500.
- Stevens, R. D., Dowdy, D. W., Michaels, R. K., Mendez-Tellez, P. A., Pronovost, P. J., & Needham, D. M. (2007). Neuromuscular dysfunction acquired in critical illness: A systematic review. *Intensive Care Medicine*, 33(11), 1876-1891.
doi:10.1007/s00134-007-0772-2 [doi]
- 鈴木 洋児, & 郡司 篤晃. (1991). Deconditioning と運動耐容能. *循環器科*, 30(6), 550-559.
- S.J.ギャリクソン編(2005).第1章不動.リハビリテーション実践ハンドブック(pp11).石田 暉、豊倉穰、花山耕三監訳.東京. シュプリンガー・フェアラーク東京.

- 田畑 稔, 新井 保久, 瀬城 亜也子, 家護谷 知子, 竹中 杏奈, 中川 晋, ... 廣谷 隆.
(2004). 急性心不全に対する急性期心臓リハビリテーション成績 心疾患リハビリテーション料適応疾患との比較. *心臓リハビリテーション*, 9(1), 63-66.
- 田畑 稔, 中川 晋, 宇井 進, 三田村 秀雄, & 山田 純生. (2007). 慢性心不全リハの阻害要因を考える 入院期心不全リハビリテーションにおける阻害要因の検討. *心臓リハビリテーション*, 12(1), 40-43.
- 武市 尚也, 井澤 和大, 渡辺 敏, 平木 幸治, 森尾 裕志, 長田 尚彦, & 大宮 一人. (2009). 回復期冠動脈疾患患者における身体活動量と下肢筋力との関連について. *理学療法*, 36(3), 109-113.
- 田中 由貴. (2014). せん妄に陥った男性高齢患者にドールセラピーを試みて日中の混乱が改善した 1 事例. *奈良県立三室病院看護学雑誌*, 30, 60-62.
- 谷口良司.(2014).第 4 章ガイドラインからみた心臓リハビリテーション.佐藤幸人(編),臨床心不全のいちばん大事なところ(pp252).メディカ出版.
- Tavazzi, L., Maggioni, A. P., Lucci, D., Cacciatore, G., Ansalone, G., Oliva, F., ...
Italian survey on Acute Heart Failure Investigators. (2006). Nationwide survey on acute heart failure in cardiology ward services in Italy. *European Heart Journal*, 27(10), 1207-1215. doi:ehi845 [pii]
- 筒井裕之, 絹川真太郎.(2010).第 3 章心不全.川名正敏,北風政史,小室一成ら(編),循環器病学(pp338).西村書店.
- 友田春夫.(2010).第 17 章その他の薬物療法.心不全—循環生理からみた診断と治療—(pp178).医学書院.
- Vaccarino, V., Kasl, S. V., Abramson, J., & Krumholz, H. M. (2001). Depressive symptoms and risk of functional decline and death in patients with heart failure. *Journal of the American College of Cardiology*, 38(1), 199-205. doi:S0735-1097(01)01334-1 [pii]

Wilson, J. R., Martin, J. L., & Ferraro, N. (1984). Impaired skeletal muscle nutritive flow during exercise in patients with congestive heart failure: Role of cardiac pump dysfunction as determined by the effect of dobutamine. *The American Journal of Cardiology*, 53(9), 1308-1315. doi:0002-9149(84)90085-7 [pii]

安村良男.(2010).第3章心不全.急性心不全の病態生理. 川名正敏,北風政史,小室一成ら編集, 循環器病学(pp308).医学書院.

山田 純生, 長谷部 武久, 井澤 英夫, 大宮 一人, 大野 三良, 加藤 林也, . . . 横田 充弘. (2007). 慢性心不全リハの阻害要因を考える 慢性心不全の不安・抑うつへの運動介入をどうとらえるか. *心臓リハビリテーション*, 12(1), 47-51.

Zannad, F., Mebazaa, A., Juilliere, Y., Cohen-Solal, A., Guize, L., Alla, F., . . . EFICA Investigators. (2006). Clinical profile, contemporary management and one-year mortality in patients with severe acute heart failure syndromes: The EFICA study. *European Journal of Heart Failure*, 8(7), 697-705. doi:S1388-9842(06)00002-X [pii]

Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, 67(6), 361-370.