

## Reference

- Akatsuka, J. (1992). Syounihinketsuno rinsyou [Anemia in pediatrics]. Tokyo;Kaneharasyuppan.pp.49-50.
- Al-Tawil, M., Abdel-Aal, R., & Kaddah, A. (2012). A randomized controlled trial on delayed cord clamping and iron status at 3–5 months in term neonates held at the level of maternal pelvis. *Journal of Neonatal-Perinatal Medicine*, 5(4), 319-326.
- Andersson, O., Hellstrom-Westas, L., Andersson, D., & Domellof, M. (2011). Effect of delayed versus early umbilical cord clamping on neonatal outcomes and iron status at 4 months: A randomised controlled trial. *BMJ (Clinical Research Ed.)*, 343, d7157. doi:10.1136/bmj.d7157 [doi]
- Andersson, O., Domellof, M., Andersson, D., & Hellstrom-Westas, L. (2013). Effects of delayed cord clamping on neurodevelopment and infection at four months of age: A randomised trial. *Acta Paediatrica* (Oslo, Norway: 1992), 102(5), 525-531. doi:10.1111/apa.12168 [doi]
- Avery, G. B. (1990). *Neonatology pathophysiology and management of the newborn*. lippincott williams and wilkins, philadelphia, USA. Philadelphia, USA: Lippincott Williams and Wilkins.
- Baker, R. D., Greer, F. R., & Committee on Nutrition American Academy of Pediatrics. (2010). Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0-3 years of age). *Pediatrics*, 126(5), 1040-1050. doi:10.1542/peds.2010-2576 [doi]
- Becher, J. C., Bhushan, S. S., & Lyon, A. J. (2012). Unexpected collapse in apparently healthy newborns--a prospective national study of a missing cohort of neonatal deaths and near-

death events. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 97(1), F30-4.  
doi:10.1136/adc.2010.208736 [doi]

Begley, C. M. (1990). A comparison of 'active' and 'physiological' management of the third stage of labour. *Midwifery*, 6(1), 3-17.

Begley, C. M., Gyte, G. M., Devane, D., McGuire, W., & Weeks, A. (2015). Active versus expectant management for women in the third stage of labour. *The Cochrane Database of Systematic Reviews*, (3):CD007412.doi(3), CD007412.  
doi:10.1002/14651858.CD007412.pub4 [doi]

Bhutani, V. K., Gourley, G. R., Adler, S., Kreamer, B., Dalin, C., & Johnson, L. H. (2000). Noninvasive measurement of total serum bilirubin in a multiracial predischarge newborn population to assess the risk of severe hyperbilirubinemia. *Pediatrics*, 106(2), E17.

Boere, I., Smit, M., Roest, A. A., Lopriore, E., van Lith, J. M., & te Pas, A. B. (2015a). Current practice of cord clamping in the netherlands: A questionnaire study. *Neonatology*, 107(1), 50-55. doi:10.1159/000365836 [doi]

Boere, I., Roest, A., Wallace, E., Ten Harkel, A., Haak, M., Morley, C., . . . te Pas, A. (2015b). Umbilical blood flow patterns directly after birth before delayed cord clamping. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 100(2), F121-5.  
doi:10.1136/archdischild-2014-307144 [doi]

Ceriani, J. M., Carroli, G., Pellegrini, L., Otano, L., Ferreira, M., Ricci, C., . . . Lardizabal, J. (2006). The effect of timing of cord clamping on neonatal venous hematocrit values and clinical outcome at term: A randomized, controlled trial. *Pediatrics*, 117(4), e779-86.  
doi:peds.2005-1156 [pii]

Chaparro, C. M., Neufeld, L. M., Tena Alavez, G., Eguia-Liz Cedillo, R., & Dewey, K. G. (2006). Effect of timing of umbilical cord clamping on iron status in mexican infants: A randomised controlled trial. *Lancet*, 367(9527), 1997-2004. doi:S0140-6736(06)68889-2 [pii]

Chien, P. C., Yang, C. C., Gau, M. L., Liu, C. Y., & Lee, T. Y. (2015). The impact of late umbilical cord clamping on neonatal jaundice and postpartum hemorrhage: A randomized controlled trial. *Hu Li Za Zhi the Journal of Nursing*, 62(4), 41-53. doi:10.6224/JN62.4.41 [doi]

Domellof, M., Braegger, C., Campoy, C., Colomb, V., Decsi, T., Fewtrell, M., . . . ESPGHAN Committee on Nutrition. (2014). Iron requirements of infants and toddlers. *Journal of Pediatric Gastroenterology and Nutrition*, 58(1), 119-129. doi:10.1097/MPG.0000000000000206 [doi]

Eggert, L. D., Wiedmeier, S. E., Wilson, J., & Christensen, R. D. (2006). The effect of instituting a prehospital-discharge newborn bilirubin screening program in an 18-hospital health system. *Pediatrics*, 117(5), e855-62. doi:117/5/e855 [pii]

Embrey,M. (1961). Simultaneous intramuscular injection of oxytocin and ergometrine: A tocographic study. *British Medical Journal*, 1737-8.

Farrar, D., Tuffnell, D., Airey, R., & Duley, L. (2010). Care during the third stage of labour: A postal survey of UK midwives and obstetricians. *BMC Pregnancy and Childbirth*, 10, 23-2393-10-23. doi:10.1186/1471-2393-10-23 [doi]

Furukawa, R. (1982). Hinketsu no seiin to shindan [the origin and diagnosis of anemia]. *Syounika MOOK*, (25), 26-34.

Goldsmith, J. Hidaka, R (trans.). Shizen syussan no chie [Child birth wisdom form world oldest societies].Tokyo:Nihonkyobunsya.

Grajeda, R., Perez-Escamilla, R., & Dewey, K. G. (1997). Delayed clamping of the umbilical cord improves hematologic status of guatemalan infants at 2 mo of age. *The American Journal of Clinical Nutrition*, 65(2), 425-431.

Gupta, R., & Ramji, S. (2002). Effect of delayed cord clamping on iron stores in infants born to anemic mothers: A randomized controlled trial. *Indian Pediatrics*, 39(2), 130-135.

- Igarashi, T. (2005). *Syounikagaku*[pediatrics] (9th edition ed.). Tokyo: Bunkodo. pp730-731.
- Imura,S.(1985). Shinseijioudanno chiryou kousenryouhou tekioukijyun to fukusayouno boushi [treatment of neonatal jaundice standard of phototherapy]. *Nihon Rinsyou*, 43(8), 1741-1748.
- International Confederation of Midwives (ICM), & International Federation of Gynaecology and Obstetrics (FIGO). (2006). Prevention and treatment of post-partum haemorrhage: New advances for low resource settings: Joint statement international confederation of midwives (ICM) international federation of gynaecology and obstetrics (FIGO). Retrieved from <http://www.who.int/pmnch/events/2006/figo2006statementeng.pdf>
- Isomura, H., Takimoto, H., Miura, F., Kitazawa, S., Takeuchi, T., Itabashi, K., & Kato, N. (2011). Type of milk feeding affects hematological parameters and serum lipid profile in Japanese infants. *Pediatrics International : Official Journal of the Japan Pediatric Society*, 53(6), 807-813. doi:10.1111/j.1442-200X.2011.03360.x [doi]
- Itou, S., Matsunaga, T., Hara, T., Hayakawa, J., & Isobe, K. (1978). Shinseijino kessyoubirirubinchito saitaikessatujikitono kankeinitsuite [relationships between serum bilirubin level and timing of cord clamping]. *Syuusanki Igaku*, 8(8), 974-978.
- Jahazi, A., Kordi, M., Mirbehbahani, N. B., & Mazloom, S. R. (2008). The effect of early and late umbilical cord clamping on neonatal hematocrit. *Journal of Perinatology*, 28(8), 523-525. doi:10.1038/jp.2008.55 [doi]
- Joo, E. Y., Kim, K. Y., Kim, D. H., Lee, J. E., & Kim, S. K. (2016). Iron deficiency anemia in infants and toddlers. *Blood Research*, 51(4), 268-273. doi:10.5045/br.2016.51.4.268 [doi]
- Kangaroo-care working group. (2010). Evidence based kangaroo-care (SSC) guidelines. Retrieved from [http://minds4.jcqhc.or.jp/minds/kc/fukyu/1\\_kc.pdf](http://minds4.jcqhc.or.jp/minds/kc/fukyu/1_kc.pdf)
- Kataoka, Y., Nakayama, K., Yaju, Y., Eto, H., & Horiuchi, S. (2015). Comparison of policies for the management of care for women and newborns during the third stage of labor among

japanese hospitals, clinics, and midwifery birth centers. *International Journal of Childbirth*, 5(4), 200-209(10).

Kimura, M. (2014). Bonyuutojinkoueiyou nyuuyoujino tetsuketsubouseihinketsuto bonyueiyou [breast milk and iron defecency anemia in infants]. *Syounikarinsyou*, 67(12), 2401-2406.

Kliegman, R., Stanton, B. F., St Geme, J. W., III, Schor, N. F., Behrman, R. E. (2011). *Nelson textbook of pediatrics* (19th ed ed.). Philadelphia, PA: Elsevier. pp.603-608e,612-615e,1655-1658e,1682-1683e.

Kobe University Medical school pediatric department. (1993). *Mijyukuji shinseijinokanri* [management of premature neonates]. (p.233).

Kubo, T.(2012).Sankanyuuincyuno shinseijinoshinryoutokea bunben cyokugono souki boshisessyokunitsuite zenkoku anketocousawofumaete [national survey for early skin to skin contact ]. *Nihon Mijyukuji Shinseijigakkai Zasshi*, 24(3), 467.

Kuboi, T., Kusaka, T., Kawada, K., Koyano, K., Nakamura, S., Okubo, K., . . . Itoh, S. (2013). Hour-specific nomogram for transcutaneous bilirubin in Japanese neonates. *Pediatrics International*, 55(5), 608-611. doi:10.1111/ped.12149 [doi]

Li, N., Yang, L. C., Wu, Q., Han, C. C., Wang, L., Rong, L., . . . Zhang, W. (2012). The effects of iron stores and growth of delayed umbilical cord clamp timing on term breastfed infants at 4 months. *Zhonghua Yu Fang Yi Xue Za Zhi [Chinese Journal of Preventive Medicine]*, 46(4), 303-306.

Lundberg, C., Oian, P., & Klingenberg, C. (2013). Umbilical cord clamping at birth--practice in Norwegian maternity wards. *Tidsskrift for Den Norske Laegeforening : Tidsskrift for Praktisk Medicin, Ny Raekke*, 133(22), 2369-2373. doi:10.4045/tidsskr.12.0707 [doi]

Maeda, M. (2015). Ketsuekishikkan no syohou tetsuketsubousei hinketsu [perscription for blood disorder -iron defecency anemia ]. *Syounika Rinsyou*, 68(4), 780-784.

Marumoto, K., Nakao, H., Azuma, A. (1994). Saitaikessatsujikiga soukishinseijikiniokeru kesseibirurubinchi oyobi honyuuryokuni oyobosueikyou. *Perinatal Care*, 13(4), 359-363.

Maruo, Y. (2012). Shinseijikino oudanto bilirubin UDP-grucronsantenikouso (UGT1A1 idenshiheni[neonatal jaundice and bilirubin UDP - glucuronic acid transferase gene mutation]. *Syounikashinryou*, 75(9), 1572-1577.

McDonald, S., Middleton, P., Dowswell, T., & Morris, S. (2013). Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes. *The Cochrane Database of Systematic Reviews*, 7, CD004074. doi:10.1002/14651858.CD004074.pub3 [doi]

Mercer, J., S., & Erickson-Owens, D. (2012). Rethinking placental transfusion and cord clamping issues. *Journal of Perinatal & Neonatal Nursing*, 26(3), 202-217.

Moghaddam Tabrizi, F., & Barjasteh, S. (2015). Maternal hemoglobin levels during pregnancy and their association with birth weight of neonates. *Iranian Journal of Pediatric Hematology and Oncology*, 5(4), 211-217.

Ministry of Health, Labour and Welfare. (2010). Heisei22nendo nyuuyouji shintaihatiku cyousa houkokusyo [2010 report for infant and toddler physical growth]. Retrieved from <http://www.mhlw.go.jp/stf/houdou/2r9852000001t3so-att/2r9852000001t7dg.pdf>

Moore, E. R., Bergman, N., Anderson, G. C., & Medley, N. (2016). Early skin-to-skin contact for mothers and their healthy newborn infants. *The Cochrane Database of Systematic Reviews*, 11, CD003519. doi:10.1002/14651858.CD003519.pub4 [doi]

Morimoto, Y. (2012). Shinseijioudanchiryouniokeru hikarinoyakuwari [role of the light in the jaundice of the newborn treatment]. *Lightedge*, 38, 17-25.

Mothers' and Children's Health & Welfare Association. (2016). Boshihoken no omonaru toukei [maternal and child health statistics of Japan]. Tokyo: Mothers' & Children's health & welfare association; Tokyo: Mothers' & Children's Health & Welfare Association.

National Institute for Health and Care Excellence. (2014). Intrapartum care: Care of healthy women and their babies during childbirth. NICE guidelines [CG190] 1.14 third stage of labour. Retrieved from <https://www.nice.org.uk/guidance/cg190/chapter/1-recommendations#third-stage-of-labour>

Newman, T. B., Easterling, M. J., Goldman, E. S., & Stevenson, D. K. (1990). Laboratory evaluation of jaundice in newborns. frequency, cost, and yield. *American Journal of Diseases of Children (1960)*, 144(3), 364-368.

Nishida, H. (2012) *Shinseijigaku nyuumon [introduction to neonatology]* (4th edition ed.). Tokyo: Igakusyoin. pp66-67, 209-305.

Nishizawa, Y., Fujimatsu, M., Fujita, T., & Matsuoka, K. (2002). Shinseijioudanto oxytocin no kankei (daiippou) shinseijikoubilirubinkessyouno sankatekirisukunitsuite. [relationships between neonatal jaundice and oxytocin (no.1) Risk factor of neonatal bilirubinemia]. *Nihon Miyukuji Shinseijigakkaishi*, 14(3), 456.

Poets, A., Steinfeldt, R., & Poets, C. F. (2011). Sudden deaths and severe apparent life-threatening events in term infants within 24 hours of birth. *Pediatrics*, 127(4), e869-73. doi:10.1542/peds.2010-2189 [doi]

Preer, G. L., & Philipp, B. L. (2011). Understanding and managing breast milk jaundice. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 96(6), F461-6. doi:10.1136/adc.2010.184416 [doi]

Raicevic, S., Eventov-Friedman, S., Bolevich, S., Selakovic, D., Joksimovic, J., Djuric, J., . . . Jakovljevic, V. (2014). Correlation between oxidative stress and G6PD activity in neonatal jaundice. *Molecular and Cellular Biochemistry*, 395(1-2), 273-279. doi:10.1007/s11010-014-2136-x [doi]

Renner, J., Broch, O., Scheewe, J., Gruenewald, M., & Bein, B. (2010). Noninvasive estimation of hemoglobin by pulse-CO-oximetry in infants undergoing major surgery [Abstract]. *2010 Annual Meeting of the American Society Anesthesiologists*, Abs 1668.

Ryu, T., Maruo, Y., & Kusuda, S. (2007). Syuusankitoudan up to date. sankaniokeru shinseijikanri [management of neonates in obstetrics]. *Syusankiigaku*, 37(10), 1275-1280.

Sakaguchi, K., Tokutake, S., Haga, A., & Kondo, R. (2013). Seikisanjini taisuru soukiboshisessyokuno koukatoanzensei [effects and safety of early skin to skin contact for term neonates]. *Sinsyu Igaku Zasshi*, 61(5), 263-272.

Samiee-Zafarghandy, S., Feberova, J., Williams, K., Yasseen, A. S., Perkins, S. L., & Lemire, B. (2014). Influence of skin color on diagnostic accuracy of the jaundice meter JM 103 in newborns. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 99(6), F480-4. doi:10.1136/archdischild-2013-305699 [doi]

Sanpavat, S., & Nuchprayoon, I. (2004). Noninvasive transcutaneous bilirubin as a screening test to identify the need for serum bilirubin assessment. *Journal of the Medical Association of Thailand*, 87(10), 1193-1198.

Setia, S., Villaveces, A., Dhillon, P., & Mueller, B. A. (2002). Neonatal jaundice in asian, white, and mixed-race infants. *Archives of Pediatrics & Adolescent Medicine*, 156(3), 276-279. doi:poa10200 [pii]

Shimada, K., & Shirakawa, K. (2011). Syuusankino ketsuekiwomeguru wadai. sekkekkyuuno ijyou shinseijino taketsusyou.[abnormality of the red blood cell- polycythemia]. *Syusanki Igaku*, 41(9), 1125-1127.

Shin, S., Ogawa, H., Araki, T., et al.(1990). Shinseiji kouhematoclitokessyouto saitaikessatsujikin okankei anteidouigenso cr-50 womochiite [assessing relationship between high hematoclit and timing of cord clamping with stable isotope cr-50]. *Biomedical Research on Trace Elements*, 1(2), 205-206.

Shinohara, E., Kataoka, Y., & Matsui, A. (2016). Prevalence and Risk Factors for Hyperbilirubinemia among Newborns from a Low-Risk Birth Setting in Japan .Manuscript submitted for publication.

Suzuki, R., Nomura, Y., Ujiie, J., & Ooto, S. (2007). Syusankitooudan up to date ketsuekigata futekigounkanri [management and treatment of blood type imcompatibility]. *Syuusanki Igaku*, 37(10), 1235-1240.

Takahashi, D., & Shirahata S. (2007). Syusankitooudan up to date oudan to yakubutsu [jaundice and medicine] . *Syuusanki Igaku*, 37(10), 1329-1334.

Takayama, T., Endo, Y., Tamashiro, Y., Tsujino, K., Tsugiko, G., Yamada, S., . . . Kawamitsu, K. (2015). Saitaikessatsujikiga seijyukujino seiritekioudanto nyuujisoukino keccyuu hemogurobinchini oyobosu eikyounitsuute[effects of the timing of umbilical cord clamping on physiological jaundice and hemoglobin levels in full term infants ]. *Boseieise*, 56(1), 77-86.

Tamura, M. (2011). Nihonban kinkyuusosei guideline 2010 ni motozuku Shinseiji soseihou text [text book of neonatal resuscitation based on Japanese resuscitation guideline 2010] (version2 ed.). Tokyo: Medical view.

Tamura, M., & Hosono, S.(2012). Consensus2010 ni motozuku atarashii nihonban shinsejisoseihou guidline nokakuritu fukyuuto sonokoukanohoukanikansurukenkyuu.[Evaluation for the effects of establish Japanese neonatal resustitation guideline based on consensus 2010]. Retrieved from <http://www.aiiku.or.jp/~doc/houkoku/h24/19006A130.pdf>

Tanaka, T., Nakamune, T., Tamanaha, E. & Yamagata, R. (2012). Nyuuyouji kenshin jyuudan data no riyouhouhou ni kansrukennkyu [research for using data of infants' check ups].

Retrieved from <http://www.aiiku.or.jp/~doc/houkoku/h24/19019A070.pdf>

Ukita, M. (1994). Ketsuekigata futekigouninshin to taiji shinseiji youketsusei shikkann (dai4hen) kouDkoutaiigaino sekkekkyuukoutainiyoru taiji shinseiji youketsusei shikkan [the blood group incompatibility pregnancy and fetus, hemolytic disease of the newborn]. *Nihon sanfujinka shinseiji Ketsuekigakkaishi*, 3(3), 55-80.

Vain, N. E., Satragno, D. S., Gorenstein, A. N., Gordillo, J. E., Berazategui, J. P., Alda, M. G., & Prudent, L. M. (2014). Effect of gravity on volume of placental transfusion: A multicentre, randomised, non-inferiority trial. *Lancet*, 384(9939), 235-240. doi:10.1016/S0140-6736(14)60197-5 [doi]

van Rheenen, P., de Moor, L., Eschbach, S., de Grooth, H., & Brabin, B. (2007). Delayed cord clamping and haemoglobin levels in infancy: A randomised controlled trial in term babies. *Tropical Medicine & International Health : TM & IH*, 12(5), 603-616. doi:TMI1835 [pii]

Watanabe, K., & Ouchi, Y. (1996). Koubilirubin kessyou no yobouni tsuite saitaikessatsu houhou no kentou [prevention for hyperbilirubinemia - considering timing of cord clamping]. *Ibarakiken Boseieisei Gakkaishi*, (16), 38-41.

World Health Organization. (2008). World prevalence of anemia 1993-2005 . Retrieved from <http://www.who.int/vmnis/publications/en/>

World Health Organization. (2012). WHO recommendations for the prevention and treatment of postpartum haemorrhage. Retrieved from [http://www.who.int/reproductivehealth/publications/maternal\\_perinatal\\_health/9789241548502/en/](http://www.who.int/reproductivehealth/publications/maternal_perinatal_health/9789241548502/en/)

World Health Organization. (2014). Guideline: Delayed umbilical cord clamping for improved maternal and infant health and nutrition outcomes. Retrieved from [http://apps.who.int/iris/bitstream/10665/148793/1/9789241508209\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/148793/1/9789241508209_eng.pdf)

Yamada, K. (2007). Syusankito oudan up to date koukan uketsu [exchange blood transfusion]. *Syusankigaku*, 37(10), 1263-1267.

Yamauchi, Y. (1994). Seiritekioudan to byoutekioudan. [Physiological jaundice and pathologicall jaundice]. Neonatal care 7, Autumn edition, 85-96.

Yamauchi, Y. (2007). Syusankito oudan up to date keihi birirubinsokutei, unboundbirirubine sokutei [transcuneous bilirubin and unbound bilirubin measurement]. *Syuusanki Igaku*, 37(10), 1247-1252.

Yamamura, Y., & Yoshiri, K. (Eds.). (1992). *Hinketsu, taketsusyuu [anemia and polycythemia]*. Tokyo: Nakayamasyoten. pp. 86-87.

Yang, Z., Lonnerdal, B., Adu-Afarwuah, S., Brown, K. H., Chaparro, C. M., Cohen, R. J., . . . Dewey, K. G. (2009). Prevalence and predictors of iron deficiency in fully breastfed infants at 6 mo of age: Comparison of data from 6 studies. *The American Journal of Clinical Nutrition*, 89(5), 1433-1440. doi:10.3945/ajcn.2008.26964 [doi]

Yao, A. C., Hirvensalo, M., & Lind, J. (1968). Placental transfusion-rate and uterine contraction. *Lancet*, 1(7539), 380-383.

Yao, A. C., & Lind, J. (1969). Effect of gravity on placental transfusion. *Lancet*, 2(7619), 505-508. doi:S0140-6736(69)90213-X [pii]

Yao, A. C., & Lind, J. (1974). Placental transfusion. *American Journal of Diseases of Children*, 127(1), 128-141.

Yigit, M. B., Kowalski, W. J., Hutchon, D. J., & Pekkan, K. (2015). Transition from fetal to neonatal circulation: Modeling the effect of umbilical cord clamping. *Journal of Biomechanics*, 48(9), 1662-1670. doi:10.1016/j.jbiomech.2015.02.040 [doi]