

References

- AbouZahr, C. (2003). Global burden of maternal death and disability. *British Medical Bulletin*, 67, 1-11.
- American College of Obstetricians and Gynecologists. (ACOG). (2006). ACOG Practice Bulletin: Clinical Management Guidelines for Obstetrician-Gynecologists Number 76, October 2006: postpartum hemorrhage. *Obstetrics & Gynecology*, 108(4), 1039-47.
- Baldisseri, R. M. (2011). Postpartum Hemorrhage. In [Jean-Louis Vincent, Edward Abraham, Frederick, A. Moore, Patrick, M. Kochanek, & Mitchell, P. Fink](#) (Eds.), *Textbook of Critical Care*. (sixth ed., pp.1192-1197). Saunders: Elsevier Inc.
- Begley, C. M., Gyte, G. M. L., Devane, D., McGuire, W., & Weeks, A. (2015). Active versus expectant management for women in the third stage of labour. *The Cochrane Database of Systematic Reviews*, 3, CD007412. doi: 10.1002/14651858.CD007412.pub4

Burbank, F. (2012). Hemodynamic changes in the uterus and its blood vessels in pregnancy. In: Arulkumaran, S., Karoshi, M., Keith, L. G., Lalonde. AB., B-Lynch, C. (Eds.), *A Comprehensive Textbook of Postpartum Haemorrhage*. (2nd ed., pp.177-184) London: Sapiens.

Carroli G, Cuesta C, Abalos E, & Gulmezoglu AM. (2008). Epidemiology of postpartum haemorrhage: a systematic review. *Best Practice & Research Clinical Obstetrics and Gynaecology*, 22(6), 999-1012.

Chua, S., Arulkumaran, S., Roy, A. C., Ho, L. M., Pathiraja, R., & Ratnam, S. S. (1996). Correlation of total uterine activity to blood loss in the third stage of labour. *Gynecologic and Obstetric Investigation*, 42(3), 171-173.

Clinical Evidence. (2012). Study design search filters. Retrieved from
<http://clinicalevidence.bmjjournals.org/x/set/static/ebm/learn/665076.html>

Combs, C. A., Murphy, E. L., & Laros, R. K., Jr. (1991). Factors associated with postpartum hemorrhage with vaginal birth. *Obstetrics and Gynecology*, 77(1), 69-76.

Cunningham, F. G., Leveno K. J., Bloom S. L., Spong C. Y., Dashe J. S., Hoffman B.

L., ...Sheffield J. S. (Eds.). (2014a). Obstetrical Hemorrhage. *Williams*

obstetrics (24th ed. pp.780-828). New York: McGraw-Hill Medical.

Cunningham, F. G., Leveno K. J., Bloom S. L., Spong C. Y., Dashe J. S., Hoffman B.

L., ...Sheffield J. S. (Eds.). (2014b). Maternal Anatomy. *Williams obstetrics*

(24th ed. pp.16-35). New York: McGraw-Hill Medical.

Du, Y., Ye, M., & Zheng, F. (2014). Active management of the third stage of labor with

and without controlled cord traction: A systematic review and meta-analysis of

randomized controlled trials. *Acta Obstetricia Et Gynecologica Scandinavica*,

93(7), 626-633. doi:10.1111/aogs.12424

Fahy, K. M. (2009). Third stage of labour care for women at low risk of postpartum

haemorrhage. *Journal of Midwifery & Women's Health*, 54(5), 380-386. doi:

10.1016/j.jmwh.2008.12.016

Faraoni, D., Carlier, C., Samama, C. M., Levy, J. H., & Ducloy-Bouthors, A. (2014).

[Efficacy and safety of tranexamic acid administration for the prevention

- and/or the treatment of post-partum haemorrhage: A systematic review with meta-analysis]. *Annales Françaises D'Anesthésie Et De Rèanimation*, 33(11), 563-571. doi: 10.1016/j.annfar.2014.07.748
- Ferrer, P., Roberts, I., Sydenham, E., Blackhall, K., & Shakur, H. (2009). Anti-fibrinolytic agents in post partum haemorrhage: A systematic review. *BMC Pregnancy and Childbirth*, 9, 29-29. doi:10.1186/1471-2393-9-29
- Ford, J. B., Shand, A. W., & Roberts, C. L. (2013). Characteristics, causes and treatment of postpartum haemorrhage in first and second pregnancies. *The Australian and New Zealand Journal of Obstetrics & Gynaecology*, 53(1), 90-93.
- Fujita, M., Manabe, E., & Morooka, T. (1994). Sangono sikuuhukkoni kannsuru inshino kentou [Investigation of factors related to involution of the after childbirh]. *Bulletin of College of Medical Technology Kyoto Prefectural University of Medicine*, 4(1), 90-93.

- Fyfe, E. M., Thompson, J. M., Anderson, N. H., Groom, K. M., & McCowan, L. M. (2012). Maternal obesity and postpartum haemorrhage after vaginal and caesarean delivery among nulliparous women at term: A retrospective cohort study. *BMC Pregnancy and Childbirth*, 12, 112-2393-12-112.
doi:10.1186/1471-2393-12-112 [doi]
- Guyatt, GH., Oxman, AD., Kunz, R., Vist, GE., Falck-Ytter, Y., Schunemann, HJ., & GRADE Working Group. (2008). What is "quality of evidence" and why is it important to clinicians? *BMJ*. doi: 10.1136/bmj.39490.551019.BE.
- Harmer, H. (1957). Local applications of heat, cold, and chemicals for circulatory effects. In Henderson V (Eds.), *Textbook of the principles and practice of nursing* (fifth ed., pp. 633-654). New York: Macmillan.
- Hayashi, M., Inagaki, E., Morita, S., Mano, M., & Amano, A. (1995). Effects of cold pack on the uterine contraction and hemostasis just after labor. *Journal of Japan Academy of Midwifery*, 9(1), 38-41.

- Heesen, M., Böhmer, J., Klöhr, S., Rossaint, R., van, d. V., Dudenhausen, J. W., & Straube, S. (2014). Prophylactic tranexamic acid in parturients at low risk for post-partum haemorrhage: Systematic review and meta-analysis. *Acta Anaesthesiologica Scandinavica*, 58(9), 1075-1085. doi:10.1111/aas.12341
- Hendricks, C. H. (1968). Uterine contractility changes in the early puerperium. *Clinical Obstetrics and Gynecology*, 11(1), 125-144.
- Hendricks, C. H., Eskes, T. K., & Saameli, K. (1962). Uterine contractility at delivery and in the puerperium. *American Journal of Obstetrics and Gynecology*, 83, 890-906.
- Higgins, JPT. (2011). In Green, S. (Eds.), *Cochrane Handbook for Systematic Reviews of Interventions* (version 5.1.0 [updated March 2011]). The Cochrane Collaboration. Retrieved from www.cochrane-handbook.org.

- Hofmeyr, G. J., Abdel-Aleem, H., & Abdel-Aleem, M. (2013). Uterine massage for preventing postpartum haemorrhage. *The Cochrane Database of Systematic Reviews*, 7, CD006431. doi: 10.1002/14651858.CD006431.pub3
- Hofmeyr, G. J., Gülmезoglu, A. M., Novikova, N., & Lawrie, T. A. (2013). Postpartum misoprostol for preventing maternal mortality and morbidity. *The Cochrane Database of Systematic Reviews*, 7, CD008982. doi: 10.1002/14651858.CD008982.pub2
- Hofmeyr, G. J., Gülmезoglu, A. M., Novikova, N., Linder, V., Ferreira, S., & Piaggio, G. (2009). Misoprostol to prevent and treat postpartum haemorrhage: A systematic review and meta-analysis of maternal deaths and dose-related effects. *Bulletin of the World Health Organization*, 87(9), 666-677.
- Hofmeyr, G. J., Mshweshwe, N. T., & Gülmезoglu, A. M. (2015). Controlled cord traction for the third stage of labour. *The Cochrane Database of Systematic Reviews*, 1, CD008020. doi: 10.1002/14651858.CD008020.pub2

- Hondo, K., Ishizuka, Y., & Kikuchi, H. (1993). Bunbengono kahukubureianpoouno sikuusyuusyukunitaisuru koukanokensho [Verification of the effect of cooling the lower abdomen for uterine contraction after the delivery]. *24th Nihonkangogakkaishuroku boseikango*. 133-135.
- Jin, B., Du, Y., Zhang, F., Zhang, K., Wang, L., & Cui, L. (2016). Carbetocin for the prevention of postpartum hemorrhage: A systematic review and meta-analysis of randomized controlled trials. *The Journal of Maternal-Fetal & Neonatal Medicine*, 29(3), 400-407. doi: 10.3109/14767058.2014.1002394.
- Joy, S. D., Sanchez-Ramos, L., & Kaunitz, A. M. (2003). Misoprostol use during the third stage of labor. *International Journal of Gynaecology and Obstetrics*, 82(2), 143-152.
- Kassebaum, N. J., Bertozzi-Villa, A., Coggeshall, M. S., Shackelford, K. A., Steiner, C., Heuton, K. R., . . . Lozano, R. (2014). Global, regional, and national levels and causes of maternal mortality during 1990-2013: A systematic analysis for

the global burden of disease study 2013. *Lancet*, doi: S0140-6736(14)60696-6

[pii]

Kataoka, Y., Nakayama, K., Yaju, Y., Eto, H., Horiuchi, S. (2015). Comparison of policies for the management of care for women and newborns during the third stage of labor among Japanese hospitals, clinics and midwifry birth centers.

International Journal of Childbirth, 5(4), 200-209.

Khan RU, & El-Rafaey H. (2012). Pathophysiology of postpartum haemorrhage and third stage of labour. In: Arulkumaran S, Karoshi M, Keith LG, Lalonde AB, B-Lynch C. (Eds.), *A Comprehensive Textbook of Postpartum Haemorrhage*. (2nd ed., pp.94–100). London: Sapiens.

Langenbach, C. (2006). Misoprostol in preventing postpartum hemorrhage: A meta-analysis. *International Journal of Gynaecology and Obstetrics*, 92(1), 10-18.

Leduc, D., Senikas, V., Lalonde, A. B., Ballerman, C., Biringer, A., Delaney, M., . . .

Society of Obstetricians and Gynaecologists of Canada. (SOGC). (2009).

Active management of the third stage of labour: Prevention and treatment of

postpartum hemorrhage. *Journal of Obstetrics and Gynaecology Canada*:

JOGC, 31(10), 980-993.

Liabsuetrakul, T., Choobun, T., Peeyananjarassri, K., & Islam, Q. M. (2007).

Prophylactic use of ergot alkaloids in the third stage of labour. *The Cochrane*

Database of Systematic Reviews, 2, CD005456. doi:

10.1002/14651858.CD005456.pub 2.

Magann, E. F., Lutgendorf, M. A., Keiser, S. D., Porter, S., Siegel, E. R., McKelvey, S.

A., & Morrison, J. C. (2013). Risk factors for a prolonged third stage of labor

and postpartum hemorrhage. *Southern Medical Journal*, 106(2), 131-135. doi:

10.1097/SMJ.0b013e3182824d1e [doi]

Masuzawa, Y. & Kataoka, Y. (2016). Uterine activity during the two hours after

placental delivery among low-risk pregnancies: An observational study. *The*

Journal of Maternal-Fetal & Neonatal Medicine, (22), 1-6.

doi:10.1080/14767058.2016.1253057

- Matsuoka, M., & Sannomiya, Y. (2002). Bunbenchokugono shikyu reianpouno koukawo kensyousuru [To examine the effect of cooling the uterus immediately after parturition]. *Nihon kangogakkai rombunsyu: Boseikango*, (33), 6-8.
- McCormick, M. L., Sanghvi, H. C. G., Kinzie, B., & McIntosh, N. (2002). Preventing postpartum hemorrhage in low-resource settings. *International Journal of Gynaecology and Obstetrics: The Official Organ of the International Federation of Gynaecology and Obstetrics*, 77(3), 267-275.
- McDonald, S. J., Middleton, P., Dowswell, T., & Morris, P. S. (2013). Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes. *The Cochrane Database of Systematic Reviews*, 7, CD004074. doi: 10.1002/14651858.CD004074.pub3

- McDonald, S., Abbott, J. M., & Higgins, S. P. (2004). Prophylactic ergometrine-oxytocin versus oxytocin for the third stage of labour. *The Cochrane Database of Systematic Reviews*, 3, CD004074. doi: 10.1002/14651858.CD004074.pub3

Database of Systematic Reviews, 2, CD000201. doi:

10.1002/14651858.CD000201.pub2.

Ministry of health, Labour and Welfare. *The National Health and Nutrition Survey in*

Japan, 2014. 2015. (In Japanese) Retrieved from

<http://www.mhlw.go.jp/bunya/kenkou/eiou/dl/h26-houkoku-05.pdf>

Mitchell, Janice L., Stecher, Jack, Crowson, James, Rich, Diana. (2015). Uterine

Cooling During Cesarean Delivery to Reduce Blood Loss and Incidence of

Postpartum Hemorrhage: A Randomized Controlled Trial. *Obstetrics &*

Gynecology: [May 2015](#) doi: 10.1097/01.AOG.0000463548.24459.52

Retrieved from

http://journals.lww.com/greenjournal/Abstract/2015/05001/Uterine_Cooling

[During Cesarean Delivery to Reduce.30.aspx](#)

Mothers' and Children's Health Organization. (2015). Boshihokenno suijunn [Level of

Mothers' and Children's Health]. In: Mothers' and Children's Health and

Welfare Association (Eds.), *Wagakunino boshihoken [Maternal health in Japan]* (pp. 16-35). Tokyo: Mothers' and Children's Health Organization.

Mori, R., Nardin, J. M., Yamamoto, N., Carroli, G., & Weeks, A. (2012). Umbilical

vein injection for the routine management of third stage of labour. *The Cochrane Database of Systematic Reviews*, 3, CD006176. doi:

10.1002/14651858.CD006176.pub2

Mustafa, S. M., & Thulesius, O. (1999). Cooling-induced bladder contraction: Studies

on isolated detrusor muscle preparations in the rat. *Urology*, 53(3), 653-657.

doi: S0090-4295(98)00568-8 [pii]

Mustafa, S. M., & Thulesius, O. (2001). Cooling-induced gastrointestinal smooth

muscle contractions in the rat. *Fundamental & Clinical Pharmacology*, 15(5),

349-354. doi:034 [pii]

Nasu, T. (1990). Effects of cooling on smooth muscle contraction. *Comparative*

Biochemistry and Physiology.A, Comparative Physiology, 95(2), 201-207.

National Institute for Health and Care Excellence (NICE). (2014). *NICE clinical*

guideline 190. Intrapartum care: care of healthy women and their babies

during childbirth. Retrieved from

<http://www.nice.org.uk/guidance/cg190/evidence>

Japan Society of Obstetrics and Gynecology. (2013). *Glossary of Obstetrics and*

Gynecology 3rd ed. Tokyo; Office of Japan Society of Obstetrics and

Gynecology.

Japan Society of Obstetrics and Gynecology, & Japan Association of Obstetricians and

Gynecologists. (2014). *Obstetrics and gynecology clinical guideline*

obstetrics version 2014.

Novikova, N., Hofmeyr, G. J., & Cluver, C. (2015). Tranexamic acid for preventing

postpartum haemorrhage. *The Cochrane Database of Systematic Reviews*, 6,

CD007872. doi: 10.1002/14651858.CD007872.pub3

Oladapo, O. T., Fawole, B., Blum, J., & Abalos, E. (2012). Advance misoprostol

distribution for preventing and treating postpartum haemorrhage. *The*

Cochrane Database of Systematic Reviews, 2, CD009336. doi:

10.1002/14651858.CD009336.pub2

Oladapo, O. T., Okusanya, B. O., & Abalos, E. (2012). Intramuscular versus

intravenous prophylactic oxytocin for the third stage of labour. *The Cochrane*

Database of Systematic Reviews, 2, CD009332. doi:

10.1002/14651858.CD009332.pub2

Olefile, K. M., Khondowe, O., & M'rithaa, D. (2013). Misoprostol for prevention and

treatment of postpartum haemorrhage: A systematic review. *Curationis*,

36(1), E1-E10. doi:10.4102/curationis.v36i1.57

Osumi, K., Horiuchi, S. (2007). The physiological and emotional responses to the care

for postpartum uterine contraction: the comparison between cold compress

application and observation without the application. *Journal of St. Luke's*

Society for Nursing Research, 11(1), 10-18.

Otsuka, Y., Fujiwara, K., Hayashi, Y., Ogawa, H., Maruyama, M., & Hamada, T.

(1990). Bunbengono kahukubureianpouni kansurukentou [Study on cooling the

lower abdomen after the delivery]. *Japanese Journal of Maternal Health*, 31(4), 629-630.

Peña-Martí, G., & Comunián-Carrasco, G. (2007). Fundal pressure versus controlled cord traction as part of the active management of the third stage of labour.

The Cochrane Database of Systematic Reviews, 4, CD005462.

Paul, M. J., & Smeltzer, J. S. (1991). Relationship of measured external tocodynamometry with measured internal uterine activity. *American Journal of Perinatology*, 8(6), 417-420.

Royal Australian and New Zealand College of Obstetricians and Gynaecologists.

(RANZCOG). (2011). *Management of postpartum hemorrhage*. Retrieved from <https://www.ranzcog.edu.au/doc/management-of-postpartum-haemorrhage>.

Royal College of Obstetricians and Gynaecologists. (RCOG). (2011). *Postpartum Haemorrhage, Prevention and Management (Green-top 52)*. Retrieved from

https://www.ranzcog.edu.au/RANZCOG_SITE/media/RANZCOG-

[MEDIA/Women's%20Health/Statement%20and%20guidelines/Clinical-
Obstetrics/RCOG-Prevention-and-management-of-postpartum-
haemorrhage.pdf?ext=.pdf](#)

Sato, Y., Noro, Y., & Nakayama, R. (1984). Kahukubureianpouto

bunbenjis Yukketuryouno kentou [The correlation of cooling the lower abdomen and blood loss at the delivery]. *Josampuzzashi*, 38(6), 509-512.

Say, L., Chou, D., Gemmill, A., Tuncalp, O., Moller, A. B., Daniels, J., . . . Alkema, L. (2014). Global causes of maternal death: A WHO systematic analysis. *The Lancet. Global Health*, 2(6), e323-e333. doi:10.1016/S2214-109X(14)70227-X

Shea, BJ., Grimshaw, JM., Wells, GA., Boers, M., Andersson, N., & Hamel, C. (2007). Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Medical Research Methodology*, 7:10. doi: 10.1186/1471-2288-7-10

Sheldon, W. R., Blum, J., Vogel, J. P., Souza, J. P., Gulmezoglu, A. M., Winikoff, B., & WHO Multicountry Survey on Maternal and Newborn Health Research

Network. (2014). Postpartum haemorrhage management, risks, and maternal outcomes: Findings from the world health organization multicountry survey on maternal and newborn health. *BJOG: An International Journal of Obstetrics and Gynaecology*, 121 Suppl 1, 5-13. doi:10.1111/1471-0528.12636

Soltani, H., Hutchon, D. R., & Poulose, T. A. (2010). Timing of prophylactic uterotronics for the third stage of labour after vaginal birth. *The Cochrane Database of Systematic Reviews*, 8, CD006173. doi: 10.1002/14651858.CD006173.pub2

Su, L., Chong, Y., & Samuel, M. (2012). Carbetocin for preventing postpartum haemorrhage. *The Cochrane Database of Systematic Reviews*, 4, CD005457. doi: 10.1002/14651858.CD005457.pub4

Suzuki, S., Hiraizumi, Y., & Miyake, H. (2012). Risk factors for postpartum hemorrhage requiring transfusion in cesarean deliveries for japanese twins:

Comparison with those for singletons. *Archives of Gynecology and Obstetrics*, 286(6), 1363-1367. doi:10.1007/s00404-012-2461-9

Tunçalp, Ö., Hofmeyr, G. J., & Gülmезoglu, A. M. (2012). Prostaglandins for preventing postpartum haemorrhage. *The Cochrane Database of Systematic Reviews*, 8, CD000494. doi: 10.1002/14651858.CD000494.pub4

Westhoff, G., Cotter, A. M., & Tolosa, J. E. (2013). Prophylactic oxytocin for the third stage of labour to prevent postpartum haemorrhage. *The Cochrane Database of Systematic Reviews*, 10, CD001808. doi: 10.1002/14651858.CD001808.pub2

World Health Organization (WHO). (2012). *WHO recommendations for the prevention and treatment of postpartum haemorrhage*. Retrieved from http://www.who.int/reproductivehealth/publications/maternal_perinatal_health/9789241548502/en/

Yaju, Y., Kataoka, Y., Eto, H., Horiuchi, S., & Mori, R. (2013). Prophylactic interventions after delivery of placenta for reducing bleeding during the

postnatal period. *The Cochrane Database of Systematic Reviews*, 11,
CD009328. doi: 10.1002/14651858.CD009328.pub2