

References

- Adib-Hajbaghery, M., Rajabi-Beheshtabad, R., & Abasi, A. (2013). Effect of whole body massage by patient's companion on the level of blood cortisol in coronary patients. *Nursing and Midwifery Studies*, 2(3), 10-15.
- Akbarian, Z., Kohan, S., Nasiri, H., & Ehsanpour, S. (2018). The effects of mental health training program on stress, anxiety, and depression during pregnancy. *Iranian Journal of Nursing and Midwifery Research*, 23(2), 93-97.
- DOI: 10.4103/ijnmr.IJNMR_207_16.
- Alonso-Alonso, M., Woods, S. C., Pelchat, M., Grigson, S. P., Stice, E., Farooqi, S., . . . Beauchamp, K. G. (2015). Food reward system: Current perspectives and future research needs. *Nutrition Reviews*, 73(5), 296-307.
- Bakermans-Kranenburg, M. J., & van IJzendoorn, M. H. (2008). Oxytocin receptor (OXTR) and serotonin transporter (5-HTT) genes associated with observed parenting. *Social Cognitive and Affective Neuroscience*, 3(2), 128-134.
- DOI: 10.1093/scan/nsn004
- ベネッセ教育総合研究所. (2007). 第1回 妊娠出産子育て基本調査 (横断調査).
- Retrieved from
https://berd.benesse.jp/jisedaiken/research/pdf/kihonC_023-047.pdf

ベネッセ教育総合研究所. (2013). 第2回 妊娠出産子育て基本調査 (横断調査).

Retrieved from https://berd.benesse.jp/jisedaiken/research/research_23/pdf/03.pdf

Carter, C. S., Pornajafi-Nazarloo, H., Kramer, K. M., Ziegler, T. E., White-Traut, R.,

Bello, D., & Schwertz, D. (2007). Oxytocin: behavioral associations and potential as a salivary biomarker. *Annals of the New York Academy of Sciences*, 1098(1), 312-322.

DOI: doi.org/10.1196/annals.1384.006

Choi, S. M., & Lee, J. E. (2015). Effects of foot-reflexology massage on fatigue, stress and postpartum depression in postpartum women. *Journal of Korean Academy of Nursing*, 45(4), 587-594.

Clements, A. D. (2013). Salivary cortisol measurement in developmental research:

Where do we go from here? *Developmental Psychobiology*, 55(3), 205-220.

DOI: doi.org/10.1002/dev.21025

Dedovic, K., Duchesne A., Andrews, J., Engert, V., & Pruessner, J.C. (2009). The Brain and the stress axis: The neural correlates of cortisol regulation in response to stress. *Neuroimage*, 47(3), 864-871.

DOI: 10.1016/j.neuroimage.2009.05.074.

de Geest, K., Thiery, M., Piron-Pussuyt, G., & Driessche, R. V. (1985). Plasma oxytocin in human pregnancy and parturition. *Journal of Perinatal Medicine*,

13(1), 3-13.

Dickerson, S. S., Kemeny, E. M. (2004). Acute stressors and cortisol responses: a

theoretical integration and synthesis of laboratory research. *Psychological Bulletin*,

130(3), 355-391.

DOI: 10.1037/0033-2909.130.3.355

Engert, V., Efanov, I. S., Dedovic, K., Dagher, A., & Pruessner, J. C. (2011). Increased

cortisol awakening response and afternoon/evening cortisol output in healthy

young adults with low early life parental care. *Psychopharmacology*, 214(1), 261-

268.

DOI: 10.1007/s00213-010-1918-4.

Feldman, R., Weller, A., Zagoory-Sharon, O., & Levin, A. (2007). Evidence for a

neuroendocrinological foundation on human affiliation. Plasma oxytocin levels

across pregnancy and the postpartum period predict mother-infant bonding.

Psychological Science, 18(11), 965-970.

DOI: 10.1111/j.1467-9280.2007.02010.x

Feldman, R., Gordon, I., Schneiderman, I., Weisman, O., & Zagoory-Sharon, O.

(2010). Natural variations in maternal and paternal care are associated with systematic changes in oxytocin following parent-infant contact.

Psychoneuroendocrinology, 35(8), 1133-1141.

DOI: 10.1016/j.psyneuen.2010.01.013.

Feldman, R., Zagoory-Sharon, O., Weisman, O., Schneiderman, I., Gordon, I., Maoz,

R., Ebstein, R. P. (2012). Sensitive parenting is associated with plasma oxytocin and polymorphisms in the OXTR and CD38 genes. *Biological Psychiatry*, 72(3), 175-181.

DOI: doi.org/10.1016/j.biopsych.2011.12.025

Ferguson, A. C. (1964). Baby talk in six languages. *American Anthropologist*, 66(6),

103-114.

DOI: 10.1525/aa.1964.66.suppl_3.02a00060

Field, T., Diego, M., Delgado, J., & Medina, L. (2013). Yoga and social support reduce

prenatal depression, anxiety and cortisol. *Journal of Bodywork & Movement Therapies*, 17, 397-403.

Fuchs, A. R., Behrens, O., & Liu, H.C. (1992). Correlation of nocturnal increase in plasma oxytocin with a decrease in plasma estradiol/progesterone ratio in late pregnancy. *American Journal of Obstetrics & Gynecology*, 167(6), 1559-1563.

Guney, E. & Ucar, T. (2018). Effect of the fetal movement count on maternal-fetal attachment. *Japan Journal of Nursing Science*, 16(1), 71-79.

DOI: 10.1111/jjns.12214.

Gordon, I., Zagoory-Sharon, O., Leckman, F. J., & Feldman, R. (2010). Oxytocin and the development of parenting in humans. *Biological Psychiatry*, 68(4), 377-382.

DOI: 10.1016/j.biopsych.2010.02.005.

花沢成一. (1992). 母性心理学. 東京: 医学書院.

肥田野直, 福原眞知子, 岩脇三良, 曾我祥子, Spielberger, C. D. (200). 新版 STAI マニュアル. 東京: 実務教育出版.

Holt-Lunstad, J., Birmingham, A. W., & Light, C. K. (2008). Influence of a “Warm Touch” support enhancement intervention among married couple on ambulatory blood pressure, oxytocin, alpha amylase, and cortisol. *Psychosomatic Medicine*, 70(9), 976-985.

DOI: doi.org/10.1097/PSY.0b013e318187aef7

堀内成子, 田所由利子, 高畠香織. (2016). 唾液オキシトシン測定のための検体採取法の検討. 第 57 回日本母性衛生学会(Vol.57, p.251). 東京, 日本.

Jing, Z., Sasaki, A., & Sato, C. (2006). Infertility treatments and anxiety and feelings for the baby in pregnant women. *Journal of Japan Academy of Midwifery*, 20(1), 99-106.

DOI: doi.org/10.3418/jjam.20.1_99

角森輝美, 山口洋史. (2012). 初産婦の妊娠期対児感情 赤ちゃんに対する正のイメージが強化される要因. リハビリテーション連携科学, 13(1), 12-20.

萱間真美. (2007). 質的研究実践ノート - 研究プロセスを進める clue とポイント . 医学書院: 東京.

Kudielka, M. B., Buske-Kirschbaum, A., Hellhammer, D. H. & Kirschbaum, C. (2004). HPA axis responses to laboratory psychosocial stress in healthy elderly adults, younger adults, and children: Impact of age and gender. *Psychoneuroendocrinology*, 29(1), 83-98.

DOI: 10.1016/S0306-4530(02)00146-4

Kudielka, M. B., Hellhammer, H. D., & Wust, S. (2009). Why do we respond so differently? Reviewing determinants of human salivary cortisol responses to challenge. *Psychoneuroendocrinology*, 34(1), 2-18.

DOI: 10.1016/j.psyneuen.2008.10.004.

熊谷さなえ, 大沢妙子, 大崎美紀子. (2004). 切迫早産で入院した妊婦への"新生児ふれあい体験"の効果. 日本看護学会論文集: 母性看護, 35, 90-92.

厚生労働省. (2003). 厚生労働白書 第2章子どもをとりまく現状/課題.

Retrieved from <http://www.mhlw.go.jp/wp/hakusyo/kousei/03/>

Kusaka, M., Matsuzaki, M., Shiraishi, M. & Haruna, M. (2016). Immediate stress reduction effects of yoga during pregnancy: One group pre-post test. *Women and Birth*, 29(5), 82-88.

DOI: 10.1016/j.wombi.2016.04.003.

Levine, A., Zagoory-Sharon, O., Feldman, R., & Weller, A. (2007). Oxytocin during pregnancy and early postpartum: Individual patterns and maternal-fetal attachment. *Peptides*, 28(6), 1162-1169.

Light, K. C., Grewen, K. M., & Amico, J. A. (2005). More frequent partner hugs and higher oxytocin levels are linked to lower blood pressure and heart rate in premenopausal women. *Biological Psychology*, 69(1), 5-21.

DOI: 10.1016/j.biopspsycho.2004.11.002

Lindow, S. W., Newham, A., Hendricks, M. S., Thompson, J. W., & Spuy, d. v. (1996).

The 24-hour rhythm of oxytocin and beta-endorphin secretion in human

pregnancy. *Clinical Endocrinology (Oxf)*, 45(4), 443-446.

松岡治子, 行田智子, 今関節子, 横田正夫. (2002). 妊娠期・産褥期・育児期の母

親の不安について 日本版 STAI を用いた横断的研究. *母性衛生*, 43(1), 13-

17.

Migeon, C. J., Bertrand, J., & Wall, P. E. (1957). Physiological disposition of 4-C14-

cortisol during late pregnancy. *Journal of Clinical Investigation*, 36(9), 1350-1362.

Moberg, K.U. (1996). Neuroendocrinology of the mother-child interaction. *Trends in*

Endocrinology & Metabolism, 7(4), 126-131.

Muramatsu, K., Muyaoka, H., Muramatsu, Y., Yoshida, M., Ostubo, T., & Gejyo, F.

(2007). The patient health questionnaire, Japanese version: Validity according to

the mini-international neuropsychiatric interview-plus. *Psychological Reports*,

952-960. DOI:10.2466/pr0.101.3.952-960

中嶋カツエ, 小林益江, 田中佳代. (1998). 初産婦の妊娠期の不安. *母性衛生*,

39(2), 250-254.

Nestler, J. E. (2004). Historical review: Molecular and cellular mechanisms of opiate

and cocaine addiction. *Trends in Pharmacological Sciences*, 25, 210–218.

Doi: 10.1016/j.tips.2004.02.005

西谷正太, 木田哲夫, 高村恒人, 篠原一之. (2013). 家族「愛」の神経基盤. 分子精神医学, 13(4), 236-242.

O'Connor, T. G., Heron, J., Golding, J., & Glover, V. (2003). Maternal antenatal anxiety and behavioral/emotional problems in children: A test of a programming hypothesis. *Journal of Child Psychology and Psychiatry*, 47, 1025-1036.

小川雅美. (1991). PBI (Parental Bonding Instrument) 日本版の信頼性, 妥当性に関する研究. 精神科治療学, 6(10), 1193-1201.

大森あゆ美, 石原美恵子, 杉村貴子, 福力純子. (2005). 当院における妊娠期の育児不安に関する調査~赤ちゃん同窓会に参加する意義~. 岡山県母性衛生, 21, 43-44.

Oyola, G. M., & Handa, J. R. (2017). Hypothalamic-pituitary-adrenal and hypothalamic-pituitary-gonadal axes: sex differences in regulation of stress responsivity. *Stress*, 20(5), 476-494.

Doi: 10.1080/10253890.2017.1369523

Pinar, R., & Afsar, F. (2015). Back massage to decrease State Anxiety, cortisol level, blood pressure, heart rate and increase sleep quality in family caregivers of patients

with cancer: A randomized controlled trial. *Asian Pacific Journal of Cancer Prevention*, 16(18), 127-133.

Rapaport, M. H., Schettler, P., & Bresee C. (2012). A preliminary study of the effects of repeated massage on hypothalamic-pituitary-adrenal and immune function in healthy individuals: A study of mechanisms of action and dosage. *Journal of Alternative Complementary Medicine*, 18(8), 789-797.

Doi: doi.org/10.1089/acm.2011.0071

Rubin, R. (1984). Shindo, Y & Goto, K. /新道幸恵, 後藤桂子. (1997). 母性論 母性の主観的体験. 東京: 医学書院.

Sakamoto, S., Sakamori, S., & Nakamura, J. (2018). Physiological changes related to attachment formation between parents and children by touching objects with palm (hairless part). The 59th Annual Meeting of Japan Society of Maternal Medicine (Vol 59, p. 312) Niigata, Japan.

Sanaati, F., Charandabi, M.S., Eslamlo, F.H, & Mirghafourvand, M. (2018). A randomized controlled trial on the effect of lifestyle education for Iranian women and their husbands on post-partum anxiety and depression. *Health Education Research*, 33(5), 416-428.

Doi: 10.1093/her/cyy026.

佐藤喜根子. (2006). 産褥期にある女性の不安要因の分析. 東北大学医学部保健
学科紀要, 15(2), 113-124.

佐藤奈緒子, 森岡由起子, 佐藤文, 生地新, 村田亜美. (2006). 産後うつ状態に影
響を及ぼす背景因子についての縦断的研究 (第一報) 母親自身の被養育体
験・内的ワーキングモデルおよび児への愛着との関連. 母性衛生, 47(2),
320-329.

盛山幸子, 島田三恵子. (2008). 妊娠先行結婚と妊婦の対児感情・母親役割獲
得・夫婦関係との関連. 日本助産学会誌, 22(2), 222-232.

Severi, F. M., Prattichizzo, D., Casarosa, E., Brabagli, F., Ferretti, C., Altomare, A.,
Vicino, A., & Petraglia, F. (2005). Virtual fetal touch through a haptic interface
decrease maternal anxiety and salivary cortisol. *Journal of the Society for
Gynecologic Investigation*, 12, 37-40.

Scofield, D. M., & Kalivas, W. P. (2014). Astrocytic dysfunction and addiction:
Consequences of impaired glutamate homeostasis. *Neuroscientist*, 20, 610–622.
Doi: 10.1177/1073858413520347.

園田希, 小川真世, 田所由利子, 高畠香織, 周尾卓也, 堀内成子. (2018). 妊婦を
対象としたふれて・感じる「Mama's Touch プログラム」の実行可能性 -オ

キシトシン・コルチゾールによる評価: 予備研究-. 日本助産学会, 32(1), 60-72.

DOI: doi.org/10.3418/jjam.JJAM-2017-0029

Stevens, L. F., Wiesman, O., Feldman, R., Hurley, A. R., & Taber, H. K. (2013).

Oxytocin and Behavior: Evidence for Effects in the Brain. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 25(2), 96-102.

DOI: 10.1176/appi.neuropsych.13030061.

鈴木幸子, 島田三恵子. (2013). 初めて出産を迎える妊娠末期の妊婦とその夫における夫婦の愛情と対児感情および母親役割行動との関連. 小児保健研究, 72(3), 405-412.

Tadokoro, Y. (2016). Effects of clary sage and lavender or jasmine essential oils infused in a foot bath for labor induction on oxytocin level. (Unpublished doctoral dissertation), St. Luke's International University, Tokyo Japan.

Takagi, T., Tanizawa, O., Otsuki, Y., Sugita, N., Haruta, M., & Yamaji, K. (1985). Oxytocin in the cerebrospinal fluid and plasma of pregnant and nonpregnant subjects. *Hormone and Metabolic Research*, 17(6), 308-310.

Doi: 10.1055/s-2007-1013526

Takeda, S., Kuwabara, Y., & Mizuno, M. (1985). Effects of pregnancy and labor on oxytocin levels in human plasma and cerebrospinal fluid. *Endocrine Journal*, 32(6), 875-880.

立岡弓子, 高橋真理. (2004). 妊娠経過における唾液中ストレス関連ホルモン濃度の縦断的調査. *母性衛生*, 45(1), 112-117.

Terkawi, A. S., Jackson, W. M., Thiet, M., Hansoti, S., Tabassum, R., & Flood, P. (2012). Oxytocin and catechol-O-methyltransferase receptor genotype predict the length of the first stage of labor. *American Journal of Obstetrics and Gynecology*, 207(3), 184-184.

Doi: doi.org/10.1016/j.ajog.2012.06.079

Urizar Jr, G. G., Milazzo, M., Le, H., Delucchi, K., Sotel, R., & Munoz, R. F. (2004). Impact of stress reduction instructions on stress and cortisol levels during pregnancy. *Biological Psychology*, 67, 275-282.

DOI: [10.1016/j.biopspsycho.2003.11.001](https://doi.org/10.1016/j.biopspsycho.2003.11.001)

van IJzendoorn, M. H., Bhandari, R., van der Veen., R., Grewen, K. M, & Bakermans-Kranenburg, M. J. (2012). Elevated salivary levels of oxytocin persist more than 7h after intranasal administration. *Frontiers in Neuroscience*, 6(174), 1-6.

DOI: [10.3389/fnins.2012.00174](https://doi.org/10.3389/fnins.2012.00174)

Viru, A. (1992). Plasma hormones and physical exercise. *International Journal Sports Medicine*, 13(3), 201-209.

DOI: 10.1055/s-2007-1021254

Voegtlind, K. M., Costigan, K. A., Kivlighan, K. T., Laudenslger, M. L., Henderson, J.

L., & Dipietro, J. A. (2013). Concurrent levels of maternal salivary cortisol are

unrelated to self-reported psychological measures in low-risk pregnant women.

Archives of Women's Mental Health, 16(2), 101-108.

DOI: 10.1007/s00737-012-0321-z.

Walker, S. C., Trotter, P. D., Swaney, W. T., Marshall, A., & McGlone, F. P. (2017). C-

tactile afferents: Cutaneous mediators of oxytocin release during affiliative tactile

interactions?. *Neuropeptides*, 64, 27-38.

渡邊奈津子, 田口理恵[袴田], 真子春菜, 糸井和佳, 河原智江, 臺有桂, 今松友紀,

田高悦子. (2013). 地域における初妊婦の育児中の母親との交流における体

験. *保健師ジャーナル*, 69(1), 60-66.

DOI: doi.org/10.11477/mf.1664102055

Weiland, J. B., Heitzeg, M. M., Zald D., Cumminford, C., Love, T., Zucker, A. R.,

& Zubieta, K. J. (2014). Relationship between impulsivity, prefrontal anticipatory

activation, and striatal dopamine release during rewarded task performance. *Psychiatry Research*, 223(3), 244–252.

Werker, F. J., & McLeod, J. P. (1989). Infant preference for both male and female infant-directed talk: a developmental study of attentional and affective responsiveness. *Canadian Journal of Experimental Psychology*, 43(2), 230-246.

DOI: 10.1016/j.psychresns.2014.05.015.

Wise, A. R., & Koob, F. G. (2014). The development and maintenance of drug addiction. *Neuropsychopharmacology*, 39, 254-262.

DOI: 10.1038/npp.2013.261

Xiaoxi, L., Kawamura, Y., Shimada, T., Otowa, T., Koishi, S. & Sugiyama, T., et al. (2010). Association of the oxytocin receptor (OXTR) gene polymorphisms with autism spectrum disorder (ASD) in the Japanese population. *Journal of Human Genetics*, 55(3), 137-141.