

References

Articles marked with an asterisk indicate studies included in the systematic review

Arai K. (2018). Necessity of kampo lifelong education considering the awareness of pharmacists. *The Journal of Kampo Medicine & Pharmacy*, 52,53.

*Chubachi, S. (2015). A case of mesenteric phlebosclerosis complicated by perforative peritonitis in a patient with a history of intake of Chinese herbal medicine. *Journal of Japan Surgical Association*, 76(5), 1196 - 1200. <https://doi.org/10.3919/jjsa.76.1196>

*Endo, S., Sasaki, Y., Mori, K., Ichihara, A., Inoue, S., & Iida, S. (2015). A case of asymptomatic idiopathic mesenteric phlebosclerosis associated with long-term administration of Kamishoyosan that initially manifested as a positive fecal test for occult blood. *Progress of Digestive Endoscopy*.

*Fujieda, Y., Maeda, H., Kawanishi, Y., Tokumaru, T., Kuwahara, M., Akimori, T., Kamioka, N., & Hanazaki, K. (2019). A case of mesenteric phlebosclerosis induced by long-term use of Kampo medicine. *Annals of Cancer Research and Therapy*, 27(1), 12–14. <https://doi.org/10.4993/acrt.27.12>

Hiramatsu, K., Sakata, H., Horita, Y., Orita, N., Kida, A., Mizukami, A., Miyazawa, M., Hirai, S., Shimatani, A., Matsuda, K., Matsuda, M., Ogino, H., Fujinaga, H., Terada, I., Shimizu, K., Uchiyama, A., Ishizawa, S., Abo, H., Demachi, H., & Noda, Y. (2012a). Mesenteric phlebosclerosis associated with long-term oral intake of geniposide, an ingredient of herbal medicine. *Alimentary Pharmacology and Therapeutics*, 36(6), 575–586. <https://doi.org/10.1111/j.1365-2036.2012.05221.x>

Hiramatsu, K., Sakata, H., Horita, Y., Orita, N., Kida, A., Mizukami, A., Miyazawa, M., Hirai, S., Shimatani, A., Matsuda, K., Matsuda, M., Ogino, H., Fujinaga, H., Terada, I., Shimizu, K., Uchiyama, A., Ishizawa, S., Abo, H., Demachi, H., & Noda, Y. (2012b).

Mesenteric phlebosclerosis associated with long-term oral intake of geniposide, an ingredient of herbal medicine. *Alimentary Pharmacology & Therapeutics*, 36(6), 575–586.

<https://doi.org/10.1111/j.1365-2036.2012.05221.x>

*Hoshino, N., Hasegawa, S., Hida, K., Kawada, K., & Sakai, Y. (2016). Right hemicolectomy for mesenteric phlebosclerosis potentially caused by long-term use of herbal medicine: A case report and literature review. *International Journal of Surgery Case Reports*, 24, 191–194. <https://doi.org/10.1016/j.ijscr.2016.05.051>

Im, M., Kim, A., & Ma, J. Y. (2016). Ethanol extract of baked Gardeniae Fructus exhibits in vitro and in vivo anti-metastatic and anti-angiogenic activities in malignant cancer cells: Role of suppression of the NF-B and HIF-1 pathways. *International Journal of Oncology*, 49(6), 2377–2386. <https://doi.org/10.3892/ijo.2016.3742>

Iwashita, A., Yao, T., Schlemper, R. J., Kuwano, Y., Yao, T., Iida, M., Matsumoto, T., & Kikuchi, M. (2003). Mesenteric phlebosclerosis: A new disease entity causing ischemic colitis. *Diseases of the Colon and Rectum*, 46(2), 209–220.

<https://doi.org/10.1097/01.DCR.0000044720.43258.6E>

Joanna Briggs Institute (JBI). (2020). *Checklist for case reports critical appraisal tools for use in JBI systematic reviews*. https://joannabriggs.org/sites/default/files/2019-05/JBI_Critical_Appraisal-Checklist_for_Case_Reports2017_0.pdf

Japan Kampo Medical Association (JKMA). (2011). *Summary Report of prescription status of Kampo*. <https://www.nikkankyo.org/serv/serv1.htm>

*Kakimoto, T., Takeo, S., & Miura, O. (2017). A case of idiopathic mesenteric phlebosclerosis with transverse colon cancer. *Journal of Japanese College of Surgeons*.

*Kayano, H., Nomura, E., Hiraiwa, S., Kuramoto, T., Yatabe, K., Machida, T., Tajiri, T., Mukai, M., & Makuchi, H. (2016). A case of idiopathic mesenteric phlebosclerosis with progressive intestinal necrosis. In *Tokai Journal Experimental Clinical Medicine*, 41(2).

Kinoshita Yuko. (2006). Kampo for gender specific medicine. *Obstetrical and Gynecological Therapy*.

*Kodera T., Ueo T., Ishida T., Nagamatsu H., Takahashi K., Urabe M., Narita R., Mukai T.,

Yonemasu H., & Wakasugi K. (2012). A case of idiopathic mesenteric phlebosclerosis in a young female caused by herbal medicine. *Gastroenterological Endoscopy*.

<https://www.embase.com/a/#/search/results?subaction=viewrecord&id=L364622785&from=export>

*Kuga Y, Okanobu H, & Kitamura S. (2011). A case of idiopathic mesenteric phlebosclerosis. *Clinic All-Round*.

<http://mol.medicalonline.jp/library/journal/download?GoodsID=af2sgrsa/2011/006011/034&name=2352-2355j&UserID=1100006523-81>

*Mantani, N., Minami, K., Yamamoto, N., Kimura, Y., & Ito, T. (2017). A case of mesenteric phlebosclerosis found by ultrasonography: A possible method of screening. *Japan Journal Medical Ultrasonics*.

https://mol.medicalonline.jp/library/journal/download?GoodsID=ed5choon/2017/004402/011&name=0179-0184j&UserID=1100006523-81&base=jamas_pdf

*Minami, K., Fujiiie, M., Fushimi, F., Taguchi, K., Nobutoh, Y., Yoshida, D., Ota, M., Ikebe, M., Morita, M., & Toh, Y. (2017). Ascending colon cancer coincident with mesenteric phlebosclerosis associated with the long-term oral intake of Chinese herb containing gardenia fruit: A case report and literature review. *International Cancer Conference Journal*, 6(2), 70–75. <https://doi.org/10.1007/s13691-017-0277-y>

Ministry of Health, Labor and Welfare. (2018). *Pharmaceuticals and Medical Devices Safety Information*.

Ministry of Health, Labor and Welfare (2019). *Summary of pharmaceutical affairs industry production statistics annual report in the first year of Reiwa.*

<https://www.mhlw.go.jp/topics/yakuji/2019/nenpo/>

*Miwa, W., Hiratsuka, T., Kamiya, Y., Shoji, H., Tei, S., Dairaku, K., Shiraishi, H., Fujimoto, T., Sato, K., & Kato, Y. (2015). Translated title improved mesenteric phlebosclerosis after cessation of Chinese herbal medicine, a case report with an observation period of 7 years from onset. *Gastroenterological Endoscopy*.

<https://www.embase.com/a/#/search/results?subaction=viewrecord&id=L605410715&from=export>

Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA statement. *PLoS Medicine*, 6(7), e1000097. <https://doi.org/10.1371/journal.pmed.1000097>

*Moriyama, T., Shimizu, M., Fujikawa, T., Kato, K., & Nagasu, N. (2017). Mesenteric phlebosclerosis improved by discontinuation of kamisyoyosan, presented with a positive fecal occult blood test. *Progress of Digestive Endoscopy*.

Moschik, E. C., Mercado, C., Yoshino, T., Matsuura, K., & Watanabe, K. (2012). Usage and attitudes of physicians in Japan concerning traditional Japanese medicine (kampo medicine): A descriptive evaluation of a representative questionnaire-based survey. *Evidenced Based Complementary Alternative Medicine*. *eCam* 12 139818.

<https://doi.org/10.1155/2012/139818>

*Nagasue, T., Kurahara, K., Yaita, H., Oshiro, Y., Kochi, S., Morishita, T., Kunou, N., Abe, H., Harada, A., & Iwasaki, K. (2014). Two cases of idiopathic mesenteric phlebosclerosis associated with long-term use of Chinese herbal medicine. *The Matsuyama Red Cross Hospital Journal of Medicine*.

Nagata, Y., Watanabe, T., Nagasaka, K., Yamada, M., Murai, M., Takeuchi, S., Murase, M., Yazaki, T., Murase, T., Komatsu, K., Kaizuka, M., Sano, M., Asano, K., Ando, C., & Taniuchi, N. (2016). Total dosage of gardenia fruit used by patients with mesenteric phlebosclerosis. *BMC Complementary and Alternative Medicine*, 16(1), 1–11.
<https://doi.org/10.1186/s12906-016-1182-1>

*Nakazawa K., Shingaki N., Maekita T., Enomoto S., Mori Y., Oota U., & Bunno M. (2016). A case of idiopathic mesenteric phlebosclerosis that was successfully treated with the combination therapy of warfarin and aspirin. *Gastroenterological Endoscopy*.
<https://www.embase.com/a/#/search/results?subaction=viewrecord&id=L616861291&from=export>

*Nomura K., Kikuchi D., Iizuka T., Yamada A., Furuhata T., Domon K., Yamashita S., Nakamura M., Matsui A., Mitani T., Ogawa O., & Kaise M. (2012). Idiopathic mesenteric phlebosclerosis associated with long-term use of Chinese herbs: A case report. *Journal of Japanese Society of Gastroenterology*.

<https://www.embase.com/a/#/search/results?subaction=viewrecord&id=L365771720&from=export>

*Ohe, R., Kabasawa, T., Suto, A., Yang, S., Kato, T., & Osakabe, M. (2014). A case of idiopathic mesenteric venous sclerosis leading to subtotal colectomy. *The Japanese Society of Pathology*.

Ohtsu, K., Matsui, T., Nishimura, T., Hirai, F., Ikeda, K., Iwashita, A., Yorioka, M., Hatakeyama, S., Hoashi, T., Koga, Y., Sakurai, T., & Miyaoka, M. (2014). Association between mesenteric phlebosclerosis and Chinese herbal medicine intake. *Journal of Japanese Society of Gastroenterology*, 111(1), 61–68. <https://doi.org/10.11405/nisshoshi.111.61>

*Okano, T., Ochiai, J., & Yokoshima, T. (2013). A case of idiopathic mesenteric phlebosclerosis. *Medical Journal of Iwate Prefectural Hospital*.

Okumi, H., & Koyama, A. (2014). Kampo medicine for palliative care in Japan.

BioPsychoSocial Medicine, 8(1), 4–8. <https://doi.org/10.1186/1751-0759-8-6>

*Ooki, U., Sugitani, K., Yoshida, Y., & Takahashi, T. (2014). Two cases of idiopathic mesenteric phlebosclerosis possibly associated with long-term use of a Chinese herbal medicine. *Journal of Japan Surgical Association*.

Oonishi, Y., Nakao, H., Toyama, M., Higuchi, S., Kobayashi, K., & Fukuda, M. (2014).

*Questionnaire survey of general physicians regarding herbal Japanese medicine Presented by Medical*Online*.

Pharmaceuticals and Medical Devices Agency (PMDA). (n.d.). *Search package inserts in Japanese| the Pharmaceuticals and Medical Devices Agency (PMDA)*. Retrieved January 5, 2021, from <https://www.pmda.go.jp/PmdaSearch/iyakuSearch/>

Saeki, H., Oiso, N., Honma, M., Iizuka, H., Kawada, A., & Tamaki, K. (2009). Prevalence of atopic dermatitis in Japanese adults and community validation of the U.K. diagnostic criteria. *Journal of Dermatological Science* 55(2), 140 – 141.

<https://doi.org/10.1016/j.jdermsci.2009.04.004>

Shimizu, S., Kobayashi, T., Tomioka, H., Ohtsu, K., Matsui, T., & Hibi, T. (2017).

Involvement of herbal medicine as a cause of mesenteric phlebosclerosis: results from a large-scale nationwide survey. *Journal of Gastroenterology*, 52(3), 308–314.

<https://doi.org/10.1007/s00535-016-1218-9>

*Takeda, K., Nakagawa, T., Yamada, T., Konishi, K., Okuyama, M., & Nishijima, J. (2015). A case report of idiopathic mesenteric phlebosclerosis associated with long-term use of a Chinese medicine. *Journal of Japan Surgical Association*.

*Tanikawa, A., Fujita, S., Ootsuka, H., Osaka, I., Takuma, K., & Morita, Y. (2019). A case of transverse colon cancer with idiopathic mesenteric phlebosclerosis. *Journal of Japanese Society of Gastroenterology*, 116(7), 592–596. <https://doi.org/10.11405/nisshoshi.116.592>

Watanabe, K., Matsuura, K., Gao, P., Hottenbacher, L., Tokunaga, H., Nishimura, K., Imazu, Y., Reissenweber, H., & Witt, C. M. (2011). Review article traditional Japanese kampo medicine: Clinical research between modernity and traditional medicine-the state of research and methodological suggestions for the future. *Evidenced-based Complementary and Alternative Medicine*. (2011). <https://doi.org/10.1093/ecam/neq067>

Watanabe, T., Nagata, Y., Fukuda, H., & Nagasaka, K. (2016). Screening for idiopathic mesenteric phlebosclerosis in outpatients undergoing long-term treatment at the department of kampo medicine. *Kampo Medicine*, 67(3), 230–243.
<https://doi.org/10.3937/kampomed.67.230>

Tables and Figure

Table 1: Inclusion and Exclusion Criteria

	Inclusion criteria	Exclusion criteria
Population	Any patients diagnosed with mesenteric phlebosclerosis Diagnosis with imaging and histological findings (endoscopy, barium enema X-ray, plain X-ray, CT, biopsy)	Patients with pre-existing intestinal diseases
Exposure	Having a history of intake of herbal (Kampo) medicines containing gardenia fruit	No data about duration of intake, name of herbal (Kampo) medicine containing gardenia fruit
Outcome	Mortality (all-cause) Exacerbation (imaging/subjective symptoms) Amelioration (imaging/subjective symptoms)	N/A
Study design	Case report	Other than case report