

References

- Adamson, James. (2020). Cognitive behavioural therapy for chronic fatigue and chronic fatigue syndrome: outcomes from a specialist clinic in the UK. *Journal of the Royal Society of Medicine*, 394–402.
- Andersson, Gerhard. (2014). Advantages and limitations of Internet-based interventions for common mental disorders. *World Psychiatry*, 4–11.
- Arnberg, F. K. (2014). *Internet-delivered psychological treatments for mood and anxiety disorders: a systematic review of their efficacy, safety, and cost-effectiveness*. PLoS ONE.
- Baos, Sarah. (2018). Investigating the effectiveness and cost-effectiveness of FITNET-NHS (Fatigue In Teenagers on the interNET in the NHS) compared to Activity Management to treat paediatric chronic fatigue syndrome (CFS)/myalgic encephalomyelitis (ME): protocol for a random. *Trials*, 136.
- Beck, J. S. (2021, 1 8). *A Brief History of Aaron T. Beck, MD, and Cognitive Behavior Therapy*. Retrieved from A Brief History of Aaron T. Beck, MD, and Cognitive Behavior Therapy:
<https://cpe.psychopen.eu/index.php/cpe/article/view/6701/6701.html>

Benjamin, Courtney L. (2012). History of Cognitive–Behavioral Therapy (CBT) in Youth. *Child and Adolescent Psychiatric Clinics of North America*.

Burholt, Vanessa. (2011). Short Form 36 (SF–36) Health Survey Questionnaire:

normative data for Wales. *Journal of Public Health*, 587–603. 参照先:

Wikimedia: <https://en.wikipedia.org/wiki/SF-36>

Butler, S. (1991). Cognitive behaviour therapy in chronic fatigue syndrome. *Journal of Neurology, Neurosurgery, and Psychiatry*.

CDC. (2022). *Centers for Disease Control and Prevention*. Retrieved from

Understanding History of Case Definitions and Criteria:

https://www.cdc.gov/me-cfs/healthcare-providers/case-definitions-criteria.html#anchor_1542654962

Center, U. D. (2020, July 7). *COVID fatigue' is hitting hard. Fighting it is hard, too,*

says UC Davis Health psychologist. Retrieved from UC Davis Health:

<https://health.ucdavis.edu/coronavirus/news/headlines/covid-fatigue-is-hitting-hard-fighting-it-is-hard-too-says-uc-davis-health-psychologist/2020/07>

Chand, S. P. (2021, August). *Cognitive Behavior Therapy*. Retrieved from Cognitive

Behavior Therapy: <https://www.ncbi.nlm.nih.gov/books/NBK470241/>

- Deale, A. (1997). Cognitive behavior therapy for chronic fatigue syndrome: a randomized controlled trial. *American Journal of Psychiatry*, 408–14.
- Furukawa, T. A. (2021). Dismantling, optimising, and personalising internet cognitive behavioural therapy for depression: a systematic review and component network meta-analysis using individual participant data. *The Lancet Psychiatry*, 500–511.
- Del Giacco, Luca. (2020). The Action of Verbal and Non-verbal Communication in the Therapeutic Alliance Construction: A Mixed Methods Approach to Assess the Initial Interactions With Depressed Patients. *Frontiers in Psychology*.
- Gratzer, D. (2016). Internet-delivered cognitive behavioural therapy in the treatment of psychiatric illness. *CMAJ*, 263–272.
- Green, Sally and Higgins, Julian PT. (March 2011). Cochrane Handbook for Systematic Reviews of Interventions. Reference: Cochrane Handbook for Systematic Reviews of Interventions: <https://handbook-5-1.cochrane.org>
- Haythornthwaite, Jennifer A. (2013). Assessment of Pain Beliefs, Coping, and Function. *Wall & Melzack's Textbook of Pain*, 328–338.

- Hofmann, S. G. (2015). Treatments for social anxiety disorder: considerations regarding psychodynamic therapy findings. *American Journal of Psychiatry*, 393.
- Janse A. (2018). Efficacy of web-based cognitive-behavioural therapy for chronic fatigue syndrome: randomised controlled trial. *British Journal of Psychiatry*, 112–118.
- Kawakami, Norito. (1998). Prevalence of Chronic Fatigue Syndrome in a Community Population in Japan. *The Tohoku Journal of Experimental Medicine*, 33–41.
- Kim, D.Y. (2020). *Systematic review of randomized controlled trials for chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME)*. *Journal of Translational Medicine*.
- Kim, S. W. (1992). The Symptom Checklist-90: Obsessive-Compulsive Subscale: A reliability and validity study. *Psychiatry Research*, 37–44.
- Kobayashi, Lindsay C. (2016). Aging and Functional Health Literacy: A Systematic Review and Meta-Analysis. *The Journals of Gerontology*, 445–57.
- Lynch, D. (2009). Cognitive behavioural therapy for major psychiatric disorder: does it really work? A meta-analytical review of well-controlled trials. *Psychological Medicine*, 9–24.

- Nazarboland, Neda. (2017). Effectiveness of Cognitive Behavioral Therapy for Multiple Sclerosis (MS) Fatigue. *International Journal of Behavioral Sciences*, 33–42.
- Physiopedia. (2022). Physiopedia. 参照先: Sickness Impact Profile (SIP): [https://www.physio-pedia.com/Sickness_Impact_Profile_\(SIP\)](https://www.physio-pedia.com/Sickness_Impact_Profile_(SIP))
- Price, Jonathan R. (2008). Cognitive behaviour therapy for chronic fatigue syndrome in adults. *Cochrane Database of Systematic Reviews*.
- Prins, Judith B. (2001). Cognitive behaviour therapy for chronic fatigue syndrome: a multicentre randomised controlled trial. *The Lancet*, 841–7.
- PRISMA. (2021). *PRISMA Checklist*. Retrieved from PRISMA: <http://prisma-statement.org/PRISMAStatement/Checklist>
- Bansal, Rashika. (2022). COVID-19 and chronic fatigue syndrome: An endocrine perspective. *Journal of Clinical & Translational Endocrinology*.
- Staff, M. C. (2022, 1). *Cognitive behavioral therapy*. Retrieved from MAYO CLINIC: <https://www.mayoclinic.org/tests-procedures/cognitive-behavioral-therapy/about/pac-20384610>
- Sweetman, A. (2021). Effect of depression, anxiety, and stress symptoms on response to cognitive behavioral therapy for insomnia in patients with

comorbid insomnia and sleep apnea: a randomized controlled trial. *Journal of Clinical Sleep Medicine*, 545–554.

Tate, D. F. (2009). Cost Effectiveness of Internet Interventions: Review and Recommendations. *Annals of Behavioral Medicine*, 40–45.

Thomas, P. W. (2010). *Multi-centre parallel arm randomised controlled trial to assess the effectiveness and cost-effectiveness of a group-based cognitive behavioural approach to managing fatigue in people with multiple sclerosis*. BMC Neurology .

What is ME/CFS. (2021, January). Retrieved from Centers for Disease Control and Prevention: <https://www.cdc.gov/me-cfs/about/index.html>

WHO. (March 12, 2020). WHO announces COVID-19 outbreak a pandemic.

Reference: World Health Organization Regional Office for Europe:
<https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who-announces-covid-19-outbreak-a-pandemic>

Wiborg, Jan F. (2015). Randomised controlled trial of cognitive behaviour therapy delivered in groups of patients with chronic fatigue syndrome. *Psychotherapy and Psychosomatics* , 368–76.

Wind, T. R. (2020). *The COVID-19 pandemic: The 'black swan' for mental health care and a turning point for e-health*. Retrieved from Therapists Make the Switch to Telepsychology to Safely Continue Treating Their Patients During the COVID-19 Pandemic. Virtual Reality Telepsychology May Be Next: <https://www.frontiersin.org/articles/10.3389/frvir.2020.576421/full>

Worm-Smeitink, M. (2017). The assessment of fatigue: Psychometric qualities and norms for the Checklist individual strength. *Journal of Psychosomatic Research*, 40-46.

Worm-Smeitink, Margreet. (2019). Internet-Based Cognitive Behavioral Therapy in Stepped Care for Chronic Fatigue Syndrome: Randomized Noninferiority Trial. *Journal of Medical Internet Research*.

Yudo Kasuo. (2014). 慢性疲労症候群患者の日常生活困難度調査事業 . 参照先:
慢性疲労症候群患者の日常生活困難度調査事業 :
<https://www.mhlw.go.jp/file/06-Seisakujouhou-10900000-Kenkoukyoku/0000104377.pdf>