

## REFERENCES

1. Becker BF. Towards the physiological function of uric acid. *Free Radic Biol Med.* 1993;14(6):615-31.
2. Smith EU, Díaz-Torné C, Perez-Ruiz F, March LM. Epidemiology of gout: an update. *Best Pract Res Clin Rheumatol.* 2010;24(6):811-27.
3. Hakoda M. Recent trends in hyperuricemia and gout in Japan. *Japan Med Assoc J.* 2012;55(4):319-23.
4. Perez-Ruiz F, Dalbeth N, Bardin T. A review of uric acid, crystal deposition disease, and gout. *Adv Ther.* 2015;32(1):31-41.
5. Tomita M, Mizuno S, Yamanaka H, Hosoda Y, Sakuma K, Matuoka Y, et al. Does hyperuricemia affect mortality? A prospective cohort study of Japanese male workers. *J Epidemiol.* 2000;10(6):403-9.
6. Grayson PC, Kim SY, LaValley M, Choi HK. Hyperuricemia and incident hypertension: a systematic review and meta-analysis. *Arthritis Care Res (Hoboken).* 2011;63(1):102-10.
7. Baker JF, Schumacher HR, Krishnan E. Serum uric acid level and risk for peripheral arterial disease: analysis of data from the multiple risk factor intervention trial. *Angiology.* 2007;58(4):450-7.

8. Sluijs I, Beulens JW, van der AD, Spijkerman AM, Schulze MB, van der Schouw YT.  
Plasma uric acid is associated with increased risk of type 2 diabetes independent of diet and metabolic risk factors. *J Nutr.* 2013;143(1):80-5.
9. Ben-Dov IZ, Kark JD. Serum uric acid is a GFR-independent long-term predictor of acute and chronic renal insufficiency: the Jerusalem Lipid Research Clinic cohort study. *Nephrol Dial Transplant.* 2011;26(8):2558-66.
10. Obermayr RP, Temml C, Gutjahr G, Knechtelsdorfer M, Oberbauer R, Klauser-Braun R. Elevated uric acid increases the risk for kidney disease. *J Am Soc Nephrol.* 2008;19(12):2407-13.
11. Major TJ, Topless RK, Dalbeth N, Merriman TR. Evaluation of the diet wide contribution to serum urate levels: meta-analysis of population based cohorts. *BMJ.* 2018;363:k3951.
12. Towiwat P, Li ZG. The association of vitamin C, alcohol, coffee, tea, milk and yogurt with uric acid and gout. *Int J Rheum Dis.* 2015;18(5):495-501.
13. Kushiyaama A, Tanaka K, Hara S, Kawazu S. Linking uric acid metabolism to diabetic complications. *World J Diabetes.* 2014;5(6):787-95.

14. Faller J, Fox IH. Ethanol-induced hyperuricemia: evidence for increased urate production by activation of adenine nucleotide turnover. *N Engl J Med*. 1982;307(26):1598-602.
15. Lee SJ, Terkeltaub RA, Kavanaugh A. Recent developments in diet and gout. *Curr Opin Rheumatol*. 2006;18(2):193-8.
16. Choi HK, Mount DB, Reginato AM. Pathogenesis of gout. *Ann Intern Med*. 2005;143(7):499-516.
17. Choi HK, Liu S, Curhan G. Intake of purine-rich foods, protein, and dairy products and relationship to serum levels of uric acid: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum*. 2005;52(1):283-9.
18. Choi HK, Curhan G. Beer, liquor, and wine consumption and serum uric acid level: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum*. 2004;51(6):1023-9.
19. Sugie T, Imatou T, Miyazaki M, Une H. The effect of alcoholic beverage type on hyperuricemia in Japanese male office workers. *J Epidemiol*. 2005;15(2):41-7.
20. Nakamura K, Sakurai M, Miura K, Morikawa Y, Yoshita K, Ishizaki M, et al. Alcohol intake and the risk of hyperuricaemia: a 6-year prospective study in Japanese men. *Nutr Metab Cardiovasc Dis*. 2012;22(11):989-96.

21. Yokose C, McCormick N, Rai SK, Lu N, Curhan G, Schwarzfuchs D, et al. Effects of Low-Fat, Mediterranean, or Low-Carbohydrate Weight Loss Diets on Serum Urate and Cardiometabolic Risk Factors: A Secondary Analysis of the Dietary Intervention Randomized Controlled Trial (DIRECT). *Diabetes Care*. 2020;43(11):2812-20.
22. Dessein PH, Shipton EA, Stanwix AE, Joffe BI, Ramokgadi J. Beneficial effects of weight loss associated with moderate calorie/carbohydrate restriction, and increased proportional intake of protein and unsaturated fat on serum urate and lipoprotein levels in gout: a pilot study. *Ann Rheum Dis*. 2000;59(7):539-43.
23. Gibson T, Rodgers AV, Simmonds HA, Toseland P. Beer drinking and its effect on uric acid. *Br J Rheumatol*. 1984;23(3):203-9.
24. Maxwell S, Cruickshank A, Thorpe G. Red wine and antioxidant activity in serum. *Lancet*. 1994;344(8916):193-4.
25. Nishioka K, Sumida T, Iwatani M, Kusumoto A, Ishikura Y, Hatanaka H, et al. Influence of moderate drinking on purine and carbohydrate metabolism. *Alcohol Clin Exp Res*. 2002;26(8 Suppl):20s-5s.
26. Kottgen A, Albrecht E, Teumer A, Vitart V, Krumsiek J, Hundertmark C, et al. Genome-wide association analyses identify 18 new loci associated with serum urate concentrations. *Nat Genet*. 2013;45(2):145-54.

27. Yokose C, McCormick N, Choi HK. The role of diet in hyperuricemia and gout.  
Curr Opin Rheumatol. 2021;33(2):135-44.