

引用文献

- Adewuya, A. O., Ola, B. A., Aloba, O. O., & Mapayi, B. M. (2006). Anxiety disorders among nigerian women in late pregnancy: A controlled study. *Archives of Women's Mental Health*, 9(6), 325–328.
- 新井洋子. (2010). 産後うつの予防的看護介入プログラムの介入効果の検討. *母性衛生*, 51(1), 144–152.
- Association Psychiatric Association, 高橋三郎, 大野裕 監訳, 染矢俊幸, 神庭重信, 尾崎紀夫, 三村將, 村井俊哉 訳, & 日本精神神経学会. (2014). DSM-5 精神疾患の診断・統計マニュアル / American Psychiatric Association [編]. 医学書院.
- Austin, M.-P., Frilingos, M., Lumly, J., Hadz-pavlovic, D., Roncolato, W., Acland, S., Saint, K., Segal, N., Parker, G. (2008). Brief antenatal cognitive behavior therapy group intervention for the prevention of postnatal depression and anxiety:A randomized controlled trial. *Journal of Affective Disorders* 105;35–44
- Bayrampour, H., McDonald, S., & Tough, S. (2015). Risk factors of transient and persistent anxiety during pregnancy. *Midwifery*, 31(6), 582–589.
- Beck, A. T., & Haigh, E. A. (2014). Advances in cognitive theory and therapy: The generic cognitive model. *Annual Review of Clinical Psychology*, 10, 1–24.
- Beck, C. (2001). Predictors of postpartum depression: An update. *Nursing Research*, 50(5), 275–285.
- Biaggi, A., Conroy, S., Pawlby, S., & Pariante, C. M. (2016). Identifying the women at risk of antenatal anxiety and depression: A systematic review. *Journal of Affective Disorders*, 191, 62–77.
- Bobevski, I., Rowe, H., Clarke, D. M., McKenzie, D. P., & Fisher, J. (2015). Early postnatal demoralisation among primiparous women in the community: Measurement, prevalence and associated factors. *BMC Pregnancy and Childbirth*, 15, 259–3.
- Brockington, I. F., Macdonald, E., & Wainscott, G. (2006). Anxiety, obsessions

- and morbid preoccupations in pregnancy and the puerperium. *Archives of Women's Mental Health*, 9(5), 253–263.
- Chabrol, H., Teissedre, F., Saint-Jean, M., Teisseyre, N., Rogé, B., & Mullet, E. (2002). Prevention and treatment of post-partum depression: a controlled randomized study on women at risk. *Psychological Medicine*, 32(6), 1039–1047.
- Cluxton-Keller, F., & Bruce, M. L. (2018). Clinical effectiveness of family therapeutic interventions in the prevention and treatment of perinatal depression: A systematic review and meta-analysis. *PLoS One*, 13(6), e0198730.
- Coelho, H. F., Murray, L., Royal-Lawson, M., & Cooper, P. J. (2011). Antenatal anxiety disorder as a predictor of postnatal depression: A longitudinal study. *Journal of Affective Disorders*, 129(1-3), 348–353.
- Cuijpers, P., van Straten, A., Andersson, G., & van Oppen, P. (2008). Psychotherapy for depression in adults: A meta-analysis of comparative outcome studies. *Journal of Consulting and Clinical Psychology*, 76(6), 909–922.
- Cuijpers, P., Weitz, E., Karyotaki, E., Garber, J., & Andersson, G. (2015). The effects of psychological treatment of maternal depression on children and parental functioning: A meta-analysis. *European Child & Adolescent Psychiatry*, 24(2), 237–245.
- Dennis, C. L., & Dowswell, T. (2013). Psychosocial and psychological interventions for preventing postpartum depression. *The Cochrane Database of Systematic Reviews*, (2):CD001134.
- Fairbrother, N., Janssen, P., Antony, M. M., Tucker, E., & Young, A. H. (2016). Perinatal anxiety disorder prevalence and incidence. *Journal of Affective Disorders*, 200, 148–155.
- Fairbrother, N., Janssen, P., Antony, M. M., Tucker, E., & Young, A. H. (2016). Perinatal anxiety disorder prevalence and incidence. *Journal of Affective Disorders*, 200, 148–155.

- Farrell, T., Reagu, S., Mohan, S., Elmidany, R., Qaddoura, F., Ahmed, E. E., Corbett, G., Lindow, S., Abuyaqoub, S. M., & Alabdulla, M. A. (2020). The impact of the COVID-19 pandemic on the perinatal mental health of women. *Journal of Perinatal Medicine*, 48(9), 971–976.
- Geren, A., Birge, Ö, Bakır, M. S., Sakıncı, M., & Sanhal, C. Y. (2021). Does time change the anxiety and depression scores for pregnant women on Covid-19 pandemic? *The Journal of Obstetrics and Gynaecology Research*, 47(10), 3516–3523.
- Goodman, S. H., Rouse, M. H., Connell, A. M., Broth, M. R., Hall, C. M., & Heyward, D. (2011). Maternal depression and child psychopathology: A meta-analytic review. *Clinical Child and Family Psychology Review*, 14(1), 1–27.
- Grace, S. L., Evindar, A., & Stewart, D. E. (2003). The effect of postpartum depression on child cognitive development and behavior: A review and critical analysis of the literature. *Archives of Women's Mental Health*, 6(4), 263–274.
- Gavin, N. I., Gaynes, B. N., Lohr, K. N., Meltzer-Brody, S., Gartlehner, G., & Swinson, T. (2005). Perinatal depression: a systematic review of prevalence and incidence. *Obstetrics and Gynecology*, 106(5 Pt 1), 1071–1083.
- Green, S. M., Donegan, E., McCabe, R. E., Streiner, D. L., Agako, A., & Frey, B. N. (2020). Cognitive behavioral therapy for perinatal anxiety: A randomized controlled trial. *The Australian and New Zealand Journal of Psychiatry*, , 4867419898528.
- Grote, N. K., Bridge, J. A., Gavin, A. R., Melville, J. L., Iyengar, S., & Katon, W. J. (2010). A meta-analysis of depression during pregnancy and the risk of preterm birth, low birth weight, and intrauterine growth restriction. *Archives of General Psychiatry*, 67(10), 1012–1024.
- Hahn-Holbrook, J., Cornwell-Hinrichs, T., & Anaya, I. (2018). Economic and

health predictors of national postpartum depression prevalence: A systematic review, meta-analysis, and meta-regression of 291 studies from 56 countries. *Frontiers in Psychiatry*, 8, 248.

橋本直也. (2021). With COVID-19 時代のシミュレーション教育・オンライン診療 9 オンラインでの健康医療相談（遠隔健康医療相談）. 小児科, 62(4), 371-381.

服部律子, 武田順子, 名和文香, 布原佳奈, 松山久美, 田中真理, 小森春佳, 澤田麻衣子. (2019). 助産師が認識する「気になる母子」への対応と他機関との連携に関する研究. *岐阜県立看護大学紀要*, 19 (1). 63-79.

Health Quality Ontario. (2019). Ontario Health Technology Assessment Series. Health Quality Ontario, 19(6).

堀越勝. (2015) ケアする人の対話スキル ABCD. 日本看護協会

House J. S. (1973). Occupational Stress and Coronary Heart Disease. *Journal of Health & Social Behavior*, 15, 12-27.

Hunt C, Issakidis C & Andrews G. (2002). DSM-IV generalized anxiety disorder in the Australian National Survey of Mental Health and Well-Being. *Psychol Med*. 32. 649-659

五十嵐良雄(2013). あなたも名医！「うつ状態」を知る・診る. 日本医事新法社
Jonathan Evans, Jenny Ingram, Roslyn law, Hazal Taylor, Debbie Johnson, Joel Glynn, Becky hopley, David Kessler, Jeff Round, Jenny Ford, Iryna Culpin and heather O' Mahen. (2021). Interpersonal counselling versus perinatal-specific cognitive behavioural therapy for women with depression during pregnancy offered in routine psychological treatment services:a phase II ramdomised trial. *BMC Psychiatry*. 21:504, 7-11.

我部山キヨ子, 武谷雄二, 中林正雄. (2014), 他. 助産診断・技術学Ⅱ [2] 分娩期・産褥期 (第5版) . 助産診断・技術学 / 横尾京子編 ; 横尾京子執筆 (第5版). 東京: 医学書院

金岡緑. (2011). 育児に対する自己効力感尺度 (parenting self-efficacy scale : PSE 尺度) の開発とその信頼性・妥当性の検討. 小児保健研究, 70(1), 27-38.

川上憲人. (2007). こころの健康についての疫学調査に関する研究 総合研究報告書. 平成16~18年度厚生労働科学研究費補助金 こころの健康科学的研究事業

- Kim, D. R., Sockol, L. E., Sammel, M. D., Kelly, C., Moseley, M., & Epperson, C. N. (2013). Elevated risk of adverse obstetric outcomes in pregnant women with depression. *Archives of Women's Mental Health*, 16(6), 475–482.
- Kitamura, T., Yoshida, K., Okano, T., Kinoshita, K., Hayashi, M., Toyoda, N., Ito, M., Kudo, N., Tada, K., Kanazawa, K., Sakumoto, K., Satoh, S., Furukawa, T., & Nakano, H. (2006). Multicentre prospective study of perinatal depression in Japan: incidence and correlates of antenatal and postnatal depression. *Archives of Women's Mental Health*, 9(3), 121–130.
- Kubota, C., Okada, T., Aleksic, B., Nakamura, Y., Kunimoto, S., Morikawa, M., Shiino, T., Tamaji, A., Ohoka, H., Banno, N., Morita, T., Murase, S., Goto, S., Kanai, A., Masuda, T., Ando, M., & Ozaki, N. (2014). Factor structure of the Japanese version of the Edinburgh Postnatal Depression Scale in the postpartum period. *PloS One*, 9(8), e103941.
- 桑原さやか, 新井洋子. (2019). 産後うつの予防的支援についての検討—妊娠期のピアサポートを活用して—. *母性衛生*, 60(2), 362–369.
- 厚生労働省研究事業. (2014). 妊産婦のメンタルヘルスの実態把握及び介入方法に関する研究. <https://research-er.jp/projects/view/886080>
- 厚生労働省. (2020). 産前・産後サポート事業ガイドライン 産後ケア事業ガイドライン.
- 国立成育医療研究センター. (2018). 人口動態統計（死亡・出生・死産）から見る妊娠中・産後の死亡の現状 <https://www.ncchd.go.jp/press/2018/maternal-deaths.html> 【閲覧日 ; 2021.11.13】
- 国立精神・神経医療研究センター認知行動療法センター. (2019). 平成31年度認知行動療法ベーシック研修
- Lancaster, C. A., Gold, K. J., Flynn, H. A., Yoo, H., Marcus, S. M., & Davis, M. M. (2010). Risk factors for depressive symptoms during pregnancy: A systematic review. *American Journal of Obstetrics and Gynecology*, 202(1), 5–14.
- Leach, L. S., Poyser, C., & Fairweather-Schmidt, K. (2017). Maternal perinatal anxiety: A review of prevalence and correlates. *Clinical Psychologist*,

21(1), 4–19.

- Li, C., Sun, X., Li, Q., Sun, Q., Wu, B., & Duan, D. (2020). Role of psychotherapy on antenatal depression, anxiety, and maternal quality of life: A meta-analysis. *Medicine*, 99(27), e20947.
- Lieberman, K., Le, H. & Perry, D. F. (2014). A systematic review of perinatal depression interventions for adolescent mothers. *Journal of Adolescence*, 37(8), 1227e1235.
- López-Morales, H., Del Valle, M. V., Canet-Juric, L., Andrés, M. L., Galli, J. I., Poó, F., & Urquijo, S. (2021). Mental health of pregnant women during the COVID-19 pandemic: A longitudinal study. *Psychiatry Research*, 295, 113567.
- Loughnan, S. A., Sie, A., Hobbs, M. J., Joubert, A. E., Smith, J., Haskelberg, H., Mahoney, A. E. J., Kladnitski, N., Holt, C. J., Milgrom, J., Austin, M. P., Andrews, G., & Newby, J. M. (2019). A randomized controlled trial of 'MUMentum Pregnancy': Internet-delivered cognitive behavioral therapy program for antenatal anxiety and depression. *Journal of Affective Disorders*, 243, 381–390.
- Maguire, P. N., Clark, G. I., & Wootton, B. M. (2018). The efficacy of cognitive behavior therapy for the treatment of perinatal anxiety symptoms: A preliminary meta-analysis. *Journal of Anxiety Disorders*, 60, 26–34.
- Matthey, S., Barnett, B., Howie, P., & Kavanagh, D. J. (2003). Diagnosing postpartum depression in mothers and fathers: Whatever happened to anxiety? *Journal of Affective Disorders*, 74(2), 139–147.
- Matthey, S., Kavanagh, D. J., Howie, P., Barnett, B., & Charles, M. (2004). Prevention of postnatal distress or depression: An evaluation of an intervention at preparation for parenthood classes. *Journal of Affective Disorders*, 79(1–3), 113–126.
- 間中麻衣子, 河副みゆき, 佐々木綾子. (2019). 産後うつ予防のためのリーフレットを用いた個別指導－妊娠後期から産褥 1 か月までの初産婦の EPDS の変化－.

60(2), 348–354.

Milgrom, J., Gemmill, A. W., Bilszta, J. L., Hayes, B., Barnett, B., Brooks, J., Erickson, J., Ellwood, D., & Buist, A. (2008). Antenatal risk factors for postnatal depression: a large prospective study. *Journal of Affective Disorders*, 108(1–2), 147–157.

Milgrom, J., Gemmill, A. W., Danaher, B. G., Holt, C., Holt, C. J., Seeley, J. R., Tyler, M. S., Ross, J., & Erickson, J. (2016). Internet Cognitive Behavioral Therapy for Women With Postnatal depression: A Randomized Controlled Trial of mumMoodMooster. *Journal of Medical internet research*, 18(3). e54, 1–18.

Morrell, C. J., Sutcliffe, P., Booth, A., Stevens, J., Scope, A., Stevenson, M., Harvey, R., Bessey, A., Cantrell, A., Dennis, C. L., Ren, S., Ragonesi, M., Barkham, M., Churchill, D., Henshaw, C., Newstead, J., Slade, P., Spiby, H., & Stewart-Brown, S. (2016). A systematic review, evidence synthesis and meta-analysis of quantitative and qualitative studies evaluating the clinical effectiveness, the cost-effectiveness, safety and acceptability of interventions to prevent postnatal depression. *Health Technology Assessment* (Winchester, England), 20(37), 1–414.

村松公美子. (2010). Gad-7 日本語版の妥当性・有用性の検討. *心身医学*, 50(6), 592.

National institute for Health and Care Excellence. (2014). Antenatal and postnatal mental health: clinical management and service guidance Clinical guideline [CG192], <https://www.nice.org.uk/guidance/cg192/chapter/1-Recommendations#providing-interventions-in-pregnancy-and-the-postnatal-period-2>

日本看護協会. (2020). 2019 年度改定 助産実践能力熟練段階 (クリニカルラダー) Clinical ladder Competencies for midwifery Practice CLoCMiP 活用ガイド

日本ラクテーション・コンサルタント協会. (2015). 母乳育児支援スタンダード / 日本ラクテーション・コンサルタント協会編 (第2版). 東京: 医学書院

日本産婦人科医会 (2017). 妊産婦メンタルヘルスケアマニュアル～産後ケアへの切れ目のない支援に向けて～；公益社団法人日本産婦人科医会

- 日本産科婦人科学会, 日本産婦人科医会, (2020) . 産婦人科診療ガイドライン—産科
編 2020, https://www.jsog.or.jp/activity/pdf/g1_sanka_2020.pdf
- 日本助産師会 (2021). 助産師の声明/コア・コンピテンシー2021
- 野口聰一, 菊井敬子, 中田高公. (2017). 無痛分娩による周産期予後と合併症の検討.
現代産婦人科, 66(2), 261-266.
- Oser, M., Wallace, M. L., Solano, F., & Szigethy, E. M. (2019). Guided digital cognitive behavioral program for anxiety in primary care: Propensity-matched controlled trial. *JMIR Mental Health*, 6(4), e11981.
- Petzoldt, J. (2018). Systematic review on maternal depression versus anxiety in relation to excessive infant crying: It is all about the timing. *Archives of Women's Mental Health*, 21(1), 15-30.
- Radesky, J. S., Zuckerman, B., Silverstein, M., Rivara, F. P., Barr, M., Taylor, J. A., Lengua, L. J., & Barr, R. G. (2013). Inconsolable infant crying and maternal postpartum depressive symptoms. *Pediatrics*, 131(6), 1857.
- Robertson, E., Grace, S., Wallington, T., & Stewart, D. E. (2004). Antenatal risk factors for postpartum depression: A synthesis of recent literature. *General Hospital Psychiatry*, 26(4), 289-295.
- Robichaud, M., Talbot, F., Titov, N., Dear, B. F., Hadjistavropoulos, H. D., Hadjistavropoulos, T., & Jbilou, J. (2020). Facilitating access to iCBT: a randomized controlled trial assessing a translated version of an empirically validated program using a minimally monitored delivery model. *Behavioural and Cognitive Psychotherapy*, 48(2), 185-202.
- Russell, E. J., Fawcett, J. M., & Mazmanian, D. (2013). Risk of obsessive-compulsive disorder in pregnant and postpartum women: A meta-analysis. *The Journal of Clinical Psychiatry*, 74(4), 377-385.
- Satoh, A., Kitamiya, C., Kudoh, H., Watanabe, M., Menzawa, K., & Sasaki, H. (2009). Factors associated with late post-partum depression in Japan. *Japan Journal of Nursing Science : JJNS*, 6(1), 27-36.
- 佐藤喜根子, 佐藤祥子. (2010) .妊娠期からの継続した心理的支援が周産期女性の不

安・抑うつに及ぼす効果. 母性衛生, 51(1), 215-225.

社会保障審議会, 子ども虐待による死亡事例等の検証結果等について. (2019). 社会保障審議会児童部会児童虐待等要保護事例の検証に関する専門委員会, 第 17 次報告. <https://www.mhlw.go.jp/content/11900000/000822359.pdf> 【閲覧日 : 2021. 11. 13】

周産期メンタルヘルス学会. (2017). 周産期メンタルヘルスコンセンサスガイド.
<http://pmhguideline.com/>

末継奏子, 岩元澄子. (2015). 産後の抑うつ予防を目的とした妊娠期からの介入の検討. 久留米大学心理学研究, 14, 25-35.

Scope, A., Leaviss, J., Kalenthaler, E., Parry, G., Sutcliffe, P., Bradburn, M., & Cantrell, A. (2013). Is group cognitive behaviour therapy for postnatal depression evidence-based practice? A systematic review. BMC Psychiatry, 13, 321-321.

Shirabe, R., Okuhara, T., Yokota, R., Okada, H., Goto, E., & Kiuchi, T. (2021). Changes in Anxiety and Stress Among Pregnant Women During the COVID-19 Pandemic: Content Analysis of a Japanese Social Question-and-Answer Website. JMIR Pediatrics and Parenting, 4(3), e27733.

Sim, J., & Lewis, M. (2012). The size of a pilot study for a clinical trial should be calculated in relation to considerations of precision and efficiency. Journal of Clinical Epidemiology, 65(3), 301-308.

Sockol, L. E. (2015). A systematic review of the efficacy of cognitive behavioral therapy for treating and preventing perinatal depression. Journal of Affective Disorders, 177, 7-21.

Sockol, L. E., Epperson, C. N., & Barber, J. P. (2011). A meta-analysis of treatments for perinatal depression. Clinical Psychology Review, 31(5), 839-849.

Sockol, L. E., Epperson, C. N., & Barber, J. P. (2013). Preventing postpartum depression: A meta-analytic review. Clinical Psychology Review, 33(8), 1205-1217.

Spitzer, R. L., Kroenke, K., Williams, J. B., & Lowe, B. (2006). A brief

measure for assessing generalized anxiety disorder: The GAD-7. Archives of Internal Medicine, 166(10), 1092–1097.

立森久照, 伊藤弘人. (1999). 日本語版Client Satisfaction Questionnaire の 8 項目版の信頼性および妥当性の検討, 41(7), 711–717.

Keita Tokumitsu, Noriko Sugawara, kazushi Marumo, Toshihito Suzuki, Kazutaka Shimoda & Noriko Yasui-Furukori. (2020). Prevalence of perinatal depression among Japanese women:a meta-analysis. Annals of General Psychiatry. 19:41, 3–18.

寺坂多栄子, 岡山久代. (2015). 妊娠末期・産褥早期における産後うつ予防の保健指導の効果, 56(1), 87–94.

東京都保健福祉局. (2018) 平成 30 年人口統計.

https://www.fukushihoken.metro.tokyo.lg.jp/smph/kiban/chosa_tokei/jinkodotaitokei/index.html 【最終閲覧日 2021.11.13】

Vesga-Lopez, O., Blanco, C., Keyes, K., Olfson, M., Grant, B. F., & Hasin, D. S. (2008). Psychiatric disorders in pregnant and postpartum women in the united states. Archives of General Psychiatry, 65(7), 805–815.

渡邊一代, 石井佳世子, 石田久江, 太田操, 後藤あや. (2019). 産後うつ病予防を目的とした妊娠期からの“夫婦の共感性を高めるセッション”の試行：対象者の共感性と精神健康度とセッション評価. 日本健康学会誌. 85 (2). 80–89

Wenzel, A., Haugen, E. N., Jackson, L. C., & Brendle, J. R. (2005). Anxiety symptoms and disorders at eight weeks postpartum. Journal of Anxiety Disorders, 19(3), 295–311.

Wilkinson, E. L., O'Mahen, H. A., Fearon, P., Halligan, S., King, D. X., Greenfield, G., Dunkley-Bent, J., Erickson, J., Milgrom, J., & Ramchandani, P. G. (2016). Adapting and testing a brief intervention to reduce maternal anxiety during pregnancy (ACORN): study protocol for a randomised controlled trial. Trials, 17, 156–8.

Wootton, B. M., Bragdon, L. B., Steinman, S. A., & Tolin, D. F. (2015). Three-year outcomes of adults with anxiety and related disorders following cognitive-behavioral therapy in a non-research clinical setting. Journal

of Anxiety Disorders, 31, 28–31.

Yamashita, H., Yoshida, K., Nakano, H., & Tashiro, N. (2000). Postnatal depression in Japanese women. Detecting the early onset of postnatal depression by closely monitoring the postpartum mood. Journal of Affective Disorders, 58(2), 145–154.

Yasuma, N., Narita, Z., Sasaki, N., Obikane, E., Sekiya, J., Inagawa, T., Nakajima, A., Tamada, Y., Yamazaki, R., Matunaga, A., Saitoh, T., Watanabe, K., Imaura, K., Kawakami, N. & Nishi, D. (2020). Antenatal psychological intervention for universal prevention of antenatal and postnatal depression: A systematic review and meta-analysis. Journal of Affective Disorders, 273, 231–239.

米山万里枝. (2008). 妊娠中の不安と産後うつとの関連性. 周産期医学, 38(11), 1466–1476.

湯舟邦子. (2015). 妊娠初期、中期、末期から産後 1 カ月までの抑うつ状態のスクリーニングの検討. 昭和学士会雑誌, 75(4), 465–473.