

参考・引用文献

- Alba B. M. (2021). CE: Postpartum Depression: A Nurse's Guide. *The American journal of nursing*, 121(7), 32-43.
- American College of Obstetricians and Gynecologists' Committee on Practice Bulletins-Obstetrics. (2021). Anemia in Pregnancy: ACOG Practice Bulletin, Number 233. *Obstetrics and gynecology*, 138(2), e55-e64.
- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*, Washington, DC: American Psychiatric Association.
- Azami, M., Badfar, G., Khalighi, Z., Qasemi, P., Shohani, M., Soleymani, A., & Abbasalizadeh, S. (2019). The association between anemia and postpartum depression: A systematic review and meta-analysis. *Caspian journal of internal medicine*, 10(2), 115-124.
- Beard, J. L., Hendricks, M. K., Perez, E. M., Murray-Kolb, L. E., Berg, A., Vernon-Feagans, L., Irlam, J., Isaacs, W., Sive, A., & Tomlinson, M. (2005). Maternal iron deficiency anemia affects postpartum emotions and cognition. *The Journal of nutrition*, 135(2), 267-272.
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of general psychiatry*, 4, 561-571.
- Blumfield, M. L., Hure, A. J., Macdonald-Wicks, L., Smith, R., & Collins, C. E. (2013). A systematic review and meta-analysis of micronutrient intakes during pregnancy in developed countries. *Nutrition reviews*, 71(2), 118-132.
- Brummelte, S., & Galea, L. A. (2016). Postpartum depression: Etiology, treatment and consequences for maternal care. *Hormones and behavior*, 77, 153-166.
- Camaschella C. (2015). Iron-deficiency anemia. *The New England journal of medicine*, 372(19), 1832-1843.

- Catania, G., Bell, C., Ottonelli, S., Marchetti, M., Bryce, J., Grossi, A., & Costantini, M. (2013). Cancer-related fatigue in Italian cancer patients: validation of the Italian version of the Brief Fatigue Inventory (BFI). *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer*, 21(2), 413-419.
- Corwin, E. J., Murray-Kolb, L. E., & Beard, J. L. (2003). Low hemoglobin level is a risk factor for postpartum depression. *The Journal of nutrition*, 133(12), 4139-4142.
- Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *The British journal of psychiatry : the journal of mental science*, 150, 782-786.
- Garcia-Esteve, L., Ascaso, C., Ojuel, J., & Navarro, P. (2003). Validation of the Edinburgh Postnatal Depression Scale (EPDS) in Spanish mothers. *Journal of affective disorders*, 75(1), 71-76.
- Goshtasebi, A., Alizadeh, M., & Gandevani, S. B. (2013). Association between maternal anaemia and postpartum depression in an urban sample of pregnant women in Iran. *Journal of health, population, and nutrition*, 31(3), 398-402.
- Hameed, S., Naser, I. A., Al Ghussein, M. A., & Ellulu, M. S. (2021). Is iron deficiency a risk factor for postpartum depression? A case-control study in the Gaza Strip, Palestine. *Public health nutrition*, 1-8.
- Hamilton M. (1960). A rating scale for depression. *Journal of neurology, neurosurgery, and psychiatry*, 23(1), 56-62.
- 平野まどか, 渡邊憲和, 成味恵, 出井麗, 深瀬実加, 高橋可菜子, 山内敬子, 堤誠司, 永瀬智. (2021). エジンバラ産後うつ病評価票を用いた、合併症妊娠を含めた妊娠関連項目と周産期うつ病の関連の解析. *山形医学*, 39(1), 10-15.

- Hodapp V. (1989). Theoretische und empirische Analysen der Struktur der STAI-Angstskalen [Theoretical and empirical analyses of the structure of the State Trait Anxiety Inventory anxiety scales]. *Archiv für Psychologie*, 141(1), 27-47.
- 市川ゆかり, 黒田緑. (2008). 産後うつ病に関連する要因の分析. *母性衛星*, 49(2), 336-346.
- Kang, S. Y., Kim, H. B., & Sunwoo, S. (2020). Association between anemia and maternal depression: A systematic review and meta-analysis. *Journal of psychiatric research*, 122, 88-96.
- 菊池紗耶, 小林 奈津子, 木村 涼子, 富田 博秋. (2021). 産科医のための周産期メンタルヘルスケア入門. *産婦人科の実際*, 70(6), 635-640.
- Kim, S. Y., Park, J. E., Lee, Y. J., Seo, H. J., Sheen, S. S., Hahn, S., Jang, B. H., & Son, H. J. (2013). Testing a tool for assessing the risk of bias for nonrandomized studies showed moderate reliability and promising validity. *Journal of clinical epidemiology*, 66(4), 408-414.
- McGuinness, L. A., & Higgins, J. (2021). Risk-of-bias Visualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Research synthesis methods*, 12(1), 55-61.
- Miller BJ, Murray L, Beckmann MM, Kent T, Macfarlane B. Dietary supplements for preventing postnatal depression. *Cochrane Database Syst Rev*. 2013 Oct 24;(10):CD009104.
- Moraes, G. P., Lorenzo, L., Pontes, G. A., Montenegro, M. C., & Cantilino, A. (2017). Screening and diagnosing postpartum depression: when and how?. *Trends in psychiatry and psychotherapy*, 39(1), 54-61.
- Moretti, D., Goede, J. S., Zeder, C., Jiskra, M., Chatzinakou, V., Tjalsma, H., Melse-Boonstra, A., Brittenham, G., Swinkels, D. W., & Zimmermann, M. B. (2015). Oral iron supplements increase hepcidin and decrease iron absorption from daily

or twice-daily doses in iron-depleted young women.

Nguyen, P. H., DiGirolamo, A. M., Gonzalez-Casanova, I., Pham, H., Hao, W., Nguyen, H., Truong, T. V., Nguyen, S., Harding, K. B., Reinhart, G. A., Martorell, R., & Ramakrishnan, U. (2017). Impact of preconceptional micronutrient supplementation on maternal mental health during pregnancy and postpartum: results from a randomized controlled trial in Vietnam. *BMC women's health*, 17(1), 44.

日本人の食事摂取基準策定検討会. (2020). 日本人の食事摂取基準(2020年版) 策定検討会報告書. 厚生労働省.

日本助産学会ガイドライン委員会. (2020). エビデンスに基づく助産ガイドライン—妊娠期・分娩期・産褥期 2020. 日本助産師会.

日本産科婦人科学会. (2020). 産婦人科診療ガイドライン—産科編 2020. 日本産科婦人科学会.

小川浩平, 前田裕斗, 左合治彦. (2021). 産後うつに貧血が関係? 国立成育医療研究センターが新知見を報告. *ペリネイタルケア*, 40(9) 906-910.

Peña-Rosas, J. P., De-Regil, L. M., Garcia-Casal, M. N., & Dowswell, T. (2015). Daily oral iron supplementation during pregnancy. *The Cochrane database of systematic reviews*, (7), CD004736.

Radloff LS. (1977). The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas*, 385-401.

Rodriguez-Cabezas, L., & Clark, C. (2018). Psychiatric Emergencies in Pregnancy and Postpartum. *Clinical obstetrics and gynecology*, 61(3), 615-627.

Ruiz de Viñaspre-Hernández, R., Gea-Caballero, V., Juárez-Vela, R., & Iruzubieta-Barragán, F. J. (2021). The definition, screening, and treatment of postpartum anemia: A systematic review of guidelines. *Birth (Berkeley, Calif.)*, 48(1), 14-25.

- Shea, B. J., Reeves, B. C., Wells, G., Thuku, M., Hamel, C., Moran, J., Moher, D., Tugwell, P., Welch, V., Kristjansson, E., & Henry, D. A. (2017). AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ (Clinical research ed.)*, 358, j4008.
- Sheikh, M., Hantoushzadeh, S., Shariat, M., Farahani, Z., & Ebrahimasab, O. (2017). The efficacy of early iron supplementation on postpartum depression, a randomized double-blind placebo-controlled trial. *European journal of nutrition*, 56(2), 901-908.
- Smets, E. M., Garssen, B., Bonke, B., & De Haes, J. C. (1995). The Multidimensional Fatigue Inventory (MFI) psychometric qualities of an instrument to assess fatigue. *Journal of psychosomatic research*, 39(3), 315-325.
- Soler, J., Pérez-Sola, V., Puigdemont, D., Pérez-Blanco, J., Figueres, M., & Alvarez, E. (1997). Estudio de validación del Center for Epidemiologic Studies-Depression (CES-D) en una población española de pacientes con trastornos afectivos [Validation study of the Center for Epidemiological Studies-Depression of a Spanish population of patients with affective disorders]. *Actas luso-espanolas de neurologia, psiquiatria y ciencias afines*, 25(4), 243-249.
- Sterne, J., Savović, J., Page, M. J., Elbers, R. G., Blencowe, N. S., Boutron, I., Cates, C. J., Cheng, H. Y., Corbett, M. S., Eldridge, S. M., Emberson, J. R., Hernán, M. A., Hopewell, S., Hróbjartsson, A., Junqueira, D. R., Jüni, P., Kirkham, J. J., Lasserson, T., Li, T., McAleenan, A., ... Higgins, J. (2019). RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ (Clinical research ed.)*, 366, 14898.
- Stewart, D. E., & Vigod, S. N. (2019). Postpartum Depression: Pathophysiology, Treatment, and Emerging Therapeutics. *Annual review of medicine*, 70, 183-196.

- Tokumitsu, K., Sugawara, N., Maruo, K., Suzuki, T., Shimoda, K., & Yasui-Furukori, N. (2020). Prevalence of perinatal depression among Japanese women: a meta-analysis. *Annals of general psychiatry*, 19, 41.
- Tran, T. D., Tran, T., La, B., Lee, D., Rosenthal, D., & Fisher, J. (2011). Screening for perinatal common mental disorders in women in the north of Vietnam: a comparison of three psychometric instruments. *Journal of affective disorders*, 133(1-2), 281-293.
- Vitale, S. G., Fiore, M., La Rosa, V. L., Rapisarda, A., Mazza, G., Paratore, M., Commodari, E., & Caruso, S. (2021). Liposomal ferric pyrophosphate and ascorbic acid supplementation in pregnant women with iron deficiency anaemia: haematochemical, obstetric, neonatal and psychological outcomes in a prospective observational study. *International journal of food sciences and nutrition*, 1-9.
- Ware, J. E., Jr, & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Medical care*, 30(6), 473-483.
- Woody, C. A., Ferrari, A. J., Siskind, D. J., Whiteford, H. A., & Harris, M. G. (2017). A systematic review and meta-regression of the prevalence and incidence of perinatal depression. *Journal of affective disorders*, 219, 86-92.
- World Health Organization. (1992). *The Prevalence of Anaemia in Women: A Tabulation of Available Information*. 2nd ed. Geneva: World Health Organization.
- World Health Organization. (2006). *Iron and folate supplementation: integrated management of pregnancy and childbirth (IMPAC)*. Standards for maternal and neonatal Geneva: Department of Making Pregnancy Safer, World Health Organization.
- World Health Organization. (2008). *Worldwide prevalence of anaemia 1993-2005* : WHO

Global database on anaemia. / Edited by Bruno de Benoist, Erin McLean, Ines Egli and Mary Cogswell. World Health Organization.

World Health Organization. (2016). WHO recommendations on antenatal care for a positive pregnancy experience. Geneva: World Health Organization.