

## 引用・参考文献

Akechi, T., Momino, K., Katsuki, F., Yamashita, H., Sugiura, H., Yoshimoto, N., et al. (2021). Brief collaborative care intervention to reduce perceived unmet needs in highly distressed breast cancer patients: Randomized controlled trial. *Japanese Journal of Clinical Oncology*, 51(2), 244-251. doi:10.1093/jjco/hyaa166

明智龍男.(2002).がん患者の精神症状の診断と治療.がん看護,7(6).475-481.

Akechi, T., Okuyama, T., Imoto, S., Yamawaki, S., & Uchitomi, Y. (2001). Biomedical and psychosocial determinants of psychiatric morbidity among postoperative ambulatory breast cancer patients Springer Science and Business Media LLC. doi:10.1023/a:1010661530585

Akizuki N, Yamawaki S, Akechi T, Nakano T, Uchitomi Y.(2005). Development of an Impact Thermometer for use in combination with the Distress Thermometer as a brief screening tool for adjustment disorders and/or major depression in cancer patients. *Journal of Pain & Symptom Management*. 29(1),91-99. doi:10.1016/j.jpainsymman.2004.04.016

浅野憲一,清水栄司.(2015).集団認知行動療法の特徴と新しい展開.臨床精神医学,44(8).1067-1073.

飛鳥井望.(2015).VII 精神症状の評価法:18 [トラウマ体験評価] 改訂出来事インパクト尺度 (IES-R). 山内俊雄, 鹿島晴雄(総編集), 精神・心理機能評価ハンドブック (p.262).中山書店.

Baker, F., Denniston, M., Smith, T., & West, M. M. (2005). Adult cancer survivors: How are they faring? *Cancer*, 104(S11), 2565-2576. doi:10.1002/cncr.21488

Barbara B. Germino., Merle H. Mishel., Jamie Crandell., Laura Porter., Diane Blyler., Coretta Jenerette., Karen M. Gil.(2013).Outcomes of an Uncertainty Management Intervention in Younger African and Caucasian Breast Cancer Survivors. *Oncology Nursing*, 40(1),82-92.

Beltman, M. W., Voshaar, R. C. O., & Speckens, A. E. (2010). Cognitive-behavioural therapy for depression in people with a somatic disease: Meta-analysis of randomised controlled trials. *British Journal of Psychiatry*, 197(1), 11-19. doi:10.1192/bjp.bp.109.064675

- Burgess, C., Cornelius, V., Love, S., Graham, J., Richards, M., & Ramirez, A. (2005). Depression and anxiety in women with early breast cancer: Five year observational cohort study. *Bmj*, 330(7493), 702-705. doi:10.1136/bmj.38343.670868.D3
- Chang, Y., Hu, W., & Chang, Y. (2020). Cognitive-behavioral therapy to alleviate treatment-induced menopausal symptoms in women with breast cancer: A systematic review. *Cancer Nursing*, 44(5), 411-418. doi:10.1097/NCC.0000000000000827
- Calhoun, L., Tedeschi, R. (2006/2014) 宅香菜子, 清水研(監訳). 第1章 心的外傷後成長の基礎 発展的枠組み. 心的外傷後成長ハンドブック: 耐え難い体験が人の心にもたらすもの (pp.4-8). 医学書院.
- Folkman, S. (2012/2016). はじめに. Moorey, S., Greer, S. (著), 鈴木伸一(監訳). がん患者の認知行動療法: メンタルケアと生活支援のための実践ガイド. (pp. IV-V.). 北大路書房.
- Getu, M. A., Chen, C., Panpan, W., Mboineki, J. F., Dhakal, K., & Du, R. (2020). The effect of cognitive behavioral therapy on the quality of life of breast cancer patients: A systematic review and meta-analysis of randomized controlled trials. *Quality of Life Research*, 30(2), 367-384. doi:10.1007/s11136-020-02665-5
- Götze, H., Friedrich, M., Taubenheim, S., Dietz, A., Lordick, F., & Mehnert, A. (2019). Depression and anxiety in long-term survivors 5 and 10 years after cancer diagnosis. *Supportive Care in Cancer*, 28(1), 211-220. doi:10.1007/s00520-019-04805-1
- Hamidian, P., Rezaee, N., Shakiba, M., & Navidian, A. (2018). The effect of Cognitive-Emotional training on post-traumatic growth in women with breast cancer in middle east. *Journal of Clinical Psychology in Medical Settings*, 26(1), 25-32. doi:10.1007/s10880-018-9561-z
- 平井啓. (2014). がん患者に対する問題解決療法アプローチの適用. *行動医学研究*, 20(1), 7-11.
- Huntley, A. L., Araya, R., & Salisbury, C. (2012). Group psychological therapies for depression in the community: Systematic review and meta-analysis. *British Journal of Psychiatry*, 200(3), 184-190. doi:10.1192/bjp.bp.111.092049
- Imai, F., Momino, K., Katsuki, F., Horikoshi, M., Furukawa, T. A., Kondo, N., et al. (2019). Smartphone problem-solving therapy to reduce fear of cancer recurrence among breast cancer survivors: An open single-arm pilot study. *Japanese Journal of Clinical Oncology*, 49(6), 537-544.

doi:10.1093/jjco/hyz005

Koch, L., Bertram, H., Eberle, A., Holleczeck, B., Schmid-Höpfner, S., Waldmann, A., et al. (2014). Fear of recurrence in long-term breast cancer survivors-still an issue. results on prevalence, determinants, and the association with quality of life and depression from the cancer survivorship-a multi-regional population-based study. *Psycho-Oncology* (Chichester, England), 23(5), 547-554. doi:10.1002/pon.3452

厚生労働省 (2018) .がん対策推進基本計画. <https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/000183313.html> (検索日 : 2022 年 1 月 4 日)

国立がん研究センター がん情報サービス. (2021). 院内がん登録生存率集計.<https://ganjoho.jp/public/cancer/breast/patients.html> (検索日 : 2021 年 12 月 30 日)

Komatsu, H., Hayashi, N., Suzuki, K., Yagasaki, K., Iioka, Y., Neumann, J., et al. (2012). Guided self-help for prevention of depression and anxiety in women with breast cancer. *ISRN Nursing*, 2012, 716367-8. doi:10.5402/2012/716367

小山敦子.(2014).特集:サイコオンコロジー サイコオンコロジー総論.*Japanese Society of Psychosomatic Medicine*,54,12-19.

熊野宏昭. (2012). 第 2 章 新世代の認知行動療法に共通する者. *新世代の認知行動療法*(pp.21-32).日本評論社.

白石裕子(編).(2014).第 1 章 認知行動療法を理解し看護実践に活用していくためのポイント. *看護のための認知行動療法*(pp.17-20).金剛出版.

Lengacher, C. A., Reich, R. R., Post-White, J., Moscoso, M., Shelton, M. M., Barta, M., et al. (2011). Mindfulness based stress reduction in post-treatment breast cancer patients: An examination of symptoms and symptom clusters. *Journal of Behavioral Medicine*, 35(1), 86-94. doi:10.1007/s10865-011-9346-4

松永美希,鈴木伸一,岡本泰昌,木下亜紀子,吉村晋平,山脇成人.(2007).うつ病に対する集団認知行動療法の展望.*精神科治療学*,22(9).1081-1091.

松沼晶子.,二渡玉江.(2020).ピアサポートを受けたがん患者の思いに関する研究内容の分析.

桐生大学紀要.第 31 号,99-107.

Momino, K., Mitsunori, M., Yamashita, H., Toyama, T., Sugiura, H., Yoshimoto, N., et al. (2017). Collaborative care intervention for the perceived care needs of women with breast cancer undergoing adjuvant therapy after surgery: A feasibility study. *Japanese Journal of Clinical Oncology*, 47(3), 213-220. doi:10.1093/jjco/hyw189

大谷恭平,内富庸介.(2010).総説「他領域からのトピックス」 がん患者の心理と心のケア.日本耳鼻咽喉科学会会報.113,45-52.

Park, S., Sato, Y., Takita, Y., Tamura, N., Ninomiya, A., Kosugi, T., et al. (2020). Mindfulness-based cognitive therapy for psychological distress, fear of cancer recurrence, fatigue, spiritual well-being, and quality of life in patients with breast Cancer—A randomized controlled trial. *Journal of Pain and Symptom Management*, 60(2), 381-389. doi:10.1016/j.jpainsymman.2020.02.017

Park, S., Sado, M., Fujisawa, D., Sato, Y., Takeuchi, M., Ninomiya, A., et al. (2018). Mindfulness-based cognitive therapy for japanese breast cancer patients—a feasibility study. *Japanese Journal of Clinical Oncology*, 48(1), 68-74. doi:10.1093/jjco/hyx156

Ramos, C., Leal, I., & Tedeschi, R. G. (2016). Protocol for the psychotherapeutic group intervention for facilitating posttraumatic growth in nonmetastatic breast cancer patients. *BMC women's Health*, 16(1), 22. doi:10.1186/s12905-016-0302-x

佐渡充洋.(2019a).マインドフルネス療法の基本的なアプローチ.精神医学,61(6),627-635.

佐渡充洋.(2019b).マインドフルネス認知療法-構造,エビデンス,そして効果機序-.認知行動療法の進歩：研究と臨床の最前線.分子精神医学,19(1),28-33.

Segal ZV, Williams JMG, Teasdale JD.(2002) Mindfulness-based Cognitive Therapy for Depression: A New Approach to Preventing Relapse. New York: Guilford Press.

竹内恵美,小川祐子,原沙彩,鈴木伸一.(2016).乳がんサバイバーにおける再発不安の構成要因の検討-再発の心配および対処行動の観点から-.行動医学研究,22(1),9-17.

Tedeschi, R. G., & Calhoun, L. G. (1996). The posttraumatic growth inventory: Measuring the positive legacy of trauma. *Journal of Traumatic Stress*, 9(3), 455-471. doi:10.1002/jts.2490090305

Turkington, D., Kingdon, D., Athod, R. S., Hammond, K., Pelton, J., Mehta, R. (2006). Outcomes of an effectiveness trial of cognitive behavioural intervention by mental health nurses in schizophrenia. *British Journal of Psychiatry*, 189, 36-40. doi : 10.1192/bjp.bp.105.010884