

引用文献

- Adams J.,Frawley J.,Steel A., et al.(2015). Use of pharmacological and non-pharmacological labour pain management techniques and their relationship to maternal and infant birth outcomes: Examination of a nationally representative sample of 1,835 pregnant women *Midwifery*,31(4),pp.458-463.
- Andersson L.,Poromaa I.,Wulff M.,et al.(2004).Implications of Antenatal Depression and Anxiety for Obstetric Outcome. *American College of Obstetricians and gynecologists*,104(3),pp.467-476.
- Antonakou A.,& Papoutsis.(2016). The Effect of Epidural Analgesia on the Delivery Outcome of Induced Labour: A Retrospective Case Series. *Obstetrics and Gynecology International*,pp.1-5.
- Areskog,B.,& Uddenberg,N.(1984).Postnatal emotional balance in women with and without antenatal fear of childbirth.*Journal of psychosomatic research*,28(3), pp.213-220.
- Blondel B.,Breart G., Mazaubrun C., et al.(1997).The perinatal situation in France.Trends between 1981 and 1995.*Gynecol Obstet Biol Reprod*,26(8),pp.770-780.
- Blondel B.,Supernant K.,Mazaubrun., et al.(2006).Trends in perinatal health in metropolitan France between 1995 and 2003: results from the National perinatal surveys. *Gynecol Obstet Biol Reprod*,35(4).pp.373-387.
- Brenda A.,Bucklin M.,Hawkins,J.(2005).Obstetric anesthesia workforce survey: twenty year update. *Anesthesiology*,103,pp.645-653.
- Brixval C.,Thygesen L.,Axelsen S., et al.(2016).Effect of antenatal education in small classes versus standard auditorium-based lectures on use of pain relief during labour and of obstetric interventions: results from the randomized NEWBORN trial.*BMJ Open*,6,pp.1-10.
- Carlsson M.,Ziegert K.,Nissen E.(2015).The relationship between childbirth self-efficacy and aspects of well-being,birth interventions and birth outcomes.*Midwifery*,31,pp.1000-1007.
- Carolan M.,Davey M.,Biro M., et al.(2011).Older maternal age and intervention in labor: A population-based study comparing older and younger first-time mothers in

- Victoria,Australia.*BIRTH*,38:1,pp.24-28.
- Carvalho B.,Wang P.,Cohen S.(2006).A survey of labor patient-controlled epidural anesthesia practice in California hospitals.*International journal of obstetric anesthesia*,15(3),pp.217-222.
- Cheng,Y.W.,Shaffer,B.L., et al.(2014).Second stage of labor and epidural use: a larger effect than previously suggested.*Obstet Gynecol*, 123, pp.527-535.
- Corwin E ., & Bozoky I.(2002).Fatigue as predictor of postpartum depression.*Journal of Obstetric,Gynecologic & Neonatal Nursing*,31(4),pp.436-443.
- Ding T.,Wang D.,Qu Y., et al.(2014).Epidural Labor Analgesia is associated with a decreased risk of postpartum: a prospective cohort study.*Anesthesia and analgesia*,119(2),pp.383-392.
- Fenwick J.,Toohill J.,Gamble J., et al.(2015).Effects of a midwife psycho-education intervention to reduce childbirth fear on women's birth outcomes and postpartum psychological wellbeing.*BMC Pregnancy and Childbirth*,15:284,pp.1-8.
- Ferrer M.,Jordana M., Meseguer C., et al.(2016).Comparative study analyzing women's childbirth satisfaction and obstetric outcomes across two different models of maternity care.*BMJ Open*,6,pp.1-10.
- Firoozeh M.,Fattaneh.,Lhadigeh M., et al.(2016). Respect for woman's decision-making in spontaneous birth: A thematic synthesis study.*Iranian Journal of Nursing and Midwifery Research*,21(5).pp.449-457.
- Gau A.,Chang C.,Tian S., et al.(2011). Effects of birth ball exercise on pain and self-efficacy during childbirth: A randomised controlled trial in Taiwan.*Midwifery*,27(6),pp.293-300.
- Gibbins J.,& Tomson M.(2001).Women's expectations and experiences of childbirth.*Midwifery*,17,pp302-313.
- Gomenz A.,Martinez O.,Torrecillas J.(2015).Retrospective study of the association between epidural analgesia during labour and complications for the newborn.*Midwifery*,31,pp.613-616.
- Green J.(1993).Expectation and experiences of pain in labor: findings from a large prospective study,*BIRTH*,20(2),pp.65-72.

- Greenow C.,Nassar N.,Torvaldsen., et al.(2010).Assisting informed decision making for labour analgesia: a randomized controlled trial of a decision aid for labour analgesia versus a pamphlet.*BMC Pregnancy and Childbirth*,10:15,pp.1-13.
- Hauck,Y.,& Fenwick J.(2007).The influence of childbirth expectations on Western Australian women's perceptions of their birth experience.*Midwifery*, 23,pp.235-247.
- Hasegawa Junichi.,Antonio Farina.,Hasegawa Yuko., et al. (2013).Effects of epidural analgesia on labor length,instrumental delivery, and neonatal short-term outcome.*Journal of Anesthesia*, 27(1),pp.43-47.
- 花沢成一(1977).妊娠の時期による不安の変動,母性心理学研究VI.心身医,17,51.
- Halls K.(2008).Maternal satisfaction regarding anaesthetic services during childbirth.British *Journal of Midwifery*,16(5),296-301.
- Halpern,S.H.,Leighton,B.L., et al.(1998).Effect of epidural vs parenteral opioid analgesia on the progress of labor:a meta-analysis.*JAMA*,280,pp.2105-2110.
- Hannah G.,Lesley M.,Homer C.(2010).The novice birthing: theorizing first-time mothers' experiences of birth at home and in hospital in Australia.*Midwifery*,26,pp.53-63.
- Harkins J.,Carvalho.,Evers A., et al.(2010).Survey of the factors associated with a woman's choice to have an epidural for labor analgesia.*Anesthesiology Research and Practice*,pp.1-8.
- Hatem M.,Sandall J.,Devane D., et al.(2016).Midwife-led versus other models of care for childbearing women. *Cochrane Database of Systematic Reviews*,4.
- Hawkins J.,Gibbs C.,Orleans M., et al.(1997).Obstetric anesthesia work force survey,1981versus1992.*Anesthesiology*,87(1),pp.135-143.
- Hitashu Bhatt,Sunil Pandya.,Geeta Kolar.(2014).The impact of labour epidural analgesia on the childbirth expectation and experience at a tertiary care center in Southern India.*Journal of Clinical and Diagnostic Research*,8(3),pp.73-76.
- Hodnett,E.D.(1987).The labour agency scale: Psychometric properties of an instrument measuring control during childbirth.*Research in Nursing & Health*, 10(5),pp.301-310.
- Hodnett,E.D.(2002).Pain and women's satisfaction with the experience of childbirth:a

- systematic review. *American Journal of Obstetrics and Gynecology*, 186(5), pp.160-172.
- Humburg J.(2008).Postartum urinary retention-without clinical inmac?. *Ther Umsc* h, 65(11), pp.681-685.
- Humenick.R.N.(1981).Mastery: The Key to Cildbirth Satisfaction? A Review. *BIRTH*, 8(2), pp79-83.
- 飯村直子, 檜木野裕美, 二宮啓子(2002). Wong-Bakerのフェイススケールの日本における妥当性と信頼性. *日本小児看護会誌*, 11, pp.21-27.
- 石井秀宗(2007).統計分析のここが知りたいー保健・看護・心理・教育系のまとめ方ー. 文光堂, pp. 60-62.
- Ip W., Tang C., Goggins W.(2009).An education intervention to improve women's ability to cope with childbirth. *Journal of Clinical Nursing*, 18, pp.2125-2135.
- 伊藤伸大(2016).当院産婦人科外来における無痛分娩アンケート調査(会議録). *日本周産期・新生児医学会雑誌*, 52(2), pp.662.
- 岩ヶ谷愛, 交野好子, 石津みゑ子(2005).産婦の分娩に対する満足度に関する研究-理想・予想と現実との関連性-. *日本健康科学学会誌*, 21(2), pp.269-278.
- Jones L., Othman M., Dowswell T., et al.(2012).Pain management for women in labour: an overview of systematic reviews. *Cochrane Database of Systematic Reviews*, 3.
- Kendall K., Cong Z., Hale T.(2015).Birth Interventions Related to Lower Rates of Exclusive Breastfeeding and Increased Risk of Postpartum Depression in a Large Sample. *clinical Lactation*, 6(3), pp87-97.
- 厚生労働省(2015).平成 27 年人口動態統計月報年計(概数)の概況
<http://www.mhlw.go.jp/toukei/saikin/hw/jinkou/geppo/nengai15/dl/gaikyou27.pdf>
 [アクセス日 : 2016 年 12 月 1 日]
- Kringeland T., Kjersti A., Moller A.(2010).What characterizes women who want to give birth as naturally as possible without painkillers or intervention?, *Sexual & Reproductive Healthcare*, 1, pp.21-26.
- 久保田陽子, 伊田昌功, 伊藤宏一, 加藤浩志他(2014).硬膜外麻酔による無痛分娩が分娩および新生児に与える影響について. *産婦人科の進歩*, 66(3), pp.157-264.
- Lally J., Murtagh M., Macphail S., et al.(2008).More in hope than expectation: a sy

- stematic review of women's expectations and experience of pain relief in labour. *BMC Medicine*, 6(7), pp.1-10.
- Lally J., Thomson R., Macphail S., et al. (2014). Pain relief in labour: a qualitative study to determine how to support women to make decisions about pain relief in labour. *BMC Pregnancy & Childbirth*, 14(6), pp.1-10.
- Leap N., Sandall J., Buckland S., et al. (2010). Journey to confidence: women's experiences of pain in labour and relational continuity of care. *Journal of Midwifery & Women's Health*, 55(3), pp.234-242.
- Lene A., Torset B., Bo K. (2015). What is the effect of regular group exercise on maternal psychological outcomes and common pregnancy complaints? An assessor blinded RCT. *Midwifery*, 32, pp.81-86.
- Lowe, N.K. (2002). The nature of labor pain. *Am J Obstet Gynecol*, 186(5), pp.16-24.
- Meades, R., & Ayers, S. (2011). Anxiety measures validated in perinatal populations: A systematic review. *Journal of Affective Disorders*, 133(1-2), pp. 1-15.
- 水尾智佐子, 塩野悦子 (2013). 妊娠期に無痛分娩を選んだ女性の出産に至るまでの体験. *日本助産学会誌*, 27(2), 257-266.
- 毛利多恵子 (1993). 産婦の身体拘束が陣痛体験に及ぼす影響. 聖路加看護大学大学院看護学研究科修士論文, pp.1-61.
- 中里克治, 水口公信 (1982). 新しい不安尺度 STAI 日本語版の作成. *心身医*, 22, pp.107-112.
- 日本産婦人科学会編 (2003). *産婦人科用語集・用語解説集*. 改訂新版, 201.
- 日本産業衛生学会産業疲労研究会 (1970). 自覚症しらべの使い方.
- Retrieved from <http://square.umin.ac.jp/of/> [アクセス日 : 2015年4月10日]
- 岡田景子, 村山朋子 (2007). 妊婦が無痛分娩を選択する理由. *第38回母性看護*, pp.14-16.
- 奥富俊之 (2002). 無痛分娩に対する妊婦の意識. *ペインクリニック*, 23(2), pp.154-158.
- Oliver, R.L. (1980). A cognitive model of the antecedents and consequences of satisfaction decisions. *Journal of Marketing Research*, 17, pp.460-469.
- Pettersson F., Hellgren C., Nyberg F., et al. (2016). Depressed mood, anxiety, and the use of labor analgesia. *Womens Ment Health*, 19, pp.11-16.
- Read, G.D. (1949). Observations on a series of labours. *The Lancet*, pp.721-726.
- Redshaw M., & Henderson J. (2015). Safely delivered: a national survey of women's a

- national survey of maternity care 2014. *National Perinatal epidemiology Unit*, pp.1-83.
- Rouhe H., Salmela K., Gissler., et al. (2011). Mental health problems common in women with fear of childbirth. *An international Journal of Obstetrics and Gynaecology*, pp.1104-1111.
- Ryding, E.L., & Wijima, K. (1998). Fear of childbirth during pregnancy may increase the risk of emergency caesarean section. *Acta Obstetrica et Gynecologica Scandinavica*, 77, pp.542-547.
- 鈴木恵子(1998).産痛教育が産婦の産痛イメージ及び産痛体験へ与える影響.研究実績報告書.
- 佐々木和子(2011).硬膜外麻酔使用による無痛分娩:産後1年経過した母親が語るその体験を聞いて. *助産雑誌*, 65(5), pp.418-423.
- Somuah, M., Smyth, R.M., Jones, L. (2011). Epidural versus non-epidural or no analgesia in labour. *The Cochrane Library*, pp.1-89.
- Spielberger C., Gorsuch L., Lushene, R.E.. (1970). Manual for the state trait anxiety inventory (Self-evaluation questionnaire). *Consulting Psychologists*, pp.2-24.
- 菅裕佳子, 岩間洋亮, 佐々木宏輔 (2013). 最近2年間における当院の硬膜外無痛分娩2646例に関する考察. *分娩と麻酔*, 95, pp.18-23.
- 須郷慶信, 筒井淳奈, 渡辺典芳 (2010). 症例から学ぶハイリスク妊娠、無痛分娩後の尿閉. *産科と産婦人科*, 77(4), pp.449-451.
- Sun Yi-Chin., Hung Ya-Chi., Chang Y. (2009). Effects of a prenatal yoga programme on the discomforts of pregnancy and maternal childbirth self-efficacy in Taiwan. *midwifery*, 26, pp.31-36.
- 照井克生, 上山博史, 大西佳彦, 他 (2008). 全国の分娩取り扱い施設における麻酔科診療実態調査. 厚生労働科学研究費補助金分担研究報告書, pp.433-468.
- Troy, N.W. (2003). Is the significance of postpartum fatigue being overlooked in the lives of women?. *The American Journal of Maternal Child Nursing*, 28(4), pp.252-257.
- Tsuchiya M., Mori E., Sakajo A., et al. (2016). Age-specific determinants of post-partum fatigue in primiparous women. *Japan Journal of Nursing Science*, 13(1), pp.83-94.
- 梅原永能, 角倉弘行, 他 (2011). 高齢初産婦と無痛分娩. *産婦人科治療*, 103(4), pp.369-395.
- Wong, D., & Baker, C. (1988). Pain in children: Comparison of Assessment Scales. *Pediatric*

Nursing, 14(1), pp.9-17.

Yazdanimehr R., Omid A., Sadat Z., et al. (2016). The effect of mindfulness-integrated cognitive behavior therapy on depression and anxiety among pregnant women: a randomized clinical trial. *Journal of Caring Sciences*, 5(3), pp.195-204.

柳吉桂子(1992). 初産婦の出産経験に対する評価過程. *日本看護科学会誌*, 12(3), pp.212-213.

山崎圭子, 齋藤益子(2012). わが国における産後の疲労感に関する文献検討. *日本母子看護学会誌*, 6(2), pp.31-39.

山崎圭子, 齋藤益子(2014). 「産後の疲労感」の概念分析. *日本母子看護学会誌*, 7(2), pp.1-10.

吉田和枝(2008). 欧米および日本における産痛対応法の比較的研究. *大阪大学大学院人間科学研究科紀要*, 34, pp.269-289.