

References

- von Both, C., Fleßa, S., Makuwani, A., Mpembeni, R., & Jahn, A. (2006). How much time do health services spend on antenatal care? Implications for the introduction of the focused antenatal care model in Tanzania. *BMC Pregnancy & Childbirth*, *6*, 22. doi: 1471-2393-6-22 [pii]
- Carlo, W. A., Wright, L. L., Chomba, E., McClure, E. M., Carlo, M. E., Bann, C. M., . . . Harris, H. (2009). Educational impact of the neonatal resuscitation program in low-risk delivery centers in a developing country. *Journal of Pediatrics*, *154*(4), 504-508.e5. doi:10.1016/j.jpeds.2008.10.005 [doi]
- Carroli, G., Villar, J., Piaggio, G., Khan-Neelofur, D., Gulmezoglu, M., Mugford, M., . . . WHO Antenatal Care Trial Research Group. (2001). WHO systematic review of randomised controlled trials of routine antenatal care. *Lancet (London, England)*, *357*(9268), 1565-1570. doi: S0140-6736(00)04723-1 [pii]
- Dulock, L. H., & Holzemer, L. W. (1993). Substruction: Improving the linkage from theory to method. *Kango Kenkyu*, *26* (5), 455-61. [In Japanese]
- Duysburgh, E., Ye, M., & Williams, A. (2013). Counselling on and women's awareness of pregnancy danger signs in selected rural health facilities in Burkina Faso, Ghana and Tanzania. *Tropical Medicine & International Health*, *18* (12), 1498-1509.
- Duysburgh, E., Ye, M., Williams, A., Massawe, S., Sie, A., Williams, J., . . . Temmerman, M. (2013). Counselling on and women's awareness of pregnancy danger signs in selected rural health facilities in Burkina Faso, Ghana and Tanzania. *Tropical Medicine & International Health: TM & IH*, *18*(12), 1498-1509. doi:10.1111/tmi.12214 [doi]
- Ersdal, H. L., Vossius, C., Bayo, E., Mduma, E., Perlman, J., Lippert, A., & Soreide, E. (2013). A one-day "helping babies breathe" course improves simulated performance but not clinical management of neonates. *Resuscitation*, *84*(10), 1422-1427. doi:10.1016/j.resuscitation.2013.04.005 [doi]
- Evans, C. L., Johnson, P., Bazant, E., Bhatnagar, N., Zgambo, J., & Khamis, A. R. (2014). Competency-based training "helping mothers survive: Bleeding after birth" for providers from central and remote facilities in three countries. *International Journal of Gynaecology & Obstetrics*, *126*(3), 286-290. doi:10.1016/j.ijgo.2014.02.021 [doi]
- Griscti, O., & Jacono, J. (2005). Effectiveness of continuing education programmes in nursing: literature review. *Journal of Advanced Nursing*, *55*, (4), 449-456, doi: 10.1111/j.1365-2648.2006.03940.x

- Hägström, E., Mbusa, E., & Wadensten, B. (2008). Nurses' workplace distress and ethical dilemmas in Tanzanian health care. *Nursing Ethics*, *15* (4), 478-491. doi: 10.1177/0969733008090519
- Harvey, S. A., Jennings, L., Chinyama, M., Masaniga, F., Mulholland, K., & Bell, D. (2008). Improving community health worker use of malaria rapid diagnostic tests in Zambia: Package instructions, a job aid, and job aid-plus training. *Malaria Journal*, *7*, 160-172.
- Jennings, L., Yebadokpo, A. S., Affo, J., & Agbogbe, M. (2010). Antenatal counseling in maternal and newborn care: Use of job aids to improve health worker performance and maternal understanding in Benin. *BMC Pregnancy & Childbirth*, *10*, 75-2393-10-75. doi: 10.1186/1471-2393-10-75 [doi]
- Jennings, L., Yebadokpo, A., Affo, J., & Agbogbe, M. (2015). Use of job aids to improve facility-based postnatal counseling and care in rural Benin. *Maternal & Child Health Journal*, *19*(3), 557-565. doi: 10.1007/s10995-014-1537-5 [doi]
- JHPIEGO. (2004a). *Maternal and neonatal health programme birth preparedness and complication readiness: A matrix of shared responsibilities* (Original BP/CR Matrix Poster published in 2001. English Introductory Text revised in 2004). Maryland, USA: Johns Hopkins University for International Education in Gynecology and Obstetrics.
- JHPIEGO. (2004b). *Monitoring birth preparedness and complication readiness tools and indicators for maternal and newborn health*. Maryland, USA: Johns Hopkins University for International Education in Gynecology and Obstetrics.
- Kabakian-Khasholian, T., Campbell, O., Shediak-Rizkallah, M., & Ghorayeb, F. (2000). Women's experiences of maternity care: Satisfaction or passivity? *Social Science & Medicine* (1982), *51*(1), 103-113. doi: S0277953699004438 [pii]
- Kabakyenga, J. K., Ostergren, P. O., Turyakira, E., & Pettersson, K. O. (2011). Knowledge of obstetric danger signs and birth preparedness practices among women in rural Uganda. *Reproductive Health*, *8*, 33-4755-8-33. doi: 10.1186/1742-4755-8-33 [doi]
- Kahabuka, C., Kvåle, G., Moland, K. M., & Hinderaker, S. G. (2011). Why caretakers bypass primary health care facilities for child care - a case from rural Tanzania. *BMC Health Services Research*, *11*, 315-6963-11-315. doi: 10.1186/1472-6963-11-315 [doi]
- Kearns, A., Hurst, T., Caglia, J., & Langer, A. (2014). Focused antenatal care in Tanzania. Women and Health Initiative & Maternal Health Task Force, Harvard School of Public Health. Retrieved from: <https://cdn2.sph.harvard.edu/wp->

content/uploads/sites/32/2014/09/HSPH-Tanzania5.pdf

- Kedenge, S. V., Kangwana, B. P., Waweru, E. W., Nyandigisi, A. J., Pandit, J., Brooker, S. J., . . . Goodman, C. A. (2013). Understanding the impact of subsidizing artemisinin-based combination therapies (ACTs) in the retail sector--results from focus group discussions in rural Kenya. *PloS One*, 8(1), e54371. doi:10.1371/journal.pone.0054371 [doi]
- Kim, Y. M., Davila, C., Tellez, C., & Kols, A. (2007). Evaluation of the world health organization's family planning decision-making tool: Improving health communication in Nicaragua. *Patient Education & Counseling*, 66(2), 235-242. doi: S0738-3991(06)00426-5 [pii]
- Knebel E. (2000). The use of manual job aids by health care providers: What do we know? (Issue Paper). Bethesda, MD: QAP for USAID. Retrieved from: http://www.urcchs.com/uploads/resourcefiles/JobAids_RationaleforUseandExperie nctoDate.pdf
- Leshabari, S., Koniz-Booher, P., Burkhalter, B., Hoffman, M., & Jennings, L. (2007). *Testing a PMTCT infant-feeding counseling program in Tanzania*. U.S. Agency for International Development. Retrieved from: <https://www.usaidassist.org/sites/assist/files/orrtztestingjobaids.pdf>
- Lassi, Z. S., Haider, B. A., & Bhutta, Z. A. (2010). Community-based intervention packages for reducing maternal and neonatal morbidity and mortality and improving neonatal outcomes (Review). *The Cochrane Library*, 11. doi: 10.1002/14651858.CD007754.pub2
- Magoma, M., Requejo, J., Campbell, O. M., Cousens, S., & Filippi, V. (2010). High ANC coverage and low skilled attendance in a rural Tanzanian district: A case for implementing a birth plan intervention. *BMC Pregnancy & Childbirth*, 10, 13-2393-10-13. doi: 10.1186/1471-2393-10-13 [doi]
- Manongi, R. N., Nasuwa, F. R., Mwangi, R., Reyburn, H., Poulsen, A., & Chandler, C. I. (2009). Conflicting priorities: Evaluation of an intervention to improve nurse-parent relationships on a Tanzanian paediatric ward. *Human Resources for Health*, 7, 50-4491-7-50. doi: 10.1186/1478-4491-7-50 [doi]
- Mutiso, S. M., Qureshi, Z., & Kinuthia, J. (2008). Birth preparedness among antenatal clients. *East African Medical Journal*, 85 (6). 275-283.
- National Bureau of Statistics (NBS) and ICF Marco. (2011). *Tanzania demographic and health survey 2010*. Dar es Salaam, Tanzania: NBS and ICF Marco.
- National Bureau of Statistics (NBS), Macro International. (2007). *Service provision assessment survey 2006*. Dar es Salaam, Tanzania: National Bureau of Statistics.

- National Bureau of Statistics (NBS), Ministry of Finance and Office of Chief Government Statistician President's Office, Finance, Economy and Development Planning (2013a). *Population distribution by age and sex*. Dar es Salaam, Zanzibar, Tanzania: National Bureau of Statistics.
- National Bureau of Statistics (NBS), Ministry of Finance and Office of Chief Government Statistician President's Office, Finance, Economy and Development Planning (2013b). *2012 Population and housing census*. Dar es Salaam, Zanzibar, Tanzania: National Bureau of Statistics.
- National Institute for Health and Care Excellence. (2010). *Pregnancy and Complex Social Factors: a model for Service Provision for Pregnant Women with Complex Social Factors*. Manchester, UK: National Institute for Health and Care Excellence.
- Nelissen, E., Ersdal, H., Mduma, E., Evjen-Olsen, B., Broerse, J., van Roosmalen, J., & Stekelenburg, J. (2015). Helping mothers survive bleeding after birth: Retention of knowledge, skills, and confidence nine months after obstetric simulation-based training. *BMC Pregnancy & Childbirth*, *15*, 190-015-0612-2. doi: 10.1186/s12884-015-0612-2 [doi]
- Nyamtema, A. S., Bartsch-de Jong, A., Urassa, D. P., Hagen, J. P., & van Roosmalen, J. (2012). The quality of antenatal care in rural Tanzania: What is behind the number of visits? *BMC Pregnancy & Childbirth*, *12*, 70-2393-12-70. doi: 10.1186/1471-2393-12-70 [doi]
- Nyamtema, A. S., Urassa, D. P., & Roosmalen, J. V. (2011). Maternal health interventions in resource limited countries: A systematic review of packages, impacts and factors for change. *BMC Pregnancy & Childbirth*, *11*. doi: 10.1186/1471-2393-11-30i
- Okereke, E., Aradeon, S., Akerele, A., Tanko, M., Yisa, I., & Obonyo, B. (2013). Knowledge of safe motherhood among women in rural communities in northern Nigeria: Implications for maternal mortality reduction. *Reproductive Health*, *10*(1), 57-4755-10-57. doi: 10.1186/1742-4755-10-57 [doi]
- Pelto, G. H., Santos, I., Goncalves, H., Victora, C., Martines, J., & Habicht, J. P. (2004). Nutrition counseling training changes physician behavior and improves caregiver knowledge acquisition. *Journal of Nutrition*, *134*(2), 357-362.
- Pembe, A. B., Carlstedt, A., Urassa, D. P., Lindmark, G., Nystrom, L., & Darj, E. (2010). Quality of antenatal care in rural Tanzania: Counselling on pregnancy danger signs. *BMC Pregnancy & Childbirth*, *10*. doi: 10.1186/1471-2393-10-35
- Pembe, A. B., Urassa, D. P., Carlstedt, A., Lindmark, G., Nyström, L., & Darj, E.

(2009). Rural

Tanzanian women's awareness of danger signs of obstetric complications. *BMC Pregnancy & birth*, 9. doi: 10.1186/1471-2393-9-12

Prost, A., Colbourn, T., Seward, N., Azad, K., Coomarasamy, A., Copas, A., ...

Costello, A. (2013). Women's groups practicing participatory learning and action to improve maternal and newborn health in low-resource settings: A systematic review and meta-analysis. *Lancet*, 18. 381 (9879): 1736-1746. doi: 10.1016/S0140-6736 (13) 60685-6

Rajabu, M. (2015). *Official hospital statistics, annual report 2014*. Tanga, Tanzania: The Anglican Diocese of Tanga, Hospital Management Team.

Sarker, M., Schmid, G., Larsson, E., Kirenga, S., De Allegri, M., Neuhann, F., . . .

Muller, O. (2010). Quality of antenatal care in rural southern Tanzania: A reality check. *BMC Research Notes*, 3, 209-0500-3-209. doi: 10.1186/1756-0500-3-209 [doi]

Shimpuku, Y., & Madeni, F. E. (2014). *Nne na Tano* [booklet]. Japan, St. Luke's International University. [In Kiswahili and English]

Shrestha, S., Petrini, M., & Turale, S. (2013). Newborn care in Nepal: The effects of an educational intervention on nurses' knowledge and practice. *International Nursing Review*, 60(2), 205-211. doi:10.1111/inr.12017 [doi]

Soubeiga, D., Gauvin, L., Hatem, M. A., & Johri, M. (2014). Birth Preparedness and Complication Readiness (BPCR) interventions to reduce maternal and neonatal mortality in developing countries: Systematic review and meta-analysis. *BMC Pregnancy & Childbirth*, 14, 129, doi: 10.1186/1471-2393-14-129

Srivastava, A., Avan, B. I., Rajbangshi, P., & Bhattacharyya, S. (2015). Determinants of women's satisfaction with maternal health care: A review of literature from developing countries. *BMC Pregnancy & Childbirth*, 15, 97-015-0525-0. doi: 10.1186/s12884-015-0525-0 [doi]

Tanzania Ministry of Health and Social Welfare. (2007). *Focused Antenatal Care Malaria and Syphilis in Pregnancy. National Advocacy Guide*. Dar es Salaam, Tanzania: Ministry of Health and Social Welfare.

Tanzania Ministry of Health and Social Welfare. (2011). *A performance Audit on the Monitoring, Evaluations and Budget Allocation for Maternal Health Care Activities in Tanzania*. Dar es Salaam, Tanzania: Ministry of Health and Social Welfare.

Thaddeus, S., & Maine, D. (1994). Too far to walk: Maternal mortality in context. *Social Science & Medicine* 38: 1091-1110. Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed/8042057>

- World Health Organization. (2002). *WHO antenatal care randomized trial: Manual for the implementation of the new model*. Geneva Switzerland: World Health Organization.
- World Health Organization. (2006). *Pregnancy, childbirth, postpartum and newborn care: A guide for essential practice*. Geneva Switzerland: World Health Organization.
- World Health Organization. (2013). *Counselling for maternal and newborn health care: A handbook for building skills*. Geneva Switzerland: World Health Organization.
- World Health Organization. (2014). *Focused antenatal care in Tanzania*. Geneva Switzerland: World Health Organization.
- World Health Organization. (2014). *Trends in maternal mortality: 1990 to 2013*. Geneva Switzerland: World Health Organization.
- Willis-Shattuck, M., Bidwell, P., Thomas, S., Wyness, L, Blaauw, D., & Ditlopo, P. (2008). Motivation and retention of health workers in developing countries: a systematic review. *BMC Pregnancy & Childbirth*, 8, 247, doi: 10.1186/1472-6963-8/247
- Zaman, S., Ashraf, R. N., & Martines, J. (2008). Training in complementary feeding counselling of healthcare workers and its influence on maternal behaviours and child growth: A cluster-randomized controlled trial in Lahore, Pakistan. *Journal of Health, Population, & Nutrition*, 26(2), 210-222.